



















# CHEMISTRY SIMPLIFIED.





# CHEMISTRY SIMPLIFIED.

A COURSE OF LECTURES ON THE NON-METALS,  
BASED UPON THE NATURAL EVOLUTION  
OF CHEMISTRY.

DESIGNED PRIMARILY FOR ENGINEERS.

BY

GEORGE AUGUSTUS KOENIG, Ph. D., A. M., E. M.,

PROFESSOR OF CHEMISTRY, MICHIGAN COLLEGE OF MINES, HOUGHTON, MICHIGAN.

ILLUSTRATED BY ONE HUNDRED AND THREE ORIGINAL DRAWINGS.



PHILADELPHIA :  
HENRY CAREY BAIRD & CO ,  
INDUSTRIAL PUBLISHERS, BOOKSELLERS AND IMPORTERS,  
No. 810 WALNUT STREET.  
1906,

Q.D. 31  
K 6

GENERAL

COPYRIGHT, BY  
GEORGE AUGUSTUS KOENIG,  
1905.





## PREFACE.

---

IN these lectures to mature beginners in Chemistry, the fundamental idea has been followed to unroll before the student the knowable nature of bodies as an ever-growing and spreading picture, and not as a finished work handed down by the great masters. When I say mature beginners, I mean that in my estimation chemistry should not be taught at all to boys and girls before their full mental growth has been attained, fully aware of my isolated position in the present evolution of schools. Only the mature mind follows with growing interest the unfolding of such a picture, and absorbs it as a living thing.

In following this fundamental idea, the usual systematic classification had to be abandoned. It will be seen that the beginning is made with bodies of familiar acquaintance such as the common metals, but these metals are not postulated as elements or simple bodies; they are merely objects for experimentation in allowing the equally familiar bodies of air and water to act upon them under the familiar impulse of heat. Noting the changes thus wrought, the mind questions and seeks answers. Answers come by carefully laid experiments, but always in

(v)

such a way that no agent of unfamiliar nature is called in to aid. Such a scheme is essentially historical, for the successive generations of chemists have been working in exactly this way. They saw and raised questions. Their answers, being premature in so far as they brought into play unknown agents, were often wrong, had to be set aside and much modified by subsequent investigations. Thus also in these lectures, questions are raised, but not answered at once, although perfectly well known, because the student's knowledge has not reached the required fullness. In the chapters on green vitriol and on common salt, as well as on potash, the reader will find the application of the fundamental idea fully elaborated.

Generalizations from the experiments are always drawn, though I have avoided the term law. Theorizing upon molecules and the structure of molecules, ions and electrons I have omitted altogether. No beginning student can be capable of drawing inferences for himself concerning these matters. They should only be brought before the student at the end of his school work, if he intends following chemistry. To bring them before proposing engineers, seems unnecessary, if not unwise, and these lectures are delivered to engineering candidates, more especially mining engineers and metallurgists. To them the theories can be of no help. They want to know the practical consequences which must follow from the presence of certain material conditions. They must be trained to inquire, and de-



duce from given conditions. The chief tendency of the lecture course is to evoke in the student the constant thought of Why? together with the love for the experiment. The course extends over seven months and one-half, in three lectures a week. It is supplemented by laboratory work of eight hours per week in which some fifty-odd experiments, selected from the lecture experiments, are performed by the student. The chemistry of the metals in conjunction with qualitative analysis comprises the second year's work of equal time-extension, but only two lectures a week.

The illustrations are made with the chief aim of engineering simplicity. Thanks are duly given to each and every chemist who has given a laborious life in contributions with which to build up the chemistry of to-day.

THE AUTHOR.

HOUGHTON, MICHIGAN, *November 15, 1905.*



# CONTENTS.

## CHAPTER I.

PAGE

### THE NATURE OF AIR AS A MIXTURE OF GASES.

Introductory remarks . . . . .	1
The nature of air . . . . .	3
Deductions; How much air will be absorbed by a given weight of copper? . . . . .	5
How much air will be absorbed from a given volume of copper? . . . . .	7
Calibration of the apparatus . . . . .	8
Deduction . . . . .	11
Ozone and azote; Weights of air, nitrogen and ozone; Sulfur . . . . .	13
Is the burning of sulfur similar to the scale-forming of copper? . . . . .	14
Use of litmus; Acids; Definition of an oxyd . . . . .	15
General deductions; A deoxydizing substance, carbon . . . . .	16
Metals and non-metals . . . . .	17

## CHAPTER II.

### THE NATURE OF WATER.

The ancient version of things; The kinds of water. . . . .	18
Physical properties of water . . . . .	19
Chemistry . . . . .	21
Action of steam on iron and zinc . . . . .	22
On copper; Reverse proof . . . . .	23
Hydrogen; Electrolysis . . . . .	25
Oxygen; Deductions; Inverse proof . . . . .	31
Explanation of action; Law; Idea of an atom. . . . .	33
Hydrogen peroxyd. . . . .	34

## CHAPTER III.

### GREEN VITRIOL OR COPPERAS.

Description . . . . .	35
Chemical investigation; Deduction . . . . .	37

	PAGE
Presence of sulfur established ; The oil of vitriol . . . . .	40
Preparation of a quantity of the oil . . . . .	41
Investigation of the oil . . . . .	43
A different oil . . . . .	44
Work with this other oil, the liquid residue . . . . .	45
A higher oxyd of sulfur . . . . .	46
Action of the liquid residue on lead and silver . . . . .	47
Investigation of the white fumes that crystallized . . . . .	49
Summary ; Direct proof of the presence of water in the liquid residue, the sulfuric acid . . . . .	50
Proof of the volume composition of the sulfur oxyd, $\text{SO}^2$ . . . . .	52
The dilution of sulfuric acid . . . . .	54
Molecular weight ; Hydrates of sulfuric acid . . . . .	55
Action of sulfuric hydrates on the metals . . . . .	56
The iron vitriols . . . . .	58
Preparation of sulfur dioxyd . . . . .	59
Generation of hydrogen . . . . .	61
Koenig's generator . . . . .	62

## CHAPTER IV.

## THE LESSON OF LIMESTONE.

The lime minerals . . . . .	64
Action of heat on calcite . . . . .	65
Partial proof of the lime gas . . . . .	67
Study of the residue . . . . .	70
Formation of lime hydroxyd . . . . .	71
Action towards acids . . . . .	72
Action of the lime gas upon the lime hydroxyd . . . . .	73
Deduction . . . . .	74
Summary of the limestone lesson . . . . .	75

## CHAPTER V.

## THE LESSON OF WOOD ASHES.

Wood and coal ashes . . . . .	76
Potash ; Investigation of the lye . . . . .	77
Potassium hydroxyd . . . . .	79
Hydrates . . . . .	80
Proof of the hydroxyd nature of solid caustic potash . . . . .	81

# CONTENTS.

xi

	PAGE
Action of the preceding substances on the vitriol solutions . . .	82
Potassium . . . . .	84
Physical and chemical properties of potassium. . . . .	87
Use of potassium in finding the non-metal in gas from limestone.	88
Alkaline substances . . . . .	91

## CHAPTER VI.

### THE LESSON OF COMMON SALT.

Forms of natural salt. . . . .	92
Investigation . . . . .	93
Work with the salt gas . . . . .	94
Chlorine . . . . .	98
Preparation of chlorine. . . . .	99
Chemical properties of chlorine . . . . .	104
Chlorate . . . . .	105
Process for making potassium chlorate ; Composition of salt gas.	106
Properties of hydrogen chlorid . . . . .	107
HCl + water . . . . .	108
Advantages of different driers . . . . .	111
Acid hydrometers . . . . .	112
The discovery of the metal sodium, by working with the salt cake . . . . .	113
Physical and chemical properties of sodium. . . . .	116
Oxyds of sodium . . . . .	118

## CHAPTER VII.

### THE STORY OF SODA ASH AND LEBLANC.

The Leblanc soda process. . . . .	119
Sodium carbonate and bicarbonate. . . . .	121
Caustic soda ; Lye balls ; Kelp . . . . .	122

## CHAPTER VIII.

### ON THEORIES, COMBINING WEIGHTS, ATOMIC WEIGHTS AND VALENCES.

A uniform system of writing formulas . . . . .	123
The idea of radicals . . . . .	124
The electrochemical series ; Valence . . . . .	12



	PAGE
The atomic weights of the elements; Molecular weights and volume weights . . . . .	126
Hydroxyl radical; Sulfate instead of vitriol . . . . .	129
Atomic weights of calcium, copper, lead and zinc . . . . .	130
Atomic weights of silver and gold . . . . .	131
Relation between atomic weights and specific heats of the elements . . . . .	132
Law of Dulong and Petit; Review of the action of chlorine on the alkaline hydroxyds. . . . .	133
Dichloroxyd . . . . .	134
Chlorates . . . . .	135
Oxygen from chlorates . . . . .	136

## CHAPTER IX.

## BROMINE, IODINE, FLUORINE.

Bromine from the mother liquor of the salt works . . . . .	137
Compounds and actions of bromine . . . . .	138
Iodine . . . . .	139
Properties of iodine; Starch and iodine . . . . .	141
Compounds of iodine . . . . .	142
General remarks. . . . .	143
Ismorphous substances; Fluorine . . . . .	144
Flux . . . . .	145
Investigation of the fluorite gas . . . . .	146
Composition of the gas . . . . .	148
The etching process . . . . .	149
The nature of fluorine; The metal in fluorspar . . . . .	150

## CHAPTER X.

## LECTURE ON NITER OR SALTPETER.

The kinds of niter . . . . .	152
Solubility; Investigation of the soda niter . . . . .	153
The nature of nitrogen . . . . .	154
The spirits of niter. . . . .	155
Aqua regia . . . . .	156
The properties of nitrogen . . . . .	159
Properties and composition of spirits of niter . . . . .	160
Nitrates. . . . .	164

	PAGE
Gunpowder . . . . .	166
Calculations on the explosive power of gunpowder . . . . .	168
Other powders; Investigation of the nitrous fumes . . . . .	170
Nitrogen dioxyd. . . . .	171
Nitrogen monoxyd . . . . .	176
Proof that the formula is NO . . . . .	177
Properties of NO . . . . .	178
The <i>ring test</i> for nitrates . . . . .	178
Nitrites. . . . .	179
Preparation of $\text{KNO}^2$ . . . . .	180
Dinitrogen trioxyd . . . . .	181
Nitrite test, using starch and iodine compound; Laughing gas .	182
Properties of $\text{N}^2\text{O}$ . . . . .	183
Preparation of the gas on a large scale . . . . .	184
Recapitulation of the oxyds of nitrogen. . . . .	185

## CHAPTER XI.

## AMMONIA, A VOLATILE ALKALI. A COMPLEX METALLIC RADICAL.

The <i>nascent state</i> ; Ammonia. . . . .	186
Investigation . . . . .	187
Preparation of ammonia . . . . .	188
Sal-ammoniac . . . . .	189
Composition of ammonia . . . . .	190
Proof that hydrogen is contained in ammonia. . . . .	191
Proof that ammonia contains no oxygen; Demonstration that the formula is $\text{NH}^3$ . . . . .	192
The chemical nature of ammonia. . . . .	194
Ammonium . . . . .	195
Indirect proof of the ammonium theory . . . . .	196
Formation of ammonium compounds. . . . .	197
Liquid ammonia. . . . .	200
Preparation of ammonia water . . . . .	201
Carbonates . . . . .	203
Sal-ammoniac . . . . .	204
Soldering, with use of sal-ammoniac . . . . .	205
Solvay process . . . . .	206

## CHAPTER XII.

PAGE

## ORIGIN AND OCCURRENCE OF NITER.

Sources of nitrogen . . . . .	209
Niter plantation . . . . .	210
The Chili niter deposits . . . . .	211
The origin of the deposit . . . . .	212
Conversion of soda niter into potash niter . . . . .	213

## CHAPTER XIII.

THE MANUFACTURE OF HYDROGEN SULFATE OR SULFURIC ACID  
ON A COMMERCIAL SCALE.

Direct proof that $\text{H}^2\text{SO}^4$ results from the union of $\text{SO}^3$ with $\text{H}^2\text{O}$ .	216
Cost of the sulfur-nitric acid method ; The $\text{SO}^2$ -nitric acid method . . . . .	217
The lead chamber process . . . . .	218
Sulfur from pyrite . . . . .	219
The plant needed . . . . .	220
The Glover tower . . . . .	222
Size and cost of plant . . . . .	224
The Gay-Lussac tower . . . . .	225
The Lunge-Rohrmann plate column . . . . .	226
Concentration of the chamber acid . . . . .	227
The Gridley system ; Lemaire and Co.'s stills . . . . .	228
Loss of platinum ; Manufacture of oil of vitriol by Winkler's method . . . . .	230
The direct contact method . . . . .	231

## CHAPTER XIV.

## OTHER COMPOUNDS OF SULFUR.

The sulfid ores . . . . .	233
Hydrogen sulfid . . . . .	234
Proof of the formula, $\text{H}^2\text{S}$ . . . . .	235
Generation of $\text{H}^2\text{S}$ in a steady current at a minimum of cost .	236
Properties of $\text{H}^2\text{S}$ . . . . .	237
Action of $\text{H}^2\text{S}$ on $\text{KOH}$ . . . . .	239
Action of $\text{H}^2\text{S}$ on $\text{NaOH}$ ; On $\text{Ca}(\text{HO})^2$ . . . . .	240
Calcium monosulfid . . . . .	241

# CONTENTS.

XV

	PAGE
Action of sulfur on the alkaline hydroxyds . . . . .	242
Sodium hyposulfite and its manufacture . . . . .	243
Test for thiosulfate . . . . .	244
Action of $H^2S$ upon solutions of metallic salts . . . . .	245
Examination of the precipitates obtained . . . . .	249
Hydrogen persulfid. . . . .	253
Sulfur chlorid and carbon disulfid . . . . .	254

## CHAPTER XV.

### CARBON COMPOUNDS; ORGANIC BODIES.

Native carbon . . . . .	257
Coal . . . . .	258
Carbon dioxyd . . . . .	259
Chemical properties . . . . .	260
Composition of $CO^2$ . . . . .	261
The atomic weight of carbon; Carbon monoxyd . . . . .	262
Proof of the composition of $CO$ . . . . .	263
Structure of plants . . . . .	264
Experiments with cellulose . . . . .	265
The formula of cellulose derived from the results of combustion. . . . .	266
Parchment . . . . .	270
Nitro-cellulose, gun-cotton . . . . .	271
Gun-cotton as an explosive . . . . .	272
Destructive distillation of wood . . . . .	274
Charcoal; Pyroligneous acid . . . . .	276
Acetic acid . . . . .	278
Wood tar . . . . .	279
Carbolic acid; Picric acid. . . . .	282
Paraffin . . . . .	283
Analysis of wood gas. . . . .	284
Marsh gas, methane . . . . .	294
Safety lamp . . . . .	300
The theoretical importance of marsh gas . . . . .	301
The marsh gas or paraffin series of hydro-carbons . . . . .	302
Propane; Butane; Isomerids . . . . .	303
Pentane . . . . .	304
Carbonyl-hydroxyl; Wood subjected to pressure and heat. . . . .	305

## CHAPTER XVI.

## MINERAL COAL AND ITS CHEMISTRY.

The kinds of coal . . . . .	308
Composition of coal . . . . .	310
Ultimate composition . . . . .	311
Proximate composition ; Origin of coals . . . . .	312
The coking process . . . . .	316
Coke . . . . .	318
Coal tar ; Naphthaline ; Nitrobenzol . . . . .	320
Anilin ; Complex base ; Methylanilin . . . . .	321
Rosanilin ; Mauvanilin . . . . .	322

## CHAPTER XVII.

THE GASES FROM THE DISTILLATION OF COAL ; THE MANUFACTURE  
OF ILLUMINATING GAS AND GAS COKE.

The qualitative composition of the coal gas . . . . .	324
Dissociation at high temperatures ; Plan for gas works . . . . .	325
The dimensions of the plant . . . . .	329
Water-gas . . . . .	331
Fuel-gas . . . . .	333
Rock-oil, petroleum . . . . .	335
Chemical properties ; Refining of the crude oil . . . . .	336
The flash-point ; Natural gas . . . . .	337

## CHAPTER XVIII.

## THE HOMOLOGUES OF CELLULOSE—STARCH, DEXTRIN, SUGARS.

The wheat grain . . . . .	339
Starch . . . . .	340
Chemical properties of starch ; Dextrin . . . . .	341
The sugars ; Cane sugar . . . . .	342
Grape sugar . . . . .	343
Fruit sugar . . . . .	344
Milk sugar ; Gum arabic . . . . .	345

## CHAPTER XIX.

## ALCOHOL, SPIRITS OF WINE, ETHYLHYDROXYD.

The production of alcohol . . . . .	346
Fermentation ; Yeast . . . . .	347



# CONTENTS.

xvii

	PAGE
The manufacture of ethyl alcohol . . . . .	348
Absolute alcohol . . . . .	349
Properties and chemical constitution of alcohol . . . . .	350

## CHAPTER XX.

### SOME IMPORTANT DERIVATIVES OF ALCOHOL.

Ether, ethyl oxyd . . . . .	352
Chloroform . . . . .	353
Iodoform ; Aldehyde ; Chloral . . . . .	354
Mercuric fulminate ; Preparation of the fulminate . . . . .	355
Manufacture of percussion caps . . . . .	356
Silver fulminate . . . . .	357

## CHAPTER XXI.

### SOME ORGANIC ACIDS.

Formic acid and acetic acid . . . . .	358
The manufacture of vinegar . . . . .	360
Oxalic acid . . . . .	361
Chemical properties ; Methods of manufacture . . . . .	363
Lactic acid and tartaric acids . . . . .	364
Tartar emetic . . . . .	365
Citric acid ; Malic and tannic acids . . . . .	366
Ink . . . . .	367
Leather . . . . .	368
Gallic and pyrogallic acids . . . . .	369

## CHAPTER XXII.

### THE VEGETABLE FATS AND ANIMAL FATS.

Plant oils . . . . .	370
Animal fats . . . . .	371
Chemical action of fats . . . . .	372
Glycerin . . . . .	373
Manufacture of glycerin . . . . .	374
Tri-nitro-glycerin ; Dynamite . . . . .	375
Manufacture of nitro-glycerin ; Tallow . . . . .	376
Soap ; Drying oils . . . . .	377
Turpentine . . . . .	378
Rosin . . . . .	379

	PAGE
Rubber. . . . .	380
Vulcanizing of rubber ; Ebonite. . . . .	382

## CHAPTER XXIII.

## ORGANIC ALKALOIDS.

Theobromin ; Caffein ; Urea ; Morphin . . . . .	383
Chinin ; Strychnin ; Brucin and Atropin . . . . .	384
Cocain ; Nicotin . . . . .	385

## CHAPTER XXIV.

## ALBUMEN AND ALBUMENOIDS

Albumen . . . . .	386
Haemoglobin ; Haematin. . . . .	387
Fibrin ; Casein ; Horn, hair and skin . . . . .	388
Distillation of albumenoid substances. . . . .	389
Neat's-foot oil. . . . .	390

## CHAPTER XXV.

## ORIGIN OF CYANIDS.

The discovery of the yellow prussiate of potash . . . . .	391
Potassium ferrocyanate. . . . .	392
Potassium cyanid and its preparation. . . . .	393
Chemically pure potassium cyanid . . . . .	394
Hydrogen cyanid ; Prussic acid . . . . .	395
Preparation of prussic acid and of cyanogen ; Composition of cyanogen ; Sulfocyanogen and sulfocyanates . . . . .	397
Potassium ferricyanate . . . . .	398
Potassium cyanate ; Cyanuric acid . . . . .	399
Fulminic acid. . . . .	400

## CHAPTER XXVI.

## BONE-ASH AND PHOSPHORUS.

Bone-ash . . . . .	401
Experiments on the bone-ash . . . . .	402
The discovery of phosphorus . . . . .	404
The modifications of phosphorus. . . . .	405
Red phosphorus . . . . .	406
Black phosphorus ; The atomic weight of phosphorus ; Phos- phorus pentoxyd. . . . .	407

# CONTENTS.

xix

	PAGE
Preparation of phosphorus pentoxyd ; Orthophosphoric acid. .	408
Preparation of the ortho-acid ; Orthophosphates. . . . .	409
Microcosmic salt. . . . .	410
Insoluble orthophosphates ; Molybdate test . . . . .	411
Pyro- and metaphosphoric acids and their preparation . . . .	412
Phosphorus trioxyd ; Phosphorous acid. . . . .	413
Hypophosphorous acid ; Phosphorus trichlorid . . . . .	414
Phosphorus pentachlorid ; Phosphine . . . . .	415
Preparation of phosphine. . . . .	416
Composition of phosphine ; Phosphonium. . . . .	417
Liquid hydrogen phosphid ; Metallic phosphids. . . . .	418

## APPENDIX.

The chemical elements, their symbols, equivalents, and specific gravities . . . . .	419
Table for the comparison of the scales of Réaumur's, Celsius's, and Fahrenheit's thermometers . . . . .	421
Rules for the conversion of the different thermometer degrees into each other . . . . .	422
Table of the liter weights of gases . . . . .	423
INDEX . . . . .	425





# CHEMISTRY SIMPLIFIED.

---

## CHAPTER I.

### THE NATURE OF AIR AS A MIXTURE OF GASES.

INTRODUCTORY REMARKS. I show you this small copper ingot. It was made by pouring liquid copper, at a yellow heat, into an iron mould. You notice a bright red color on the sides and bottom, a dark reddish-gray on the upper side, and a pale yellowish-pink on this spot which I scrape with the knife. By drilling a hole through the ingot, or by sawing it into two we find the pale, soft yellowish-pink color persistent throughout the interior. Hence we infer that the yellowish-pink is the *true color of copper*, while the red and gray-red at the outside must be due to a change which happened to the ingot during the time when it passed from yellow heat to the temperature of the room. If we do not take this change as a simple matter of fact, but if instead we ask for the reason of the change, then our mind is *scientific*, it is above the average, and there is hope for us. Now there are two ways for satisfying this desire for the reason for things. Along one way we just ask the nearest man or the handiest book, along the other way we set to work it out ourselves. Following the second and much harder road, we become investigators and inventors.



You have come here to learn chemistry, and my duty is to show you the way. I may lead you along the first trail by means of a book and recitations and my own acquired knowledge of the nature of things. However, I choose to take you by the second trail, difficult in steep slopes and rapid descents, but which will ultimately take you, when the pass shall have been won, into the beautiful valley of intellectual satisfaction and fruitful technical invention. Some may get exhausted on the journey, and some may not have the right eyes to see, the right hands to grapple with the difficulties; for them of course there is no hope and they must take the other trail, they will not become leaders in the profession. Those who are born with right desire will ultimately get to the goal by any road whatsoever; but they will have spent much time, much energy, in doing useless things. *My object, as your teacher, is to point out things to you so that you may save the waste and arrive much more rapidly at the pass.*

Let us return to our ingot and ask: What causes the copper to be of different colors on the outside from the inside? This question we name, by general consent, a *chemical question* because the change which happened to the outside of the ingot is a *permanent change*, by which the substance has acquired different properties throughout. If I bend this wire there is also a permanent change, but it is only a change of shape, the copper itself has not changed any of its properties. If we inquire about this change of form in the new positions of the copper

particles, we have to deal with a proposition in *physics*. There are, of course, many changes in which physics and chemistry overlap, when we must enter into both sciences. It is merely a matter of convenience, for the sake of more effective work that this division into physics and chemistry has been made. One set of men pursue the one set of phenomena, becoming more expert in observation than if they spread themselves over both sets.

### 1. *The nature of air.*

Pursuing the question concerning the changes at the surface of the copper ingot, we shall now institute experiments, every one of which will give rise to new questions.

In Figs. 1, 2, 3, 4 the trial arrangements are set up to meet those questions.

In Fig. 1, a bright piece of sheet copper *A* is placed within the non-luminous or purplish part of the flame *F*. It is evident that here the metal will be exposed to the action of heat, to the action of air and to the action of the substance of the flame. The metal at bright red heat, under these conditions covers itself with a dark scale, which peels off and is found to be very brittle. In Fig. 2, the metal *A* is within the hard-glass tube *T*, which is closed at one end, open at the other end. The substance of the flame does not come in contact with the metal, but the heat of the flame is conducted through the glass, the metal becomes red hot and covers itself with a very thin film. Here the air has access, but

limited. Fig. 3, shows the metal strip *A* within a tube *T*. The tube was then drawn out at *C*, leaving a narrow neck, was then connected with an air pump until all air was drawn out, when the neck *C* was allowed to close up by means of a sharp, hot flame (blow-pipe flame). If then the tube *T* be

FIG. 1.



FIG. 2.




FIG. 3.





FIG. 4.




made red hot in the flame *F* and kept so for any length of time, we find, on cooling, that no change has taken place, the copper is bright with its fine pale yellowish-pink color. Lastly in Fig. 4, we have the copper strip in a glass tube which is open at both ends. Owing to inclined position a lively air current will set in in the direction of the arrows as

soon as the tube is heated by the flame. We find quickly that the last conditions produce the most change on the copper—the most rapid development of the scale.

Deduction from the four experiments: 1. Copper takes from the air, at red heat, a portion of the air and changes to a brittle scale. 2. The substance of the flame does not enter into this change, only the heat of the flame. Two questions now suggest themselves: 1. How much will be absorbed by a given weight of copper? 2. How much of a given

FIG. 5.




volume of air will be absorbed? In order to answer the first question rig an apparatus as shown in Fig. 5, where *T* is an infusible glass tube (so-called combustion tube) not less than  $\frac{1}{2}$ " inside diameter. The tube is bent over the blast lamp. Place the tube in a horizontal position by means of clamp-stand 5. Select a porcelain boat sufficiently narrow to pass into the tube easily. Clean and weigh the boat. We choose glazed porcelain because it will not become soft except at white heat, because it does not change in weight when heated

in air. Place into this boat about 0.100 gram of copper filings, that is to say you must take the weight very accurately. But it does not matter whether you take 0.095 or 0.102, only by taking 0.100 accurately you will save figuring the percentage. You are told to take filings, because in this form the metal will be exposed with a maximum of surface and a minimum volume, which is most desirable, since the intended action proceeds from the surface. Shove the boat into the tube as shown in the figure, and place the lamp so that the entire length of the boat can be brought to redness. The bent part of the tube being bent upwards, a so-called draught will arise as by a chimney, because a column of hot air being in the vertical tube, this hot air being lighter than the cold outer air, the latter will pass in at the lower end and push out the warm air, hence a steady air current will pass through the tube in the direction of the arrow's pointing. Allow this action to proceed for 30 minutes. Do not forget to place a few fibres of infusible asbestos between the tube and the boat, for the tube may become soft and cause the boat to stick. Should the tube exhibit a tendency to sag you will prop it up at the end by means of a piece of brick. Remove the flame at the expiration of the half hour, let the tube cool down until it can be held in the hand. Pull out the boat and weigh. Return the boat to the tube, replace the lamps and work for another half hour. Then weigh again. If the second weight is not equal to the first, you cannot be certain that the



copper has saturated itself, and a third period of heating becomes necessary, and perhaps a fourth until you get constant weight. Many trials, made with greatest care, show that 0.100 copper will in-

FIG. 6.



crease to 0.1254 and no further, and thus the resulting dark gray-black product of change will contain in 100 : Copper 79.6 and air 20.4.

2. *How much will be absorbed from a given volume of air?*


For an answer to this question let an apparatus

be rigged as in Fig. 6. We must make provision to pass and repass the same volume of air over the heated copper without any air coming from the outside, nor any air escaping to the outside. Let  $T$  again be of hard glass  $\frac{1}{8}$  or  $\frac{3}{16}$  inch internal diameter and 5 inches long. Introduce a roll of fine copper gauze, 5, then bend the tube into a double  $L$ . Paste on a mark at  $M$ . Determine or gauge the volume of the tube by filling it with mercury up to the mark  $M$ , and then weigh the mercury. If the weight of the mercury be  $g$ , then  $g/13.6 = V$  (in cubic centimeters). Provide two soft glass tubes,  $S, S'$ , with stoppers, the upper ones to receive the tube  $T$ , the lower ones to receive narrow glass tubes, which are connected by rubber tubing, 2, 4, with the glass syphons 1, 3. These stand in beaker glasses,  $R, R'$ . Calibrate the tube  $S$  into cubic centimeters, the lower side of the upper stopper forming the 0 mark.

*Calibration.* Stand  $S$  upside down, after a solid stopper 1, Fig. 7, has been pushed into the 0 mark. Then weigh out (to 0.1 gram)  $5 \times 13.6 = 68.0$  grams of mercury and pour this into the tube. Make a scratch (with a glazier's diamond, with a sharp splinter of quartz, or with a hard file) tangent to the meniscus. Measure the distance from 0 to 5 with a scale or pair of dividers. Lay this distance down a strip of white paper and divide into 5 equal parts. Lay on these divisions beyond five for the whole length of the tube. Then paste the strip upon the tube so that the 0 marks fall together, and then cover the paper on the outside with molten

paraffin. Thus your tube is calibrated into cubic centimeters. Weighing out a defined quantity of mercury is not quite so easy as it sounds. Proceed as follows: Place a very small beaker glass upon a so-called pulp balance (it were foolish to use a fine balance because we only want to weigh as close as 0.1 gram). Why? Because 0.1 gram of mer-


FIG. 7.



cury corresponds to less than 0.01 cubic cent., and we do not care to read a volume closer than 0.1 cubic cent. It would in fact be quite sufficient accuracy if we got any weight of mercury within 1.0 gram. Make a pipette with capillary outlet by drawing out an 8" x  $\frac{1}{8}$ " soft glass tube over the lamp flame as shown in Fig. 8, and cut off with file at point *a*. Fill this tube—by sucking—with mercury, and let this latter flow into the beaker, until the balance tips. The beaker has been previously balanced by shot, and the required weight, 68 grs., has been placed upon the weights' pan. The first finger closes the upper end of the tube at the moment of tipping. After the apparatus (Fig. 6)

has been set up—the tubes  $S$ ,  $S'$  being held in place by clamp-stands or tripods or under propings—the beaker glasses  $R$ ,  $R'$  are filled with water, the syphons  $1$ ,  $3$  are filled by sucking at the end of the detached rubber tubing until the water comes to the mouth, the rubber being then joined to the tubes  $S$ ,  $S'$ .  $R'$  is then raised upon

FIG. 8.




blocks until the water reaches the mark  $M$ , while  $R$  is lowered until its water level falls together with a division of the scale on  $S$ , say for instance 10. We are having then a volume of air  $V$  equal to 10 plus  $v$  ( $v$  being volume of tube  $T$ ). Let  $v$  be 5 c.c. for example; then the total volume will be 10 plus 5 equal 15 c.c. Bring now the burner  $B$  under  $T$  so that the copper gauze 5 becomes red hot. Cause the air to move slowly over the hot copper, by lowering beaker  $R'$  and raising  $R$  at the same time. Repeat this movement 10 times. Then remove the burner and let the tube  $T$  return to the temperature of the room. Bring the water to the mark  $M$  and adjust  $R$  in such a way that its water-line and the water-line in  $S$  fall into one level. On reading off the position of the water-line  $S$  on the scale, we will read 7, that is, 3 c.c. of air have disappeared; 15 c.c. of air have lost 3 c.c., 5 c.c. have lost 1 c.c. Bring the flame

back and again pass 10 times over the hot copper. On cooling will find the same result as before; hence the loss of the air, under these conditions, is constant.

*Deduction.* Air is composed of at least two parts, which possess each decidedly different properties, or in other words: Air consists of two different gases. Now it is well known that men and animals must have air to breathe. Query? Which one of the two gases do they require, or do they need both? As the one gas has disappeared into the copper, we shall have to experiment with the  $\frac{4}{5}$  residuum. Let a glass jar *J*, Fig. 9, be filled with the gas. How? A two-foot length of  $\frac{3}{4}$ " gas pipe *2* (Fig. 10), is charged

FIG. 9.

FIG. 10.



with copper gauze, and placed in a charcoal furnace *4*. Rubber tube *5* leads to bottle *6* through a two-hole stopper; so that the funnel stem *7* may pass through second hole. If water be poured into the funnel the air will be driven through the heated copper, and the  $\frac{4}{5}$  residuum will pass through *1* into the inverted and water-filled jar *J* whose open end is supported under water by metal blocks in the basin *B*. We will now do a little figuring. We

found above that 1 gram of copper can absorb 0.25 gram of air. One litre equals 1000 c.c. of air and weighs 1.2932 gram, hence  $\frac{1}{5}$  of this weight is 0.258, that is one single gram of copper is sufficient to take the absorbable gas from one litre or one quart of air. 50 grams of copper will furnish  $(50 \times 4) / 5$  equal 40 litres of the gas we desire. But for our experiment, that is to fill the jar, we will not need more than one or at the most two litres supposing that the whole of the water has been displaced in *J* so that the bubbles will come on the outside. We remove first the tube *I*, then the supporting blocks and push a flat glass plate under the water, and press it with the hand against the jar rim, lift the jar out and stand it bottom down on the table as in Fig. 9. A mouse having been trapped, we drop the animal from the trap into the jar and replace the glass cover plate. At once the mouse shows distress, tumbles into a heap, and after a few spasmodic kicks lies quiet and is dead. Hence we draw the conclusion that the part of the air which is absorbed by the copper, is the part also which sustains animal life. If we introduce a burning taper into the jar it will become at once extinguished, and hence it follows with some considerable probability that breathing, burning and scale forming of copper are *similar* processes, being in fact, fundamentally, the same phenomenon.

It will be necessary, for clearness and conciseness of expression, to distinguish from now on these two parts of the air by separate names. Properly we



should say—in English—life-air and death-air; instead Greek words are chosen by scientists. Life air equals ozone; death air equals azote. The root of both words is Zoë equals life (zoölogy equals the science of living things). Ozone equals intense life; azote equals no life. German chemists made and use the term, stickstoff equal to suffocating stuff; the French hold on to azote, English and Americans now use the word *nitrogen* which equals niter-producer.

Weight of one c.c. of nitrogen equals 0.001256 gram (Regnault).

Weight of one c.c. of air equals 0.001293 gram.

Hence it follows that ozone must be heavier than air. The weight of ozone follows by calculation.

We reduce azote to terms of air by dividing the weight of air into that of azote,  $0.001256 / 0.001293$  equals 0.97137 which equals the *specific* weight of azote or its specific gravity, then we have


$$4 \times \frac{0.9713}{5} + \frac{x}{5} = 1; \quad x = (1 - 0.777)5 = 1.1145,$$

the specific gravity of ozone, and  $1.1145 \times 1.293 = 1.441$ , the weight of 1000 c.c. of ozone, and 0.00144, the weight of one c.c. of ozone.

Among the substances more or less familiar to every one of us is sulfur, which I show you now as crystal and massive as a broken lump. Light yellow color and translucency distinguish it from other bodies. If washed in water first, it yields no taste, it is insoluble in water. Against heat it is quite

susceptible. I place some of it upon this porcelain crucible lid and put the flame underneath. We see it melt quickly into a dark red-brown liquid, and then vaporize. Shortly after a blue flame appears and a strong, pungent odor permeates the air. The sulfur disappears slowly. Queries. 1. Is this phenomenon similar to that of the scale forming of copper, to the burning of the taper, to the breathing? Has either the ozone or the azote part in the disappearance? Let the flask 1 be filled with air and a few c.c. of water as shown in Fig. 11. Take

FIG. 11.



a strong iron wire 2, hammer it flat at one end and rivet it to a small iron dish 3. Put sulfur into the dish, heat the latter over a flame until the blue flame is strong, then lower the dish into the flask until the plate covers the mouth of the flask. The sulfur keeps on burning. Soon a white cloud obscures the flame. After some minutes we remove the spoon and notice that it begins to burn when it reaches the outer air. (Why? Because it again finds the ozone, which had become used up in the

flask.) We now shake the water and the gases together in the flask and pour out the water. It is somewhat turbid or milky ; after filtering it is clear, it has a sour taste and changes the purplish color of litmus to bright *onion red*. Litmus is obtained by extracting a lichen named by botanists *Roccella tinctoria*. The collected lichen is allowed to ferment in heaps, is then torn by a machine and allowed to stand with water, which assumes a deep purplish color ; is used as a dye stuff under the name of *orseille*. Now vinegar has the same action, *i. e.*, a sour taste and turns litmus red. Vinegar (from French *vin-aigre* equal wine sour) arises when the sugary juice of grape or any other fruit, is allowed to stand in a warm place.

Wine results in a cool place. (Details will be given later.) The Latin name for vinegar is *acidum*, hence it came that later on all *sour* substances were and are called acids in English, French, Spanish, Italian ; but in German the word *säure* = a sourness, and in Swedish *syra*, only differing in the vowel.

When Lavoisier, in 1781, had observed the above phenomena, he proposed the name *oxygène* in French, *oxygen* in English for what we have named ozone or life-air. The word oxygen is derived from the two Greek words *oxus* = sour, sharp, biting and *genomein* = to produce, to generate. This name we shall use in the future.

*Definition* : The combination of oxygen with a metal or non-metal is hereafter to be designated an

*oxyd—oxid—oxide.* The first spelling is the grammatically more correct, but the last spelling *oxide* is at present mostly used by English and American chemists. I myself use *oxyd*—the derivation is from *oxydatus* = oxydized.

Hence copper scale is now copper oxyd, the penetrating gas from burning sulfur is sulfur oxyd.

*General deductions.* Air being a compound of azote + oxygen and copper oxyd of copper + oxygen, they both seem to be representatives of the same type of bodies *i. e.*, *compounds*. Whether all compounds are of one kind or not, we cannot, at this stage demonstrate. The term *compound* requires *single*, or *simple* as opposite. We will define as *simple bodies* or *elements* all those bodies which can be brought back from their compounds to their first condition *i. e.*, with all their previous physical properties. This definition is raised in our minds by the behavior of copper oxyd when heated in a closed tube with splinters of charcoal. At a red heat, the scale of copper oxyd becomes yellowish-pinkish. On cooling a spongy mass of the metal is seen, and the metal possesses all previous properties. Hence copper is a simple body or element. The question arises: Are all our well-known metals elements? Experiment on the same line as that with the copper, answers the query in the affirmative, though there will be found experimental difficulties with some. When zinc has been burnt in air, at high heat, into white zincoxyd, and when we heat this oxyd with charcoal, we will get no satis-

faction because the zinc only gives up its oxygen to the coal at a *white heat* which the glass tube cannot stand; and moreover, at this high temperature zinc itself is a vapor. Special arrangements must be made in this case to catch the zinc vapors, which will first condense to a liquid and the latter will solidify into metal with all previous properties. Though we know as yet nothing of the nature of charcoal, we must surmise that it contains a body which has a stronger attraction or affinity for oxygen, at high heat, than the metal. Owing to this we designate charcoal as an *deoxydizing* substance, and the process itself we call *deoxydation*.

Copperoxyd + charcoal + red heat = deoxydation. That the charcoal forms a new oxyd during this action we surmise at once. Because we see no deposit of any kind on the charcoal or on the tube, we infer that this new oxyd is a gas at the ordinary temperature of the air. By using only the first letters as symbols, we can represent the processes thus:

$\text{Cu} + \text{O} + \text{red heat} = \text{CuO} = \text{oxydation}.$

$\text{CuO} + \text{C (charcoal)} + \text{red heat} = \text{Cu} + \text{CO} = \text{deoxydation}.$

Sulfur oxyd can be decomposed by passing it in a glass tube over heated charcoal. Sulfur will deposit in the tube; hence sulfur is an element, according to our definition. Yet sulfur is not at all like the metals. As mentioned above, it is transparent and very brittle, melts at a very low heat and becomes a vapor at  $250^{\circ} \text{C}$ . Hence we at once make two classes of elements: *metallic elements*: copper, tin, lead, iron, and so forth; *non-metallic elements*: sulfur, oxygen, azote, etc.



## CHAPTER II.

### THE NATURE OF WATER.

To us who are dwelling on the shores of a great lake, or to the inhabitants of the seashore, the idea must early present itself that water must be of much importance in the household of our planet, but even to him, the savage, or half savage, who uses water merely to quench his thirst, water must be an object of veneration; and the inquisitiveness of the human mind must bring forth even in him a desire to learn more about it. The ancient thinkers, without experiments, said water is one of the primary, elementary things, and together with air, fire and earth, constitutes the great Quartette out of which comes the harmony or disharmony of all things. Man's body is made of earth and water, his soul of air and fire, thus combining within himself the four primary things. The elementary nature of water was undoubted up to the end of the 18th century. The traveler becomes acquainted with good water and bad water; water which has a bitter taste, an astringent taste, a salty taste, a nauseous taste. He learns to distinguish between Spring water, River water, Swamp water, Sea water. It was early observed that water combines with fire, and thus produces the scalding steam which would lift the cover off a pot.



Water becomes a solid, transparent block under the influence and sway of chilling Boreas, the Northwind. To this splendid and fleeting thing the Greeks gave the name *Krystallos*, sometimes *Kryos*, and thus became the founders of *Crystallography*, because when they found the beautiful transparent *Rockcrystal*, which we now call *Quartz*, they considered it as a sort of *permanent ice*, and also called it *Krystallos*. Thus the name was applied much later to all bodies which exhibit geometrical outlines. Remember this connection between ice and crystal. More wonderful seemed the snow, and it took much hard thinking and much controversy until snow was admitted to be merely frozen rain. You throw your snowballs and skate over the ice and never once think anything at all, except that it is a part of Winter's fun, and yet for a hundred years men have been studying the form of snow-flakes, and each one found something new.

*Physical properties of water.* Reference is here made only to so-called distilled water, that is, water which has been in the form of steam at least twice, and which has only been kept in *porcelain* crocks. All other water is impure in differing degrees.

Water is a liquid between the temperature of  $0^{\circ}$  and  $100^{\circ}$  C. Below  $0^{\circ}$  C. it is solid—ice—brittle, and yet to some extent plastic, that is, capable of deformation under strong, slow pressure. (Bending of glaciers when plowing over undulating rocks.) Ice is colorless in small pieces, in great masses it shows a green or sometimes a deep blue color.

Water is without taste (distilled water); hence not pleasant to the palate. It is without color when seen in a bottle, flask or jar; but when looking through a long column it becomes more and more blue, that is, all but the blue part of the sunlight becomes absorbed. The finest effect is seen off the Savoy shore on the Lake of Geneva (Switzerland), where the rocks fall abruptly into the water to a depth of 980 feet. The ocean looks blue-black, but this is not water which can be compared to distilled water, whereas the water of the Geneva Lake is remarkably pure, having a steady and strong outflow in the Rhone River.


A litre of water at  $+4^{\circ}$  C. weighs sensibly more than at  $0^{\circ}$  C., hence ice will float on water. If, however, the temperature of water is raised by heating it will loose in density and soon the block of ice will sink to the bottom.

Water *dissolves* all solids, some very slightly, others in large quantities. Note this very important property. The purer the water the more energetic is the solving action. It attacks and slowly dissolves ordinary glass, but porcelain very much less, and platinum and gold still less.

Water absorbs all gases, some to a large extent, others to a very small extent. Hence the purest distilled water will not remain so but for a very short time. For absorption begins as soon as the stopper is removed, in fact has already set in before, unless the vessel had been full up to the stopper. As a rule all absorbed gases can be expelled by prolonged boiling.

*Chemistry.* The first query will be: Is water a simple, elementary body or is it a compound? Having observed that air can be broken up by metals and by sulfur at a red heat, we shall be consistent if we subject water to the same treatment. We start, therefore, with the assumption that water is *not* a simple body. The experiment will presumably proceed under proper conditions if a hard glass tube *T*, Fig. 12, with perforated stoppers be placed horizontally with a boat holding finely granulated zinc or iron filings in the middle of the tube. A glass syphon *5* reaching into the beaker is con-

FIG. 12.



nected by rubber *1* with stopper *2* while a clamp *3* permits a stop to the flow of water as well as a regulation of the rate of inflow. A delivery tube *5* leads just under the surface of the water in basin *6* and a test-tube *7* filled with water stands ready to receive any gases which might result from the action.

Let the flame from the burner *B* be now put under the boat and water admitted drop by drop at *2* so that a small pool forms behind the cork. As the heat extends along the glass tube the water will

begin to pass into steam and the steam will push out the air. When the latter is all out the steam will condense in tube 5 and of a sudden water will rush from 5 into T, will run to the red hot place at the boat and the T cracks. The apparatus is not up to the requirements. Why should not an iron gas pipe do in place of a glass tube? Since we are studying the action of steam upon the metals a metallic tube can do no harm, and any sudden in-rush of water cannot injure the tube. Let therefore a gas-pipe of similar dimensions be substituted. A further advantage of such a tube will be that a higher degree of heat can be applied—say by means of a gas blow-pipe.

### 1. *Action on bright iron turnings.*

A gas is generated which we collect in the test-tube. It possesses an unpleasant odor and burns with a faintly colored flame. The chips have lost their brightness, they have become gray-black, and when struck with the hammer a brittle scale comes off. The scale is attracted by the magnet the same as the chips themselves. *Deduction:* This scale, resembling in every respect ordinary blacksmith's hammer scale, which is made by heating iron in air, like the other metal scales, leads us to conclude that water must contain oxygen.

### 2. *Action upon granulated zinc.*

Gas of the same character but odor less pronounced. The zinc is either converted wholly or

partly into white or grayish-white powder, or into a shining lustrous crust, which under the microscope shows six-sided prisms and pyramids. Character same as *zinc oxyd* obtained in air.

### 3. *Action upon copper turnings.*


Gas does not appear until tube has become yellow hot, and comes sparingly. Has no pronounced odor, but burns. The copper has become bright red from the brittle *copper oxyd*.

The three trials demonstrate that water is a compound, not an element; and further, that it is an *oxyd*, in which oxygen is united with the other body, the latter appearing in the free state as a light, odoriferous gas.

*Reverse proof.* Collect the gas in a large bottle (use arrangement for collecting as in the case of azote). Stand the bottle 1 (Fig. 13) containing the gas upright. Its stopper holds the long funnel tube 2, with rubber connection and clamp 3 and the delivery tube 4 which connects with the drying tube the latter being filled in the bulb with absorbent cotton and in the cylinder with burnt lime. Tube 6 is drawn out into a fine point (not too fine), and the glass cylinder, open at both ends, is held by a clamp and stand as shown in figure. If now water be run from the funnel into bottle, the water will drive the gas through, and if a burning taper be held to the tip of 6 two things can happen: either a steady flame 8 or an explosion which may shatter the tubes. (1 volume of the gas mixed with

$2\frac{1}{2}$  volumes of air explodes violently.) To avoid explosion, let the gas go through the tube for a while—until the water has risen in the bottle about one inch—and then only apply match or taper. The flame is first colorless, but soon burns of an orange-yellow color. Why? Because the glass, at a red heat, gives particles to the flame. Prove this

FIG. 13.



by placing a nozzle of silver, platinum or gold over the tube. The flame will then be invisible or nearly so, faintly purplish.

But the important part is that immediately when the flame appears the glass cylinder 7 becomes clouded and soon large drops of liquid condense upon the glass which will even collect and run from the lower edge of the tube. We find all the physi-




cal properties of distilled water in this liquid: Equal weight per equal volume; absence of taste and smell; equal resistance to passage of light; change into solid at  $0^{\circ}$  C.; change into vapor at  $100^{\circ}$  C.; very high resistance to the passage of the electric current; equal capillarity, by which is meant that both liquids rise in a very narrow or hair tube equally high above the outer level. Therefore, it is proper to apply the name *hydrogen* to this gas. *Hydor* = water, *genomein* = to produce: the body which produces water. It follows that water must be *hydrogen oxyd*.

#### ELECTROLYSIS.

Next to heat we find electricity as the most intense force, or probably both are only different manifestations of the same force, since heat can be made to generate electric current, and an electric current converts itself into heat under proper conditions. We will not investigate the generation of a current here and simply assume it as given; you will get the explanation in physics. We simply accept the fact that if I fasten this wire to the one binding post on the table and this other wire to the second post, and if the wire ends be brought in contact there is now flowing a current from one post to the other. If I break the contact a minute blue spark is the visible proof that a current was passing, but does not pass now. Why? The air space between the wires does not carry the current, air being a *non-conductor*.

In Fig. 14 a beaker glass is filled with distilled water; after bending the wires at right angles I sink the vertical portions into the water. The needle on this galvanometer does not move, hence it follows that the water between the two wires is as much a non-conductor as air; of course I speak relatively; it is a question of tension or pressure, for the current passes from cloud to earth in a

FIG. 14.




thunder-storm, and to say that air and pure water are bad conductors, will be a better expression than non-conductors.

If we substitute this Houghton spring water for the distilled water, we see the needle move, and hence deduce that the earthy materials which are dissolved in the water cause the better conductivity. An addition of salt to the water proves the correctness of the deduction. We see not only the diversion of the needle, but we see gas bubbles rise from the wires. The liquid conductor of the current is called *electrolyte*. The wires dipping into the latter are known as *electrodes*. The wire leading to the positive pole + is the *anode*, the one leading to the negative pole — is the *cathode*. They are always designated by the + and — signs.

If we arrange the apparatus as shown in Fig. 15, where 2 means a test-tube, filled with the electrolyte, and where the electrodes are spread, having strips of platinum sheet soldered to themselves, we will perceive at once—the current going through—that much more gas arises from the cathode than from the anode. On applying a match to the gas, after the test-tube has been lifted out, there is a strong

FIG. 15.



explosive sound ; the tube may be splintered. We call the gas *fulminating* (lightning like). What is the nature of this gas? That two different gases are produced we find indicated in the larger volume of gas coming from one of the electrodes ; and thus we are led to collect the two portions *separately*. Let us take two large test-tubes *B*, *B'* (Fig. 16) and blow into the closed end two narrow tubes *1*, *2* before the blow-pipe. This requires some skill acquired by practice. To acquire such skill is very useful to any one who wishes to become a chemist ; for glass-blowers are not always in the neighborhood.

However, in this case, we can avoid glass-blowing altogether by cutting off the closed end of the tubes by means of a file and a red-hot glass or iron rod. In

FIG. 16.





Fig. 17 you see the tube before and after cutting, and only a perforated stopper is needed with a narrow glass tube *t*, to which the short rubber tube with spring or screw clamp and a second bit of glass tube are attached. Calibrate now the tubes roughly into cubic centimeters. Fill the tubes with the electrolyte, hold them by a suitable stand in the basin *A* (Fig. 16), and insert from below the electrodes + —. The current being turned on it soon becomes evident that the gas volume in *B'* is much larger than in *B*, that in fact the two volumes are as two

to one. But in  $B'$  we have the negative pole, the cathode; in  $B$ , the positive pole, the anode. When  $B'$  is filled with gas to the electrode we stop the current; draw out the electrode, shove a small cup under the tube, lift the latter and bring it into a


FIG. 17.



wide cylinder (a pail will answer, in default). Now if we press the tube  $B'$  down into the water until there be a difference,  $H$ , Fig. 18, between the outer and the inner levels, then the gas will be under a pressure equal to the weight of a column of water of the height  $H$ , and if the clamp  $C$  be cautiously opened the gas must issue at  $P$ . We notice first the *absence* of any smell; the absence of acidity, and then apply a taper or match. A flame appears at the point. A cold porcelain dish held in the flame covers itself with moisture. In fact the gas acts exactly like the Hydrogen which we obtained previously, except that it has no odor. The former gas must have held some odoriferous body admixed. (Supposition which must be verified at a future

stage. As careful investigators we must heed every difference of action. In this instance we shall find that zinc and iron contain small quantities of cer-

FIG. 18.



tain other elements, which combine with hydrogen, producing strongly smelling gases.)

We proceed to operate now with tube *B* (Fig. 16) by transference to the pail or large beaker. The gas has neither odor nor taste. On bringing a match over the jet no flame, but the coal of the match burns with intense light. . A piece of red-hot copper




or iron wire held into the current, burns with strong light into a scale which has all the properties of copper oxyd, iron oxyd.

The gas shows therefore in an intensified form the behavior of air. It must be *oxygen*. That it is only oxygen we can prove by heating a given volume of iron or copper filings over mercury. On cooling the mercury will fill the entire tube; all the gas has been absorbed.

*Deductions.* 1. Water is a chemical compound of the two simple bodies, hydrogen and oxygen. It must be a chemical compound because water is a body showing altogether different properties from those of a mere mixture of the gases as we have it in fulminating gas.

2. Elements combine according to simple ratio of volumes, as water is composed by volume of 2 hydrogen, 1 oxygen. If the letters H and O stand for the two elements then the symbol




expresses this relation.

*Inverse proof* of this proposition.

Let *E* (Fig. 19) be a graduated or calibrated glass tube, near the closed end of which are fused into the glass 2 platinum wires (+ —). Such a tube is known as *eudiometer* = splendid or perfect measure. Let it be furnished with a jacket *J* made of a wide glass tube (lamp chimney); a cork *C* holds it against the eudiometer. A narrow glass tube leads from the boiling flask *F* through the cork into the jacket; *th* is a

thermometer. Introduce about 10 c.c. of fulminating gas into the eudiometer, the mercury standing as in the figure. If now the water in *F* be kept hard boiling until the thermometer shows  $98^{\circ}$  to  $100^{\circ}$  C., the volume of the gas will have increased considerably. Let us say it reads 11.0 c.c. Connect the platinum wires with poles of an induction coil, also known as Rhumkorf coil from the inventor's

FIG. 19.



name; a spark will pass between the platinum wire ends in the eudiometer with a considerable shock. (Hold the eudiometer with the hand to prevent its being lifted clear of the mercury in *P*.) On reading now the volume, we find it shrunk to 7.3 c.c. A contraction has taken place of 3.7 c.c. or *one-third* of the original volume. Stop the steam, draw off condensed water, and let come to the temperature of the room. The mercury will rise steadily until it

fills the entire tube, except a bubble of water at the very apex.

*Explanation.* The high temperature of the spark causes the union of the hydrogen with the oxygen, and at the temperature of  $100^{\circ}$  C. water is in the form of steam. The steam occupies two-thirds the space which was occupied by the mixture of gases. The union therefore caused the mass particles of the gases to approach each other at a fixed distance, so that three occupy now the space of two. 2 vols. H + 1 vol. O give 2 vols. steam. As the temperature sinks the steam becomes water and thus the mercury can fill the tube because the quantity of steam expressed by 7.3 c.c. corresponds only to a few milligrams of water, for 1 c.c. of steam at  $100^{\circ}$  C. weighs 0.0005896 gram, hence  $7.3 \text{ c.c.} = 0.0043$  or 4.3 milligrams, which in the form of a drop of water is barely distinguishable with the naked eye. The 11 c.c. of gas must have possessed the same weight of 4.3 milligrams.

*Law.* Whenever gaseous elements unite in the ratio of 2 : 1 a contraction of one-third ensues.

Comparing the weights of the 3 gasiform elements which we have now discovered and studied to some extent, we find that hydrogen : oxygen : azote = 1 : 16 : 14. Oxygen is 16 times heavier in equal volume than hydrogen, and azote 14 times heavier. Therefore  $\text{H}^2\text{O}$  represents  $2 + 16 = 18$  weight units and if the *smallest mass of each of these bodies, still capable of receiving a chemical impetus and acquiring thereby an active force or momentum*, be de-

signated as *atom*, then the figures 1, 16, 14 may be designated *atomic weights*, because the ratio will be the same as long as the volume is the same, for the largest as well as the smallest units.

*Hydrogen peroxyd*,  $\text{H}^2\text{O}^2$ .

Under certain conditions H and O can form a higher oxyd or *peroxyd* (*per* meaning beyond). This is a body of unstable nature ; it falls to pieces easily but possesses extraordinary activity as an oxydizing agent. We shall study the mode of its preparation later as well as its application.

## CHAPTER III.

### GREEN VITRIOL OR COPPERAS.

UNDER the name *copperas*, which is the corrupted form of the French *couperose*, and the latter even a corruption of the Latin *cupriaerosa*, which means the "Rose of Cyprus," a substance of most marked properties has been known from time immemorial. We see it as bluish-green fragments and also in the form of large crystals. The symmetry of the faces can be reduced to the monoclinic system. Thin pieces are quite transparent and show very little color; because the depth of color is determined by the partial absorption of certain portions of sun light. The thicker pieces absorb more light in the transit of the latter and therefore have a deeper color. The substance is very brittle; it can be crunched between the fingers. Has a strongly astringent, bitter-sour taste, and is quite soluble in cold water, but very much more soluble in boiling water. The solution is not clear, but murky from a brownish-yellow, suspended substance. By filtering through paper the solution becomes clear. If such a clear solution be left standing exposed to the air, one notices the rapid forming of a yellow film on the exposed surface. The film becomes thicker and in time the liquid turns into a yellow brown,

mud-like fluid. Query? Has the oxygen of the air anything to do with this change, or the azote? This is easily settled by bringing some of the fresh, clear solution into a tube filled with azote and closing the tube tight. No change occurs. Another portion in a tube filled with oxygen, the film appears. Oxygen does the work. If the solution of copperas comes together with oak bark, the latter turns speedily black (in reality deeply blue-purple). It also turns leather black because the latter forms when raw hide is allowed to soak in water and ground oak or hemlock bark. We can extract the leather-forming substance from the bark by means of water and this solution turns black with copperas. Thus came into existence our black writing ink, already known to the ancient Egyptians. When the extract of bark is evaporated on a water-bath, a faint yellowish scale, very light and fluffy, remains: *tannin* (from *tan*). Copperas + water + tannin = ink: mucilage is added to give the ink a better body.


The Germans use the word *vitriol* instead of copperas. This name was first used by Pliny who lived 1900 years ago. He describes the substance as "*vitriolus quasi vitrum*." Vitrum is Latin for glass, the crystals resembling green glass were yet easily soluble in water, while glass is not soluble. Glass is fairly hard, this substance not, hence *vitriolus*, a kind of glass, like glass in some ways. We shall use the word *vitriol* as being better expressive of the substance.



## CHEMICAL INVESTIGATION OF GREEN VITRIOL.

Let a hard glass tube ( $\frac{1}{8}$  to  $\frac{1}{4}$ " internal diameter and 6 to 8 inches long) (Fig. 20) be closed at one end over the gas blow-pipe. Hold the tube in nearly horizontal position or rather lower at the open end, the

FIG. 20.




closed end being charged with several small pieces of vitriol. Let the tube be supported at *S*, because it has a tendency to sag when the end comes to a red heat. It stands to reason that the tube may be laid upon a couple of bricks for support. Let heat be applied slowly with the burner *B* to the closed end, and we will be enabled to make the following observations: 1. A plentiful condensation of a liquid in the cool part of the tube. This liquid will appear very mobile, little or no taste and by applying electrolysis will give  $H + O$ ; its identity with water is fixed. 2. The vitriol meanwhile, is partly melting, raising blisters and turning into a white chalky mass.

Deduction 1: The loss of water causes the vitriol to become white opaque. We raise the temperature to a low redness: A strongly smelling gas appears re-


calling the smell of burning sulfur, that is the oxyd of sulfur  $S^nO^m$ . We cause it to act on indigo and litmus solutions, the former becomes bleached, the latter turns red; probability strong that vitriol contains sulfur oxyd though not certain yet. But remembering that copper oxyd is decomposed by charcoal at red heat, we may attempt the decomposition of this suppositious oxyd in the same way—and without much of an outlay in apparatus. Let some

FIG. 21.



vitriol be heated in a porcelain crucible until it has become white, that is, until the greater part of the water has been removed, then fill it into a tube exactly the same as before, that is, at the closed end *a*, Fig. 21. Let some splinters of charcoal be heated in a *covered* crucible at full red heat, and when cooled down introduce them into the tube at *b*; stopper the tube with cork and gas evolving tube *t*, the latter leading into basin *B* filled with water. The water filled test-tube *T* is held ready to be shoved over the end of *t*, when it may be assumed that the air has been quite driven from the apparatus by the

evolved gases. The closed end *a* is first brought to redness, then the charcoal at *b* is brought to dull redness, or any other degree of temperature we may find advisable. If the gas be decomposable exactly as copper oxyd is decomposed, we must get




but since sulfur is a *non-metal*, it is quite probable that compounds of *S* and *C* are generated, for which we must look out. The glass tube is best supported by two bricks, *P*, *P'*, stood up on edge. In order to facilitate the deposition of sulfur we protect the part of the tube beyond the charcoal with a diaphragm or screen *d*. The latter can be a slotted piece of sheet-iron or a piece of asbestus board. As soon as the charcoal becomes red at one spot, we notice a film forming before the diaphragm, which increases in bulk, forming yellow and yellow-brown drops. The water in the basin becomes turbid, milky, and a gas of peculiar smell collects in the test-tube. What causes the milkiness? What is the peculiar smell due to? These questions we shall not attempt to answer at this juncture, but place them on the calendar for future study and explanation. We let the tube cool down, cut it with file and hot glass rod at the diaphragm, and subject the yellow film to two tests. (a) We scratch out a portion of the unknown yellow substance, place it upon a piece of bright silver foil, and heat it over a flame, until the foil is barely red hot. When cold a black spot or possibly a hole will appear, where the unknown

substance was placed, the silver has formed a black compound and this is *characteristic for sulfur*. (b) We hold the tube inclined to get draught and heat the film in the flame; we get the pungent odor of burning sulfur. No doubt can remain, vitriol contains sulphur oxyd, or at any rate we can say that this gas is one of the products when vitriol is broken up by heat.

Now let us return to the original experiment which we left in order to prove the sulfur oxyd. We apply now a very hot flame (best from a Bunsen gas blow-pipe) to the vitriol. White clouds appear in the tube and roll out of the latter; the substance of this fume is evidently heavier than air. The action of it upon the mucous membrane is energetic, for a suffocating sensation is caused by inhaling it. It causes the muscles of the larynx to contract vehemently. At the same time, with the appearance of the fumes, we notice a thick liquid condensing at a little distance from the heated end of the tube. We let the latter cool down and cut it off just at the liquid ring; as the two pieces come apart another installment of white fume appears. Why? We apply litmus-paper to the liquid; it turns intensely red, but soon after turns brown, and finally the rim gets black; the paper has been charred. Deduction: The liquid possesses qualities of an extraordinary nature. The great experimenter who discovered it in the 10th century, the Moor Geber in Spain, gave it an Arabic name which was translated into Latin: *oleum vitrioli*—*the oil of vitriol*,

owing to its sluggishness in flowing like a thick olive oil, and because it produced a lubricating of the skin when rubbed between the fingers. In order to study the substance, we must design an apparatus which will allow the treatment of a considerable quantity of the vitriol, say one pound. Since we noticed that the glass became quite soft and even collapsed when exposed to the high temperature of the blow-pipe, we must look out for a material which will remain solid as well as rigid at such a temperature. Fire-clay is such a material. Its

FIG. 22.



plasticity when mixed with water makes it capable of being moulded into any desired form. When carefully dried and then equally carefully baked, it becomes hard, fire-resisting, and fairly gas-tight.

In Fig. 22 we see the section of a fire clay *retort* inside of a fire-brick lined furnace, which is heated by a gasoline flame *F*. Before filling the vitriol into the retort, we heat it in an open dish or large crucible until all the water is driven out, and until it has turned a light brick red, that is, at a low red heat, since we only care for the white, cloudy fumes



and not for the sulfur oxyd. Then we fill the material *V* through the tubulature at *S* and fasten the stopper or plug by means of pasty fire-clay. Over the neck of the retort *N*, we pass the neck of the large glass flask *R* loosely, to furnish exit for the gases. The flask we bed into a mixture of salt and broken ice (freezing mixture) contained in a basin *I* and over the upper side of the flask we spread a muslin bag *I'* filled with the same mixture. Why? In order to expose a maximum of cold surface to the fumes, having seen in the preliminary experiment that the fumes do not condense at the ordinary temperature. The lid of the furnace having been put in place, we light the flame under strong air pressure and soon the retort will show cherry redness on the outside. But since the fire-clay wall of the retort is but a poor transmitter of heat, some further time must elapse until the material *V* gets to that temperature. Then we see the white fumes pouring into the flask, falling like a foaming cataract to the bottom and also issuing at the neck of the flask. Hence the apparatus should stand under a good up draft or alongside a good down draft. The heat will be moderated or increased according to the volume of white vapor issuing from the neck *N*. If no more fume comes—say after two hours—even at a yellow heat of the retort, we stop the operation. We remove the basin *I* and the bag *I'* and wipe the outside of the flask dry. Three facts are observable: A brownish thick liquid about 50 c.c.; snow-like crystals of needle shape all over the flask; and a dense fume still filling



the flask. We pour the liquid into a large test-tube, or into a smaller flask. During this operation quantities of the white fumes arise suffocatingly from the liquid, wherever the air touches it. Query? Which constituent of the air causes this action? Azote, oxygen or the water vapor? By experimental elimination we fix the action upon the water vapor = moisture, of the air. *Without moisture no fumes.* Hence it follows that the roasted vitriol must still contain *some* water, or else we could not see the fumes inside the glass tube in our first experiment.

Two immediate investigations must now be undertaken; we must first try to get at the inwardness of the crystallized substance, and secondly, of the liquid substance.


*a.* Investigation of the *oil of vitriol*, the liquid substance. If put upon the skin, a drop will raise a blister and actually burn a hole into the flesh, destroying the tissue thoroughly.

Brought together with a drop of water a hissing sound is caused. If water be splashed or poured into a larger quantity of the oil, so much steam is generated by the evolution of heat, that the liquid is thrown violently from the vessel and has often injured the careless operator. Paper is charred into a slimy, black substance by the oil. White sugar and starch are changed into the same black substance.

The oil smells strongly of the sulfur oxyd gas, and when heated emits white fumes, until at the end a liquid results, nearly colorless, which does

not fume at the air, and does not smell of sulfur oxyd. This colorless liquid boils at  $326^{\circ}$  C. (a very high temperature), giving likewise dense white fumes to the air, when boiling, which condense into a *colorless heavy liquid*, thus differing from the white fumes of the oil, which condense into a *snowy solid*. With an arrangement as in Fig. 23, we can prove

FIG. 23.




these points. In the small retort on the left of the figure we place some 10 c.c. of the oil of vitriol, using a long-stemmed funnel, so that the neck shall not be moistened with the liquid.

At 3 we have a thermometer not reaching into the liquid ; fasten it into the tubulature with asbestos thread. The retort is held by a clamp from stand 4. At 5 we have a U-tube connected by a narrow tube with the neck of the retort through a cork. The U-tube is also fitted with perforated stoppers and stands between broken ice—or snow—in the beaker glass 6. Any liquid, condensing in the retort's neck,

will flow back. Heating with burner 2 using a very small flame, we soon get the smell of sulfur oxyd alone at 7, and later on associated with white fumes. At the same time the walls of the U-tube become frosted over with white, needle-shaped crystals. When the temperature has risen to  $300^{\circ}\text{C}$ ., we stop heating and remove the U-tube; noticing the fumes arising from the cold substance in the tube, as soon as moist air comes in contact with it. Now let us change the position of the retort by turning the swivel of the

FIG. 24.




clamp holder. The neck points downward, Fig. 24, and reaches into a dry test-tube. Raise the heat so that the liquid gets into boiling commotion. Streaks and streamers of thick liquid will run down the neck and collect in the test-tube 8. Hence it follows that by heat we can split the oil of vitriol into three parts: a gas (sulfur-oxyd); a solid (?); a liquid residue (?). The question-marks stand for: *must be found out*.

*The liquid residue.* Let it be acted upon by the metals. Bring about 2 c.c. of the liquid in a test-tube

together with a piece of copper foil. No action is noticeable until we heat, when effervescence ensues. (Effervescence means frothing due to rapid production of gas bubbles in a liquid.) The smell characterizes the escaping gas as sulfur oxyd. Further we notice the forming of a grayish, granular substance, and the copper foil has disappeared; it has become changed or converted into the white granular substance. These interesting facts lead us to a supposition or hypothesis that our liquid must contain an *oxyd of sulfur* containing *more oxygen* than the gasiform oxyd. If the latter be symbolized as  $S^nO^m$ , then the hypothetical oxyd would be  $S^nO^{m+p}$  in which symbol  $p$  denotes the mass units of oxygen which were given over to the copper enabling the latter to go into solution, through the abstraction of this oxygen from the higher oxyd to form the lower sulfur oxyd with its characteristic smell. But in order to verify the supposition we must examine the white granular product of the action. We pour off the liquid, add water to the granular residue, and see it go rapidly into solution with a pale-blue color. The solution we evaporate on a water-bath. At a certain point of the evaporation a solid begins to form. We allow the liquid to cool and a larger crop of blue crystals will form. The crystals are translucent, show an oblique symmetry, in fact resemble the crystals of our original vitriol. Hence the deduction will be rational that the crystals are *copper vitriol*. (A crystal placed upon a knife blade with a drop of water produces a bright copper spot



upon the blade.) We dry the crystals between filter paper and bring some into a closed tube. On heating, as in the original experiment, we observe water first, and the vitriol turns white. At higher heat, full redness, we note some sulfur oxyd gas, and at still higher heat, when the glass begins to melt, a ring of oily liquid (oil of vitriol). On cooling the vitriol has become *jet black*, and the black body, under the circumstances is evidently *copper oxyd*. The chain of evidence is complete. This copper oxyd must have been contained in the copper vitriol. And having put metallic copper into the liquid, the copper oxyd can only have formed by taking oxygen from a higher oxyd present. The latter is an oxyd of sulfur, because sulfur oxyd forms during the operation. We symbolize the operation or action thus :



A vitriol thus defines itself as the combination of a metallic oxyd with *sulfur peroxyd*, the syllable *per* meaning more. Both vitriols contain water, but that does not mean that *all* vitriols must contain water.

#### ACTION OF THE LIQUID RESIDUE UPON LEAD AND SILVER.

If we bring a piece of lead foil or very thin sheet lead into a test-tube with the liquid residue, there is no action at the ordinary temperature. On heating we notice a whitish film on the metal which again disappears. But on further heating there is at once



an impetuous action, with formation of sulfur oxyd gas and a very fine granular white substance. Likewise a yellow streak appears in the upper part of the tube. Lead vitriol is white and remains white when we add water, and furthermore does not dissolve even in a very large amount of water. We collect the white substance on a filter, wash it thoroughly and let it become dry in the air. In the closed tube, when heated to redness it gives *no water*, but over the gas blow-pipe it decomposes leaving yellow lead oxyd, the latter entering into combination with the glass at this high heat, forming a yellow transparent compound.

The yellow streak on the upper tube we can readily prove to be sulfur. We crack off the tube near the streak, wash thoroughly with water, dry and then heat over a flame. The yellow streak will melt and then burn with a blue flame, giving the smell of sulfur oxyd. This fact teaches us that, while under ordinary conditions a metal takes from sulfur peroxyd only enough oxygen to convert the peroxyd into the oxyd, under extraordinary conditions of stimulated chemical activity, the oxygen may be taken away altogether from the peroxyd, leaving sulfur in the free state.

We bring a piece of silver foil into the liquid residû and heat, when action sets in, the foil disappearing, without being changed into a solid residû; in other words, the silver vitriol is *soluble* in the liquid. If water be added to the cooled liquid, cautiously, a white crystalline powder will



soon fall out, but will again dissolve when much water is added, that is, the silver vitriol is soluble in water and soluble in the concentrated liquid residue, but not soluble in the moderately dilute liquid residue. Of the solution of the silver vitriol we will soon be able to make some important use. Gold is *not* attacked by the liquid residue, and hence a very important deduction follows that if gold and silver are united in the form of an alloy, they may be separated by means of this liquid residue, and are thus separated in the Mint and the Metal Refineries.

#### INVESTIGATION OF THE SOLIDIFIED OR CRYSTALLIZED WHITE FUMES.

We take now the U-tube which contains the snowy deposit. We notice the inner surface of the corks strongly blackened (same action as shown by the oil as well as the fumes, even at ordinary temperature). On pulling the cork white fumes develop (again like the oil). We let a drop of water run down the side of the glass. A hissing noise is produced, and a drop of heavy, oily liquid, runs to the bend of the U. By adding drop after drop of water, we convert all the solid, snowy substance into the brownish-colored liquid, which latter, by this time, has become burning hot. We act with this liquid upon the metals as we did with the Liquid Residue. The action is exactly the same in both cases, forming vitriol and sulfur oxyd. Hence it follows that the solid product from the distillation of the oil of vitriol, or in other words, the *white fume* must be *sulfur peroxyd*. But

then it follows inversely that if the sulfur peroxyd be converted by water into a liquid, whose action is the same as the Liquid Residue, then the latter must be sulfur peroxyd plus water. Stated symbolically this means—

White fume =  $S^nO^{m+p}$  = Sulfur peroxyd.

Liquid Residue =  $S^nO^{m+p}.H^2O$  = Sulfur peroxy-hydroxyd.

Sulfur oxyd =  $S^nO^m$ .

Oil of Vitriol =  $S^nO^{m+p}.H^2O + S^nO^{m+p} + S^nO^m$ .

And since sulfur oxyd and sulfur peroxyd are expelled from the oil by heat, we express the previous condition by saying: Sulfur oxyd and sulfur peroxyd are *dissolved* in the *sulfur peroxy-hydroxyd*, and thus form the oil of vitriol. The latter is a stable compound of the two oxyds, which cannot be broken up by boiling; it distills over at  $326^\circ C$ . without decomposition. Instead of saying sulfur peroxy-hydroxyd we will say in future *sulfuric acid*, because the substance has eminently the characters of an acid body; but the other name expresses better its make-up.


#### DIRECT PROOF OF THE PRESENCE OF WATER IN THE SULFURIC ACID.

By holding before us the actions of this substance upon the metals which we have observed we can deduce, among others, the following reasoning: (a) If lead is converted by the acid into a vitriol, *i. e.*, a combination of lead oxyd with sulfur peroxyd plus sulfur oxyd we must get lead vitriol also,

by bringing together lead oxyd plus sulfuric acid without the forming of sulfur oxyd. (b) Lead vitriol does not take up water. (c) Therefore if we mixed thoroughly an excess of yellow lead oxyd with a certain quantity of sulfuric acid, then the water must be liberated if any be contained in the acid.


In order to reduce the argument to trial, we shall mix 5 c.c. of concentrated sulfuric acid with 50 grams of yellow lead oxyd in a small wedge-wood mortar until the paste is almost powder. We fill this into a test-tube. Place the latter in a horizontal position as shown in Fig. 25, where *T* is the tube

FIG. 25.



holding the mixture *M* with a delivery tube passing through the jacket *J* fed with cold hydrant water *H*, which runs through the discharge tube *D* into the sink. Almost at once, after applying a flame gently back and forth under *M*, a condensation of mobile liquid, water in fact, takes place at *W*, much steam passes into the tube, is condensed by the cold jacket and collects in test-tube *R*. The first distillate is


nearly, not altogether, pure water. It will show acid reaction to litmus; but if one cubic centimeter be drawn out and weighed, the weight will be so nearly one gram, that for our purposes we may accept it as one gram and thus experimentally show the sameness with water. We have proved that sulfuric acid is



or in other words that it is a vitriol in which hydrogen oxyd takes the place of other metallic oxyds.

$\text{SO}^2$ . We have arrived at a stage of development when we will be able to reason out the ratio in which sulfur and oxygen are united in the gasiform sulfur oxyd. Let us make a bent (knee-shaped) tube *K*, Fig. 26. Fill it with mercury and hold it with a

FIG. 26.



proper clamp and stand in the mercury trough *P*. By heating a certain peroxyd (known as potassium chlorate) in a test-tube we can easily make pure oxygen gas and cause it to rise in the knee-tube until the mercury shall drop to an arbitrary mark *M*. We will then shove a piece of sulfur under the opening and let it rise to the surface. A piece of soft copper wire rolled into a spiral at one end will

enable us to push the sulfur to the end of the tube at 5. A strip of gummed paper can now be used to mark the new mercury level. If a flame be now brought under 5, by slow degrees to avoid cracking the glass, then 5 be strongly heated to the point of ignition of sulfur, we will suddenly see the mercury get into strong up-and-down motion. This means that union between sulfur and oxygen has taken place. Heating is stopped when the mercury has become quiet. When the apparatus has returned to the temperature of the room, we find that the mercury has risen to the mark. Hence it is evident that oxygen changes into sulfur oxyd without change of volume. Or we can say one volume of  $\text{S}^n\text{O}^m$  contains one volume of O. But how much sulfur has entered into the volume? Evidently we find this by subtracting the weight of one volume of oxygen from the weight of one volume of  $\text{S}^n\text{O}^m$ . Let the latter be  $W'$  and the former  $W$ , then

$$W' - W = S' \text{ (weight of sulfur in the gas).}$$

Now we need only to know how much one volume of sulfur weighs to solve our problem. These values have been determined and reduced to air as unity; they are

$$\begin{array}{lcl} 1 \text{ vol. } \text{S}^n\text{O}^m & = 2.210 : & \text{hence } W' - W = (2.21 - 1.105) \\ 1 \text{ vol. } \text{O} & = 1.105 : & = 1.105 \\ 1 \text{ vol. } \text{S} & = 2.200 : & = S' \end{array}$$

$$: \text{ so } S' = \frac{1}{2} \text{ S}$$

$$: \text{ because } \frac{2.20}{2} = 1.10$$

hence follows  $\text{SO}^2$  as expressing the volume ratio between the two bodies.

Thus we see that  $1\frac{1}{2}$  volumes of the free gases when combined only occupy one volume—condensation of one-third. The same was true of hydrogen and oxygen when they combined to water.

If the true numerical symbol for sulfur oxyd be  $\text{SO}^2$ , which we necessarily pronounce *sulfur dioxyd*, what is the symbol of the *sulfur peroxyd*? The answer is  $\text{SO}^3$ . You will prove this yourselves at a later stage of this course, and just simply accept the fact. The vitriols are therefore :

$\text{H}^2\text{O}.\text{SO}^3 = \text{Hydroxyd vitriol} = \text{Sulfuric acid}.$

$\text{CuO}.\text{SO}^3 = \text{Copper oxyd vitriol} = \text{Copper vitriol}.$

$\text{PbO}.\text{SO}^3 = \text{Lead oxyd vitriol} = \text{Lead vitriol}.$

$\text{ZnO}.\text{SO}^3 = \text{Zinc oxyd vitriol} = \text{Zinc vitriol}.$

$\text{FeO}.\text{SO}^3 = \text{Iron oxyd vitriol} = \text{Iron vitriol} = \text{copperas}.$

$\text{Ag}^2\text{O}.\text{SO}^3 = \text{Silver oxyd vitriol} = \text{Silver vitriol}.$

The writing of silver oxyd  $\text{Ag}^2\text{O}$  is based upon experiments and reasonings which we cannot now go into, but which will appear shortly.

*The dilution of sulfuric acid.* If we make that experiment in which we proved the presence of water in the liquid residue or the *concentrated* sulfuric acid, with great care, weighing the acid taken and weighing the water which results, we obtain in a given experiment, when we took 20.54 grams of the acid, 3.77 grams of water; hence the concentrated acid contains in 20.54 grams:  $\text{SO}^3$  equal 16.77;  $\text{H}^2\text{O} = 3.77$ . We also know that  $\text{S} = 20$ ,



that is, the sulfur as gas weighs for equal volumes twice the oxygen,  $S = 32$ , therefore

1 sulfur	$= 32 \times 1 = 32.$	Also	$2H = 2 \times 1 = 2$
3 oxygen	$= 16 \times 3 = 48.$	1 oxygen	$= 16 \times 1 = 16$
	—		—
	Sum 80		Sum 18

The numbers 80 and 18 stand for the mass units of  $SO^3$  and  $H^2O$ , or as other chemists say, they represent their *molecular weights*. The experiment gave us

$$SO^3 = 16.77 : 80 = 0.2096 : 1$$


$$H\ O = 3.77 : 18 = 0.2094 : 1$$

By dividing into these numbers the molecular weights, we change the weight numbers into molecular quantities. We get *equal* quotients and have thus established that the symbol  $H^2O.SO^3$  is the true quantitative expression for that concentrated acid, which distills without breaking up.


We take this concentrated acid and add water, mixing the two; the mixture becomes hot. We conclude of necessity that a chemical union has been effected. Let the addition of water be rational instead of arbitrary. Let the quantity of concentrated acid taken be 100 grams, which contain :  $SO^3 = 81.6$  ;  $H^2O = 18.4$ . By adding with a graduated tube or cylinder 18.4 c.c., we will have added just one other mass-unit of water; the resultant liquid will be

$H^2O.SO^3.H^2O = \textit{sulfuric acid monohydrate}$ ,  
containing in 100 parts :  $SO^3 = 68.9$  ;  $H^2O = 31.1$ .  
After this liquid has resumed the normal tempera-

ture of the room, we take of it again 100 grams, add to this 18.4 grams of water, and mixing, find again a rise of temperature, but much less than before. Unquestionably another chemical union of lesser strength of hold, the



containing in 100 parts:  $\text{SO}^3 = 58.2$ ;  $\text{H}^2\text{O} = 41.8$ . Repeating the operation a third time a very slight rise in temperature follows. Hence we conclude that the



containing in 100 parts:  $\text{SO}^3 = 49.2$ ;  $\text{H}^2\text{O} = 50.8$  is the last hydrate. Beyond this figure water is not held in chemical, only in mechanical, union, which we can very properly express by the symbol




in which aq. stands for Latin aqua = water; this is *dilute* acid.


#### ACTION OF HYDRATES OF SULFURIC ACID UPON THE METALS.

The mono- and di-hydrate act in a boiling solution the same as the acid upon lead and copper, that is,  $\text{SO}^2$  is disengaged and the vitriols form. The trihydrate does not act upon lead or copper. But if we bring this trihydrate upon zinc or iron a violent evolution of gas results. The gas, however, is not  $\text{SO}^2$ . It has a peculiar odor, more marked with the iron than with the zinc. But the gas is inflammable, and if the flame is inside a cold test-tube, water

condenses fast. The gas is, therefore, mostly hydrogen; the odor coming from other bodies contained in the iron and zinc, so-called *impurities*. The process may be represented by symbols, thus



for if the liquid be evaporated after the action, zinc vitriol crystallizes. In other words, we may describe the process thus: Zinc oxyd has a stronger tendency to form vitriol than hydrogen oxyd. The zinc therefore takes the oxygen away from the latter and thus liberates hydrogen. We may extend this idea to the former reaction, where  $\text{SO}^2$  is evolved, thus



The power inherent in “nascent” hydrogen is such that it will decompose  $\text{SO}^3$  into  $\text{SO}^2 + \text{H}^2\text{O}$ , but this power does not exist in the dilute solution. The result is just the same as if we say the metal, zinc for instance, takes the oxygen from  $\text{SO}^3$ .


Acting upon iron chips with dilute sulfuric acid, we get finally a muddy, dark-colored solution and the hydrogen gas is strongly tainted with other gas of a fetid, unpleasant odor. When the liquid stands, it gradually becomes clear with a bluish-green color, a black sediment having fallen to the bottom. We remove this by filtration and find that the dried, black substance can be burnt like coal; it is a peculiar kind of coal called graphite, and the

fetid gas is a combination of the coal with hydrogen ( $C^nH^m$ ). Zinc likewise leaves a residue, but this does not burn. We shall see at a later stage that the residue is due to certain metals and non-metals, always present in the crude, commercial zinc. But let us return to the bluish-green liquid. We will evaporate the liquid upon a water-bath until a crust forms over the surface, which signifies the beginning of crystallization. Then we set it away over night and find next morning a crop of greenish crystals. They are identical in form with the vitriol we started our experiments with; they also act in the closed tube in the same way, when heated. We have thus a direct proof that vitriol, the so-called copperas, is iron vitriol. Yet when we take the red solid residue obtained from the destructive heating of the copperas, and boil that body with sulfuric acid and trihydrate, we get a *yellow* solution, from which, by evaporation, yellow, scaly crystals are deposited. This is an undoubted vitriol, but not copperas, yet containing all the elements of the latter. Hence no other deduction is possible than the following: There must be *two oxyds of iron*, as we found a red and a black oxyd of copper, an  $SO^2$  and an  $SO^3$ . These oxyds are presumably  $FeO$  and  $FeO^2$ . But which of these is in copperas, and which is in the yellow crystals? A very neat little process of inductive reasoning will give us the answer. It has been observed at the beginning of this chapter that when a solution of copperas in water stands exposed to the air, a yellow film will form at

the surface. That this must be due to the action of oxygen, because in azote, no such film comes into being. We can accelerate the forming of the yellow precipitate by blowing air into the liquid. If we dissolve this yellow solid (after separating it from the liquid by filtration) in dilute sulfuric acid, we get, after evaporation, the same yellow crystals as we did from the red oxyd + sulfuric acid, and if we heat the yellow substance (not the crystals) we get in fact the red oxyd. Therefore it follows that copperas contains the lower oxyd, and the red oxyd is the higher oxyd. Just how the proportions of oxygen and iron stand we cannot, now, prove; but let us assume that this proportion is 1/1 for the lower and 2/3 for the higher oxyd. Then we can write



but we can explain, even now, why we get so much  $\text{SO}^2$  in the earlier stages of vitriol distillation. Namely:  $\text{SO}^3$  we saw is a strong oxidizing agent, and hence the lower iron oxyd changes at its expense into the higher oxyd.




We must of necessity get just as many molecules of  $\text{SO}^2$  as we can get of the more useful oil forming  $\text{SO}^3$ .

*Practical application* of some parts of the lesson of oil of vitriol. 1. How to prepare, on occasion of need, sulfur dioxyd ( $\text{SO}^2$ ). A 250 c.c. flask 2 is filled to about one-half with thin lathe chips of copper. The stopper funnel 1, Fig. 27, is filled with

sulfuric acid monohydrate. *B* is a burner; 4 a wash-tube partly filled with water; 5 a drying tube, one-fourth filled with concentrated sulfuric acid, and 6 a bulb tube filled with glass beads which have been moistened with concentrated sulfuric acid. Why? Because up to a certain point we have

FIG. 27.




found an extraordinary attraction of this acid for moisture. Now if the acid flows from 1 into 2 upon the chips, while heat is properly applied, a stream of pure  $\text{SO}_2$  will soon displace all the air from flask and tubes, so that presently pure, dry gas issues from 6 ready for any desired purpose. When *dry* gas is not required, the tubes 5 and 6 are left out. After the chips are used up, the flask must be cleaned out; it is best though to clean it out immediately after the



need is past, because the vitriol will become as hard as rock and the flask be much in danger.

2. *How to Generate Hydrogen.* This substance is very often required, and in large quantities, for filling a balloon, for instance. For small quantities you may use the apparatus just preceding, Fig. 27. The flask 2 is partly filled in this case with granulated zinc, and dilute acid is fed into the funnel 1. Make the acid 1:20, that is, 20 volumes of water for one volume of the acid. You will find that such a simple contrivance is all right for a small volume of gas, but not for more, and what you do get comes very fast at first, then ever slower. Why? Because the dilute acid becomes a solution of zinc vitriol, which renders the acid more and more weak. Many devices have been designed to get a better result. The adjoining figure, Fig. 28, shows my own design, which has given complete satisfaction to all those who gave it an intelligent trial. The principal part of the apparatus is a glass tube *G* drawn slightly into a neck *N*, and at the lower end into a narrow tube *O*, to which latter is fitted a stout rubber tube *R*. The latter forms a U, has a glass nozzle *P*, and thus discharges the dilute solution of zinc vitriol into the glass jar *W*. This tube *G*, which we better name the "generator" is filled with the granulated zinc. It is surrounded by a wider tube *J* which is melted at both ends together with the generator, thus forming a jacket. The latter can be filled with water through the tubulature *E*. The function of the jacket is to keep up a

FIG. 28.



KOENIG'S GENERATOR.

temperature of about  $70^{\circ}$  C. in the generator. To this end a 5-millimeter glass tube *t* has been melted into the jacket obliquely. The middle portion of the tube is of brass and is here heated by a little flame which burns from a glass tube or from a Bunsen burner. A rubber stopper closes the neck *N* and through it passes the funnel tube *F* which should be 12 inches long. The whole apparatus is held in vertical position upon two brackets which may either be attached to a portable stand *Q* or may be put against the wall under the hood permanently. The zinc is prevented from falling into the rubber tube by the "false bottom" *D* made of porcelain. After the jacket has become hot you fill the bulb of the funnel with the dilute sulphuric acid 1:20, and also fill water into the generator until it begins to run from the nozzle *P*. Now open the stopcock of the funnel so much that a drop falls from the stem every second, and soon a steady stream of hydrogen will issue from the tubulature *L*. The gas will be free from air sooner than in any other form of generator. The gas will be washed and dried as shown above. When the column of zinc has fallen by three inches, it should be filled up; but that includes a steady running for a whole day. For much use, a large reservoir bottle, holding the dilute acid, should be rigged up above the funnel, so that the acid can be drawn into the cup of the funnel by means of a syphon.

## CHAPTER IV.

### THE LESSON OF LIMESTONE.

WE have before us three productions of nature, which to the eye are very different. The first is a large crystal of *dog-toothed spar*, *calcspar* or *calcite*, quite common in our copper mines, notably at the Quincy Mine, where it is intimately associated with the native copper. This particular crystal comes from the zinc-lead mines of Joplin, S. W. Missouri. You will notice that one end of the crystal shows three lustrous faces intersecting over the edges at an angle of a hundred and five degrees nearly—a rhombohedron—whilst the other end shows six faces intersecting at alternately different angles—a scalenohedron because the faces are scalene triangles—thus producing the impression of a dog's fang. Parallel to the faces of the rhombohedron the mineral cleaves perfectly. I cleave off a piece and you see a small rhombohedron exactly *similar* to the original one. It is both colorless and transparent, and if placed upon this cross upon white paper we see the lines of the cross *double*; the two images are close together. The physicists say the mineral has double refraction and give you an explanation according to the present state of their knowledge. From this property the mineral is

sometimes called double-spar (the latter word, *spar*, is given to all minerals which are more or less transparent and show strong cleavage).


The second specimen is this whitish rock which is known as *crystalline limestone*. It shows numbers of small but splendid faces, each of which is in reality the face of a rhombohedron similar to that of the calcite: or in other words the rock is made up of innumerable calcite rhombohedrons which prevented each other from developing individual geometric independence.

The third specimen is this dull gray rock commonly known as *limestone*. In its appearance there is nothing in common with the previous specimens. But all three possess nearly the same relative weight (specific gravity), and an equal power of resistance to a penetrating steel point (equal hardness). The comparison thus far made we call *physical*, meaning therewith that all the given properties have been ascertained without destroying the identity of the original substance. Let us now act upon these materials with the powerful agents in our possession, whereby the original identity will become modified or wholly destroyed, new bodies being produced. The knowledge thus gained will be *chemical* knowledge, and will greatly widen out our horizon as to the nature of things.

1. *Action of heat.* Place a fragment of calcite in a glass tube closed at one end, and heat this end to high redness. No odor; fragment turns dull chalky. It might nevertheless be that an odorless

gas is evolved. Contrive a rig as sketched in Fig. 29, wherein  $t$  is the hard glass tube with the fragments.  $B B$  are two bricks to concentrate and reflect the heat rays upon the tube.  $T$  is a test-tube filled with water and  $t'$  a short bent glass tube attached to  $t$  by a short piece of rubber tubing. The heat may be produced by means of a blast lamp or by placing over the tube  $t$  several pieces of charcoal made incandescent and by then fanning them into combustion by an air blast or by means of an ordi-

FIG. 29.




nary fan. The heating with coal is much more satisfactory than with the blast lamp.

When the tube has come to red heat, we observe a steady current of gas bubble through the water into the test-tube. The gas is odorless and evidently not soluble in water to any considerable extent. Now we found the gas from the copperas to give a sour taste to the water, aside from the strong, pungent odor; hence we examine the present gas in the same direction. Water is not changed to the taste but blue litmus paper is slightly reddened. We say the *lime gas* has a weak acid nature. It does not burn or explode as hydrogen and does



not stimulate or increase a burning as oxygen. It is of all gases thus far encountered most like azote. But in making soap bubbles with it, the latter immediately fall to the floor; hence the gas must be much heavier than air or azote (the latter being  $\frac{4}{5}$  of the air) and moreover azote does not impart sour or acid properties to the water. Thus we are forced to the conclusion that this lime gas is a new body. But if so, we ask, is it a simple or compound body? In order to find answer to this question, let us do some reasoning by comparison: The copperas or iron vitriol is crystallized or transparent; it has been proven to be composed of two oxides + water. Calcite is crystallized and transparent, gives off a slightly acid gas and leaves a white opaque solid. Hence we will be justified in the assumption that calcite also is composed of two oxyds, one metallic, the other non-metallic.




This is an *hypothesis* (the word is the Greek equivalent for either supposition or assumption).

*Proof for the gaseous or volatile part.* If the gas be an oxyd like water (as steam) it may be possible to break it up by a metal such as zinc, or, as we saw with the oxyds of iron and copper, heat and charcoal might do it. Let us choose zinc in the form of these bright chips. It will be necessary to produce a small but steady current of the lime gas. Having seen that both dilute sulfuric acid and vinegar can decompose the calcite and the limestone, we choose


the vinegar. Why? Because the latter gives a soluble product, whilst the sulfuric acid gives a milky solution or a white mush, an insoluble vitriol. In the flask *F* (Fig. 30) bring about 20 grams of finely ground calcite or limestone with about 50 c.c. of water. Into the funnel *V* pour concentrated vinegar acid, acetic acid, because the Latin word for vinegar is *acetum*. With the help of a small flame a steady current of gas can be made for quite

FIG. 30.




a while, admitting more acid in small portions as needed. In the hard-glass tube *T* at *Z* lay the zinc chips between two asbestos plugs. The flame *L* will bring *Z* to red heat. Stand *S* with clamp *K* holds *T* in position. But since we want to prove an oxyd, it is evident that such proof would become impossible if the gas were to enter *T* at once, for the gas is charged with water vapor and we know that water will yield oxyd when brought together with red-hot zinc. We therefore interpose the test-tube *D* which is partly filled with concentrated sulfuric

acid, through which the gas will have to rise in bubbles. We do this because we found that oil of vitriol absorbs water with very great energy. It is a *dryer*. Had we not observed well and noted this property we would now stand before an impassable obstacle. But going as we do, the discoveries come apace with their immediate practical application. Now we begin the generation of the gas. The latter, being heavier than air, forms a steadily thickening layer over the liquid in *F*, driving the air before it and out of the entire apparatus. We keep on patiently until at least two volumes equal to *F* have been generated to make quite sure that the air is completely displaced. For if any remain our results could not be conclusive, since some zinc oxyd would surely be formed, whether the lime gas were an oxyd or not. Zinc is coming to redness; a white cloud appears on the glass. But in order to find what becomes of the gas let a rubber tube be brought under a test-tube *G* filled with water, and it will be seen that the tube soon fills with a colorless, inflammable gas. The zinc is oxydized. Now we may assume that the inflammable gas is the element Y or a lower oxyd of Y and symbolize thus:



or



This alternative brings us up against the wall once more. Unless we shall happen to discover a metal with greater attraction for oxygen than the zinc, we shall not be able to prove directly that the com-

bustible gas is an element or a lower oxyd. Let us not despair; of wonders and signs of wonders there is no end.

*Study of the residuum.* In order that this residue may be as much freed from the gas as possible, let us pour it from the glass tube into this platinum crucible. This latter being infusible we may concentrate upon it a much higher heat than was possible in the glass tube. Lifting the lid we see the pieces emit a white glow, whilst the metal of the crucible is only yellow. We say the lime is highly *incandescent*. The pieces still show the rhombohedrons of cleavage; there is even luster upon the faces. The volume is the same but the weight has decreased 44 per centum. Cohesion has decreased. This residue is known as *burnt lime*—*caustic lime* (“caustic” being merely the Greek for burnt). It has been known for ages to both civilized and savage peoples. Since limestone forms the surface rock for many square miles in large tracts of country all over the earth, the first burnt lime was made when the first man—living in a limestone country—made a rough fireplace with the pieces of rock around him. According to our hypothesis burnt lime ought to be an oxyd  $X^nO^m$ , the oxyd of a metal  $X$  whose properties we do not know. For if we try upon this lime the same agents which yielded us the metal from the oxyds of iron and of copper, namely intense heat (by the blow-pipe) and charcoal, our trial will end in failure; the white material remains quite unchanged. What was said of

the final nature of the element Y above, applies here in regard to X. Let it stand for the present as X, or since we called Y the lime gas, let X be the *lime metal*. We may even designate by the symbol Ca (the first letters of the word calcite), since we designate iron by the symbol Fe (the first letters of the word *ferrum*); yet ever bear in mind that it is a suppositious, a hypothetical simple body of whose properties, in the first state, we are ignorant. This ignorance does not prevent us from studying the behavior, the properties of the supposed oxyd.

*First towards water.* Let a drop of water fall upon some of the caustic lime, a hissing noise ensues; a slight cloud of steam arises; the lime swells up and falls into an extremely fine powder: *Flour of lime*. The flour can be dried at steam heat and yet this dry flour will yield water in the closed tube, at red heat, and lose 24.3 of its weight. After cooling moisten the lime and it will hiss with water as before and fall into flour. Hence we say the burnt lime has a very strong affinity for water, and will form with it a true union, a chemical compound, a *hydroxyd*, very much as the sulfur oxyd SO which forms the hydroxyd we called sulfuric acid. Deduction: *Both metallic oxyds and non-metallic oxyds form hydroxyds.*




Flour of lime = lime hydroxyd =  $\text{Ca}^n\text{O}^m.\text{H}^2\text{O}$ .

Adding more water the lime hydroxyd turns into a *white paste*: *Slaked lime*. If we add much water, shake and stir thoroughly and then let stand, we notice that apparently the whole of the white paste



will settle, leaving a clear liquid above. Deduction : Lime hydroxyd is very little—if any—soluble in water. However, the water has a decided taste, and turns reddened litmus paper to *blue*. Hence it is an agent, and we call it *lime water*. By evaporating 100 c.c. in a weighed dish, we obtain a residue of hydroxyd of 0.1 gram—1000 lime water hold 1 hydroxyd.

*Second, towards acids.* Make a  $\frac{1}{10}$  p. c. solution of sulfuric acid and of vinegar or acetic acid. Because sulfuric acid has a specific gravity of 1.83, we will require 0.054 c.c. to give us 0.1 gram ( $1.83 : 1 = 0.1 : 0.054$ ), that is, just one small drop for the 100 c.c. of water. The strongest acetic acid has sp. grav. 1.05, nearly the same as water. Two small drops of it will give the strength wanted with 100 c.c. of water. Pour 25 c.c. of each of these very diluted acids into 2 beaker glasses, add a little *litmus* solution to each, which will produce red color. Now slowly add clear lime water from a graduate into the first beaker glass. All at once the red color changes to blue, when a certain number of c.c. have been added. The same with the acetic acid. We deduce: The two bodies of opposite action to litmus saturate each other so that a *neutral* body results, the neutral body is the *salt*. Let M stand for any metal, and N for any non-metal, then





The three conceptions are fundamental in chemistry. Base, acid, salt. Yet neither base nor acid is to be understood in the absolute sense. Two metallic hydroxyds may act as base and acid towards each other, or in other words one is more basic than another. We shall find examples as we proceed. This much, however, impress upon your mind: An oxyd is rarely an active agent; in order to make one oxyd act upon another oxyd strong external impulse is needed, such impulses being heat and electricity. The internal activity becomes manifest towards the surroundings with the formation of the hydroxyd. This will become clearer in the next chapter.

Allow a portion of the liquid, in which you have saturated the lime hydroxyd with the sulfuric acid, to evaporate on a watch crystal. Groups of crystals will be formed, colorless, needle-shaped. Examine them with a microscope. They are often stellar, that is, radiating from a centre. When heated these crystals will loose water readily; they are  $\text{CaO} \cdot \text{SO}^3 + 2\text{H}^2\text{O}$ —a vitriol in which two molecules of loosely-bound water of crystallization are tacked on to the salt proper as in iron vitriol and copper vitriol.

*Action of the lime gas upon the lime hydroxyd.*—Lime water is the  $\frac{1}{10}$  per cent. solution of the lime or calcium hydroxyd in water. Generate the gas as above and let it bubble through some of the solution in a test-tube; a turbidity appears at once which gradually turns to a milky white color and white

sediment. Filter this upon a small paper filter, dry it. It has no taste. Heat some of it to yellow heat upon a bit of sheet-iron (platinum preferable but too costly); after cooling transfer it to a slip of reddened litmus paper and moisten with one drop of water. Observe that it slakes and the litmus turns blue under the white mass. Place another minute quantity of dry precipitate upon a watch glass or plain glass slide and examine, after moistening it, with a high power of the microscope. Minute but perfect transparent rhombohedrons appear. Hence deduction: *Lime gas converts lime hydroxyd into calcite* from which we started. High heat breaks up the combination of the two oxyds; at low heat (temperature of room) they recombine. Conclusion: Very high heat always counteracts the chemical attraction, tends to separate the minute mass-units. Now let the lime gas bubble through another portion of the lime water and leave it for some time, being called away. On returning we find the original milkiness all gone; evidently the calcite has been dissolved in an excess of gas. Boil the clear solution and shortly milkiness as well as sediment reappears. Conclusion: Boiling heat expells the solving excess of gas, the dissolved calcite being itself almost insoluble (one part in 250,000 parts water), must precipitate. This is the reason why the *hard water* of limestone regions always gives a sediment after boiling and the boiled water becomes *soft*. Here is some lime water which has been standing for quite a while in the open beaker glass.

Note that a film has been forming on the surface. The film shows *iridescence* in strong sunlight. Remove some to a glass slide and you will find with a high power the identical rhombohedrons. Deduction: *If calcite forms under these conditions it is evident that the lime gas must be part of the atmosphere.* And since the film forms much more rapidly in a crowded room than outside, we must reason that lime gas forms part of the effluvia or gaseous emanations of the human body. Another important side-gain is that the hardening of *mortar* must be owing to the absorption by the slaked lime of the lime gas in the air. Mortar is a mixture of 80 to 85 parts of sharp sand with 20 to 15 parts of quick-lime in the slaked condition.

All the actions observed are identical for calcite, crystalline limestone and common limestone; their substance is identical, though their look is very different. This must be noticed, however, that the gas evolved from common limestone possesses a fetid odor which is owing to oily matter often contained in the limestone.

*Summary of limestone lesson.* Discovery of a gaseous oxyd whose non-metal Y is at present unknown. Of a metallic oxyd (probably) whose action towards the acids is equal to the oxyds of iron and copper, though we do not know the metal contained therein. The oxyd differs by its tendency to form hydroxyd and by its action on litmus paper, which we call *basic* action.

## CHAPTER V.

### THE LESSON OF WOOD ASHES.

WE have before us the familiar and homely material—ashes. There are evidently two kinds. For in the one we find fragments of partly coaled wood, and in the other hard, glassy nodules known as clinkers. The first is the remnant from burning wood, the second the remnant from burning coal in a kitchen range. Why should I bring these materials before you? They look unpromising enough. Because I find, upon trial, that the wood ashes produce a strong, biting taste similar to that of slaked lime, whereas the coal ashes show no taste whatever. There must be something in the wood ashes outside of the ordinary earth calling for investigation. The Roman historians tell us that when their armies came in contact with the Teutonic tribes on the Rhine, they found it a custom among these for the women to do up their hair with a kind of ointment, and this they made up from fat and wood ashes. It was indeed what later on was called *soft soap*. Whilst these people were at that time, 2,000 years ago, barbarians much like the American Indians, they had the spirit of investigation stalking among them, and this spirit is stalking among them now. When linen and woollen

cloths had superseded the skins of animals, the original hair ointment was found to possess excellent cleansing properties ; the manufacture of soap became a separate trade, and wood ashes came to be an article of commerce. The early pioneer in America, for many years, had nothing to exchange for groceries and other store goods but the ashes which he collected from burning out his clearings in the forest. However, the valuable part of wood ashes is only about 30 per cent.; the storekeeper had no use for the 70 per cent. of waste. The farmers were made to extract the valuable portion with hot water, to strain the liquid through canvas, to boil down the liquid to solidity, in iron kettles or *pots*, hence the commercial product came to be called *potash*. This material is not yet pure ; it is of brown color, and contains other soluble parts of the ashes. It undergoes a refining process, becomes white, and goes under the name of *pearl-ash*. However, wood has become so scarce everywhere that it can no longer be burnt for the sake of the ashes. How means were found to replace it successfully, in more recent times, we shall see in a following chapter.


*Investigation.* We have boiled down the liquid—the *lye*, “lie” (pronounce lee) in French, “lauge” in German. In Germany potash was called “laugen-salz”—salt made from the lye.

First let us see how the potash acts at high heat. We place some in a hard-glass tube closed at one end. First some water condenses in the upper tube.



At red heat the substance becomes liquid, and then we observe small gas bubbles arising from the contact between the liquid and the glass, the mobility is changed to sluggish flow. Question: Have the escape of gas bubbles and change of flow anything to do with an action upon the glass? or is it inherent in the potash itself? To answer, let the glass tube be substituted by a metallic vessel, say an iron crucible. The potash melts as before but no bubbles come; the liquidity remains the same. It follows that escape of gas is caused by interaction or reaction upon the glass. Potash is fusible at red heat unchanged: remember this important fact.


Act with sulfuric acid or acetic acid upon the dry potash and upon its solution in water. In either case there is strong effervescence. We apply the same procedure to the examination of the gas which we used with the lime gas. We find the gas in all its actions like the lime gas. Moreover, a white granular salt falls out when sulfuric acid decomposes the potash, a vitriol. Therefore we will be justified in the assumption that potash is composed of two oxyds:




in which P as the first letter of potash, stands for a metal as yet unknown to us, because its vitriol is unlike the known vitriols in its crystal form, unlike also as to solubility, and especially unlike in this, that heat does not break it up; the vitriol  $\text{PO.SO}^3$  stands the heat of the blast-lamp, as I here




show you in the glass tube. The vitriol is more like the lime vitriol than like the iron and copper vitriols; since the action of potash and lime hydroxyd are both *basic* to litmus. Since heat neither breaks up the potash nor the potash vitriol, how shall we get at the hypothetical oxyd PO? Let us reason: We found calcite insoluble, but convertible into oxyd. Suppose we bring the solution of potash together with the lime oxyd in this test-tube, and let this be represented by the scheme:



CaO will become slaked lime with the water, we will then get




We notice turbidity at once, then flocculency, then a granular precipitate. We filter. The filtrate with sulfuric acid does *not* give gas, but a granular salt falls out slowly as the liquid cools. The action must, therefore, have been




the non-metallic oxyd  $Y^nO^m$  has gone to the lime, and only the hydroxyd  $PO.H^2O$  remains in solution. The filtrate causes deeper action on the skin of the fingers, on litmus paper and on the tongue. It also follows that  $PO.H^2O$  is much more soluble in water than  $CaO.H^2O$ .

*Potassium hydroxyd, caustic potash, caustic potassa, potassium hydrate.* The first of these names I want you to use. The second and third names are older





and still used in the drug trade ; the third name was the current scientific name and is used by the majority at present. But I want it to apply to a separate conception. To make myself clearly understood we will return to the action of water upon oil of vitriol. Represented symbolically we have there to start with :







Adding water, little by little, there is much heat, also hissing noise, this lasting until the  $n\text{SO}^3$  have combined with  $n\text{H}^2\text{O}$  and we have now only



our concentrated sulfuric acid—the true hydroxyd. But when you add to this more water, both being at ordinary temperature—the liquid warms up and can even reach boiling heat. This heat means *more chemical union* and may be scheduled




and similarly :



The  $n$ th hydrate may be, in fact, what we would

otherwise designate a *dilute water solution* of the hydroxyd. Returning to the matter immediately before us, we evaporate the water solution of  $H^2O.PO$  to dryness in an iron, copper or silver dish. Glass and porcelain are strongly attacked. Prove this statement by using a small beaker glass and a porcelain crucible; they are not destroyed but lose the lustrous surface and some of their material enters the liquid. After the mass has become dry at boiling-point of water, heat over an open flame. Soon fluidity will occur, more steam will be given off; at red heat white vapors appear and, using a small portion, it will slowly disappear: the hydroxyd is *volatile* at red heat. But how do we know that this material is hydroxyd still; why is it not the oxyd, when the lime hydroxyd looses its water so readily at red heat? Revolving in our minds all the actions heretofore performed, we remember that both iron and zinc decompose water at red heat; it may even do so when the water is united strongly to another oxyd. The rig will be simple. A short piece of hard-glass, thick-walled tubing closed at one end; a perforated stopper, a narrow tube drawn into a fine opening and inserted into the stopper will probably suffice. We introduce a piece of the problematic hydroxyd with some zinc shavings, insert the stopper, hold the tube by means of a clamp in inclined position and apply heat. With the melting of the hydroxyd, gas bubbles appear, and ere long the mass will want to froth out of the tube, the escaping gas burns, the flame deposits

drops of water against a cold dish—the gas is hydrogen. In symbols the action is



After cooling we find that the mass is quite soluble in water, all but some remaining zinc chips. Here is one example of two metallic oxyds combining to form a salt, because one, PO, is more basic than the other, ZnO. Thus, whilst proving the hydroxyd, we have incidentally discovered that this latter is a most powerful agent, rivalling the sulfur hydroxyd. It corrodes the skin rapidly; it destroys paper and sawdust; it dissolves wool and hair, horn chips and many other bodies. In these actions many interesting and useful new substances are formed, some of which we will inquire into hereafter. We find ourselves now in possession of the two most powerful agents  $\text{H}^2\text{O.SO}^3$  and  $\text{PO.H}^2\text{O}$ . Acting upon each other they produce the neutral vitriol  $\text{PO.SO}^3$  and water. Acting separately, they lend us their latent power.

*Action of calcite, of potash, of lime hydroxyd, of potassium hydroxyd and their hydrates upon the water-soluble vitriols of iron, zinc, copper.* The vitriols are in dilute solution ( $\frac{1}{100}$ ).

1.  $\text{CaO.Y}^n\text{O}^m + \text{CuO.SO}^3 + \text{boiling heat}$ , escape of gas, green precipitate.  
     +  $\text{FeO.SO}^3$ , slight brownish precipitate.  
     +  $\text{ZnO.SO}^3$ , no precipitate.

2.  $\text{PO.Y}^{\text{n}}\text{O}^{\text{m}} + \text{CuO.SO}^3$ , at ord. temp. blue precip.,  
     at boiling turns black.  
     +  $\text{FeO.SO}^3$ , at ord. temp. light precip.,  
     at boiling turns dark.  
     +  $\text{ZnO.SO}^3$ , at ord. temp. white precip.,  
     at boiling remains white.
3.  $\text{CaO.H}^2\text{O} + \text{CuO.SO}^3$ , precipitate at ord. temp.  
     and complete at boiling heat.  
     +  $\text{FeO.SO}^3$ , precipitate at ord. temp.  
     and complete at boiling heat.  
     +  $\text{ZnO.SO}^3$ , precipitate at ord. temp.  
     and complete at boiling heat.
4.  $\text{PO.H}^2\text{O} + \text{CuO.SO}^3$ , first blue precipitate which  
     turns black.  
     +  $\text{FeO.SO}^3$ , light green precipitate  
     which turns black.  
     +  $\text{ZnO.SO}^3$ , white precipitate which  
     dissolves in excess.

It can easily be proved that the precipitates formed under (2) are the combinations of  $\text{CuO}$ ,  $\text{FeO}$ ,  $\text{ZnO}$  with  $\text{Y}^{\text{n}}\text{O}^{\text{m}}$ , whilst the  $\text{PO}$  combines with  $\text{SO}^3$ .  $\text{CuO.Y}^{\text{n}}\text{O}^{\text{m}}$  turns black on boiling, because the  $\text{CuO}$  is not very basic and cannot hold on to the non-metallic oxyd when the shattering power of heat-waves pounds upon the compound. The same is to be said about  $\text{CuO.H}^2\text{O}$  under (4) the  $\text{CuO}$  cannot hold on to the  $\text{H}^2\text{O}$ . The white precipitate (4) in zinc solution— $\text{ZnO.H}^2\text{O}$  dissolves in excess of the agent because the soluble salt  $\text{PO.ZnO}$  forms, thus

First:  $\text{ZnO.SO}^3 + \text{PO.H}^2\text{O} = \text{ZnO.H}^2\text{O} + \text{PO.SO}^3$ .

Second:  $\text{ZnO.H}^2\text{O} + \text{PO.H}^2\text{O} = \text{PO.ZnO} + 2\text{H}^2\text{O}$ .

Of iron there are two vitriols—a green and a yellow, the latter produced by acting upon iron with the hydroxyd  $\text{H}^2\text{O}.\text{SO}^3$  when  $\text{SO}^2$  escapes instead of hydrogen.

The green vitriol contains the oxyd  $\text{FeO}$ , the yellow vitriol contains the oxyd  $\text{Fe}^2\text{O}^3$ . The two vitriols act differently on our four agents. But the one important action is that of *calcite* for  $\text{FeO}.\text{SO}^3 + \text{water} + \text{CaO}.\text{Y}^n\text{O}^m = \text{no precipitate}$ .


$\text{Fe}^2\text{O}^3.3\text{SO}^3 + \text{water} + \text{CaO}.\text{Y}^n\text{O}^m = \text{brown precipitate}$ , the action being slow at ordinary temperatures, rapid when heat is applied. Hence we have here a means of separation for the two oxyds of the same metal. (Details for this in the Chemistry of the Metals.) Likewise if the vitriols of Cu, Fe, Zn were mixed together (the iron vitriol being of the green kind), we would be able to separate the oxyds—for (Cu and Zn) vitriols precipitate by calcite or chalk and heat, separating the iron. The precipitate boiled with  $\text{PO}.\text{H}^2\text{O}$  will leave the  $\text{CuO}$  as a solid and take the  $\text{ZnO}$  in solution. Our wealth is increasing as we go along.

*The metal potassium.* Having seen that the oxyd  $\text{PO}$  is not obtainable, but only the hydroxyd  $\text{H}^2\text{O}.\text{PO}$ , and that this latter is quite volatile, their remains only one raw material, the potash. Though it be a salt, the peculiar nature of its non-metallic oxyd makes it possible to be broken up by coal, at a white heat. Cut a piece of  $\frac{3}{4}''$  or  $1''$  gas pipe  $P$  (Fig. 31) six inches long, fit on a cap  $C$ , an elbow  $E$  and a pipe  $R$  of same length, all joined by thread.



Fill *P* with a mixture of charcoal-coated iron chips, 30 grams of fused potash, 10 grams of burnt lime. Screw on *E* and *R* and set into furnace *F*. The latter is heated with gasoline burner *B*, but a good wind furnace and coke will give a suitable heat also. In order to prevent the pipe from burning through, it is well to coat it over with several coats of *char-motte* or *braise*, a mixture of three parts of ground brick and one part of fat fire-clay. The pipe *R*


FIG. 31.




is kept cool by a wrapping of blotting paper upon which water drops from the hydrant or spigot. A cork stopper 5 with glass tube permits the gases to escape. The burnt lime was added to the charge to make the potash less fusible; the iron to hold the charcoal down, preventing it from floating to the top. This charcoal-coated iron is made by heating iron chips and sugar together in a covered crucible until no more gases escape. After yellow heat has been reached, combustible gas will appear and burn

at 5 with either a purplish or yellow flame. Keep up the fire for an hour, then cool down. A gray and black loose mass will be found in *R*. If some be thrown into water a hissing will be heard and for a short time a fine purple flame. The black material is a mixture of small metallic pellets and a spongy substance. By returning this mass to a smaller apparatus, of exactly the same form, the metal can be distilled from the sponge and appears then almost pure in *R*. The electric current also decomposes the hydroxyd  $\text{PO.H}^2\text{O}$  thus: The metal deposits on the

FIG. 32.



negative pole (cathode) whilst hydrogen forms at the same pole and oxygen escapes at the positive pole.




It is difficult to keep the metal from burning up again in presence of the oxygen surrounding it. The difficulty may be avoided by pouring some mercury into the crucible (Fig. 32) and some of the third potassium hydrate over it. If now the cruci-


ble be made the negative pole and the positive pole be a platinum wire, the metal potassium in the moment of liberation combines with the mercury, alloys with it, and thus is kept from the air. The product is the potassium amalgam  $\text{Hg}^n\text{P}^m$ . We place the latter in a small glass retort and distill off the mercury at  $300^\circ \text{C}$ . Potassium remains as a liquid. At a red heat it also becomes volatile and will fill the flask as a fine green vapor. The retort must be kept filled with hydrogen to keep out the oxygen of the air.

*Physical properties of the metal.* Potassium has a silver-white color, strong metallic lustre. But in presence of air tarnishes at once, becoming covered with a gray film. Therefore, the metal must be kept under a liquid which does not contain oxygen—such as kerosene. The metal is soft, like fresh putty at ordinary temperature, becomes brittle below the freezing point; that means it crystallizes, appearing in tetragonal pyramids. It melts at  $62.5^\circ \text{C}$ .; at red heat,  $730^\circ \text{C}$ ., the liquid boils like water; the vapor is green. Its specific gravity is 0.865 (water 1), hence, it floats on water, but sinks in kerosene, sp. gr. 0.76—0.78. The specific heat or heat capacity is 0.166 (water = 1).

*Chemical properties.* Potassium decomposes water with great energy, because of all metals it has the strongest affinity for oxygen.



but since an attraction exists between the oxyd and the water the action really is




For instance, if you wish to ignite coal-oil which floats on water, you would just throw a piece of potassium on the water, and the oil would be on fire at once.

I said potassium had the strongest attraction for oxygen of any metal. Why, then, were we able to dislodge it by iron and by charcoal? The answer is found in that potassium is so easily *volatilized*. We could not separate the calcium from the lime oxyd, because the calcium is *not volatile*.

*Proof of the nature of the unknown Y in lime gas.* Once in possession of potassium, we will try it upon the lime gas, which zinc only changed into combustible gas (see above), and of which we remained

FIG. 33.




doubtful whether it was Y or a lower oxyd of Y. Let the lime gas be generated from calcite, chalk, or limestone by means of the acetic acid as before. Let the gas bubble slowly through H<sup>2</sup>O.SO<sup>3</sup>, Fig. 33, thence pass it into the combustion-tube t, in

which a piece of potassium has been placed at *P*. If the dish *D* contains the 6th potassium hydrate, as also the test-tube *t*, then there will be perfect absorption of the gas as soon as all the air is expelled from the apparatus; because we know that the potassium hydroxyd as well as the different hydrates, absorbs the gas energetically in order to become potash  $\text{PO.Y}^{\text{n}}\text{O}^{\text{m}}$ . Now let the potassium be heated with the lamp *L*. It melts, spreading over the glass and forming a perfect metallic mirror. Then it ignites and burns with the characteristic purple flame almost the same as in air, a white smoke developing. Soon the potassium becomes incrustated with a white material and a black material. All the gas becomes absorbed for a while, then reappears in the tube *t*. This designates the end of the action. After cooling, we remove the mass from the tube. The white portion gives all the actions of potash: Strong, bitter taste, easily soluble in water, and gas evolution with acid. The black portion is not soluble in water; we separate it by filtration. It looks like lamp-black or soot. Under the microscope we find it to be of brown color and translucent with yellow or brown color. At red heat it burns and disappears. If the burning be done in an open tube, one end leading by rubber tube into lime water, Fig. 34, and air being sucked through by means of an *aspirator*, then the lime water will become milky; the white sediment being rhombohedrons. Hence we deduce that the black substance is the Y in the lime-gas, because from it

comes lime-gas by combustion. Now we have many times observed that charcoal burns and disappears, leaving a slight residue of ashes, and we naturally will ask: Is there a communion between the black Y and charcoal? To satisfy the query we place a splinter of charcoal in the tube *T*, and fresh lime water in the test-tube. As the charcoal burns, the lime water becomes milky with calcite. Hence we

FIG. 34.



will make the deduction that charcoal is either wholly or partly made up of the black substance Y, and we resolve the Y into C, which is the first letter of the Latin word *carbo*, the equivalent of the English charcoal. With characteristic inconsequence the English language makes *carbon* out of *carbo*. Our lime-gas becomes now  $C^nO^m$ ; the calcite becomes  $CaO.C^nO^m$ ; potash  $PO.C^nO^m$ , pronounced *calcium carbonate*, and *potassium carbonate*.

The symbol for potassium should be P as we adopted it. But here again the English chemists are inconsequent in choosing K, which is the first letter of the Arab word *kali*, which means "burnt," but was given by the Arab chemist Geber to our potash. This word was taken up by all chemists



with the Arab prefix "al," *i. e.*, *alkali*, to mean any body which shows the essential action of potash, these actions being called *alkaline* actions. *A solution is said to be "alkaline" when it turns red litmus to blue.* German and Swedish chemists call the metal potassium *kalium*, English, American, and French chemists stick to potassium, but they accept the symbol K.

## CHAPTER VI.

### THE LESSON OF COMMON SALT.


COMMON salt is before us in three forms. 1. *Rock salt* as mined at the mouth of the Mississippi, in Canada, and other places in America. It is beautifully transparent; colorless or colored, sometimes intensely blue. It is found as large, perfect cubic crystals, and again as immense, solid, irregularly shaped masses like ice; or as fine grained, snow-white or dirty gray, or yellow, or red masses. It cleaves perfectly in three directions at right angles—cubical cleavage. It presents slight resistance to the knife or to the drill. It has a strong taste; is readily dissolved by water. 2. Salt from evaporation of salt springs and salt wells, in snow-white cubic crystals, which are all hollow on the faces. 3. Salt from evaporation of sea-water. It is certain that all the deposits of salt, found in nearly all geological formations, were at one time dissolved in the sea. It is likewise certain that salt springs and wells dissolve the rock salt they find in the rocks, so that, in the end, our three kinds of salt are practically only one kind, *sea-salt*. Man and animals crave salt, their bodies must have salt or die. Hence all the languages derived from some primitive language have nearly the same name for

the substance: Greek: *hal*, Latin: *sal*, French: *sel*, German: *salz*, English: *salt*. The plant-eating animals get some salt with the grass and leaves, but it is known that they will travel 100 miles and more to get a salt spring or salt-lick, to satisfy the craving for salt. Carnivorous animals get their salt in the blood of the grass-eaters on which they prey. Man takes his from both plants and herbivorous animals. Bloody wars have been fought for the possession of a salt spring or salt mountain.

The United States are well supplied with salt, and need not go to the evaporation of sea-water. Germany is very rich in salt; exports much to other less favored nations. Where the sun's heat can be made use of for evaporation, the sea-water gives very cheap salt.

*Investigation.* Heat does not break up the salt. Holding a piece of salt in the flame we see that it melts quickly and colors the flame a deep orange-yellow. Heating the salt in a glass tube we notice *decrepitation* (enclosed mother liquor escaping explosively), then a slight water condensation, then melting, then boiling and forming of a white sublimate. No odor, nor gas.

The sublimate shows itself made up of cubic crystals, tastes and acts the same as the original salt. Deduction: Salt is volatile without decomposition. It may be a simple body as far as heat action shows. But not so by its behavior towards electricity. We place salt upon a platinum crucible lid, the lid being in contact with a sheet of copper and the



latter forming the positive pole of the current. As soon as we bring the other pole wire in contact with the salt, the salt melts and an evolution of gas ensues accompanied by a very strong and peculiar odor. The same odor is produced if we mix salt with blue vitriol and heat in closed tube. But let us try our two strong agents: Sulfur hydroxyd and potassium hydroxyd.


$\text{Salt} + \text{K.O.H}^2\text{O} + \text{heat} = \text{no gas, no apparent effect.}$

$\text{Salt} + \text{H}^2\text{O.SO}^3 + \text{heat} = \text{copious evolution of colorless gas possessing a strong, pungent odor, and reddening blue litmus.}$  Let this gas be called *salt-gas*. We let the gas pass into water: it is absorbed eagerly, the solution becoming warm. Looking at this phenomenon we cannot deny that it is similar to the action of  $\text{H}^2\text{O.SO}^3$  upon calcite, hence that salt must be composed of two oxyds



in which neither the metal M nor the non-metal N is known to us. With this supposition as a guiding thread, we proceed with experiments until we shall have established the truth or falsity of the conception, and recognized the properties of both M and N.


*Salt gas, spirits of salt.* If our assumption be true, the production of salt gas must occur thus



the salt gas must be the hydroxyd of the non-metal oxyd  $\text{N}^{\text{SO}}$ . The gas contains *hydrogen*. For if we expose zinc to it hydrogen is evolved at once. Let


*F* be a flask holding about 500 c.c. Let it be fitted as in Fig. 35, with funnel, stopper and escape tube, all standing on a tripod so that heat may be applied. Let *C* be a bulb tube filled with cotton, to retain any particles carried over by the gas. *T* is a hard-glass tube with two perforated stoppers. *B* a bulb tube providing for any liquid mounting back from *D*. *t* is a test-tube to receive any gas which may be

FIG. 35.



generated. Remove stopper *S'*, introduce 50 grams of salt into *F* at *S*, fill funnel with  $\text{H}^2\text{O}.\text{SO}^3$  and drop the latter slowly into *F* for some time, until upon adjusting *S'* the gas bubbles are all absorbed in the water in dish *D*. Open at *S'* and introduce zinc at *Z*. Close *S'* and adjust tube *t*. The speed of generation of the gas is to be judged by the frothing of the salt. *T* being cold (flame *L* not having been lit) at start, will now warm up: zinc decomposes

gas at ordinary temperature, the combination produces heat. Prove the gas in test-tube is hydrogen by showing it inflammable. At *Z* a new body has been formed which is fusible, proven by means of lamp *L*. By withdrawing it from the tube it is found to be soluble in water. (Prove solution by testing with potash and also with  $\text{KO.H}^2\text{O}$ ), therefore this substance cannot be zinc oxyd for this latter is neither fusible nor soluble in water; it must be a combination of zinc with the unknown *N*. Acting upon it with  $\text{H}^2\text{O}$ .  $\text{SO}^3$  the pungent gas forms same as with the salt. Nevertheless the combination may contain oxygen. As we know the great avidity of potassium for oxygen, let us act as follows: We bring into a test-tube a part of the unknown compound  $\text{ZnN}^s\text{O}$ ; we fill the tube with hydrogen, excluding thus any air; we then add a piece of bright, carefully cleaned potassium and apply heat, hydrogen still passing in. The action would have to be thus:




It will evidently not matter which of the reactions ensues, or whether all three occur at the same time. The essential point is the forming of  $\text{KO}$ ; because if water be now brought into contact with the mass,  $\text{H}^2\text{O.KO}$  will be formed and will cause the change in litmus from red to blue. The experiment will give true information only, if  $\text{ZnN}^s\text{O}$  be in excess, for otherwise *K* would be left and in contact with



water would give the hydroxyd + hydrogen. The experiment must be made with much judicious care, and will prove that KO is not formed and, hence, that *oxygen is not present* in the compound and therefore not present in the salt gas. Our preliminary supposition was *wrong*: Salt gas must be  $N^sH$ . How are we to set free  $N^s$ ? In the case of the limestone gas we were successful with potassium because the C (carbon) does not combine with potassium; but in this instance, when we find the unknown  $N^s$  to combine with zinc, it seems more than likely that it will combine with potassium also. However, let the trial be made. The apparatus, Fig. 35, will be quite suitable. We simply substitute a piece of potassium at Z in place of zinc. We note a strong action even at ordinary temperature, a white substance forming and hydrogen evolving. The white substance forms cubes and octahedrons, it tastes like common salt; hence we conclude that the metal M in salt must be similar in its nature to K. Whilst this is welcome information, it is not what we are looking for. The unknown *is not set free*.


Another train of thought is needed; we know the unknown to be a hydrogen combination. Now the metals having failed us, let us try the non-metals. Of these we know oxygen, azote and sulfur, and of these oxygen seems the most active. We reason along the scheme



Mixing oxygen and salt gas at ordinary tempera-

ture has no effect as seen in this cylinder, which holds the mixture; therefore let the mixture, one volume of each, be passed through the glass tube *T* (Fig. 36) and let asbestos *A* (mineral wool) partly fill the tube. The asbestos being brought up to redness, the gas mixture slowly passes through it in measure as the cylinder *C'* is raised and the cylinder *C''* is lowered. While *C* fills with mercury the pipette *P* fills with a gas of pale-green color.

FIG. 36.



This gas we shall hereafter name *chlorine*, (*Cl.*) from Greek *chloros* = green. For consistency's sake the name should be *chlorium*—as *hydrogenium*, *ferrum*, *stannum*, *potassium*. (The great chemist, Berzelius, never gave up his belief that this chlorine was a compound and not an element, that it does contain oxygen, but that we are not able to separate the two. The demonstration we have gone through above leaves very little doubt that the salt gas and hence chlorine do not contain oxygen.)


*Chlorine, physical properties.* A green gas of

peculiar, very strong odor; offensive to the mucus membrane, producing ulceration upon the latter down into the bronchial tubes and capillaries of the lungs, painful coughing; must be careful not to breathe the gas; if necessity compels to stay in a place filled partly with chlorine must keep mouth and nose covered with a wet sponge. The gas is very heavy. It is 35.5 times heavier than hydrogen, 2.45 times heavier than air, and 2.216 times heavier than oxygen. One litre of the gas at  $0^{\circ}$  C. and 760 mm. mercury pressure weighs 3.178 grams. 1 c.c. equals 0.003178 grams, roundly 3 milligrams. The gas dissolves somewhat in water. The maximum solubility lies at  $9.5^{\circ}$  C., and corresponds to 2.75 times the volume of the water. Hence 10 c.c. of water will absorb at best 27.5 c.c. of the gas. But since 1 c.c. of gas weighs three milligrams, the 27.5 c.c. will weigh 82.5 milligrams, or 0.0825 grams, and hence in weight per cent. 0.825. It is well to remember that the most concentrated water solution of chlorine does not contain quite one per cent. of chlorine; at  $30^{\circ}$  C. only 1.75 volumes equals 0.525 weight per cent. At boiling heat the chlorine is *completely* driven from its solution.

Chlorine gas condenses into a mobile yellow liquid under a pressure of 8.5 atmospheres (127.5 pounds per square inch).

*Preparation of chlorine.* In the practical preparation of chlorine it is more advantageous to supply the oxygen in combined form, as a superoxyd, an oxyd which contains more oxygen than it can hold


firmly. Nature furnishes us with such in the so-called soft *manganese ore*  $\text{MnO}^2$ , the *pyrolusite* of mineralogists. The salt-gas acts upon this oxyd even at ordinary temperature; very energetically at about  $60^\circ \text{C.}$ ; thus



Only one-half of the chlorine contained in the salt-gas appears as *free chlorine*, the other half forming, with the metal manganese, a pale, rose-colored salt—*manganese chlorid*,  $\text{MnCl}^2$ .

Neither is it convenient nor economical to act with the salt-gas upon the superoxyd. A solution of the gas in water is much better adapted to this purpose, the best concentration is 20 per cent. How to make such a solution will be explained in the next paragraph below. The most suitable apparatus is the Koenig generator, represented in Fig. 37. *G* is the generating tube drawn out at lower end *O*, where a stout rubber tube *R* is wired onto it. The tube *R* has at its other end a bent glass spout *P*, which rests upon the edge of the waste vessel *W*. The upper end of *G* is somewhat restricted, and a glass stopper *S* is ground into the restriction. This stopper is fused to the funnel tube *F*, the latter carries a stop-cock and a receiving bowl for the salt-gas solution. *L* is the outlet for the chlorine. *J* is a wider glass tube, fused onto *G*, so as to form a complete jacket around the latter. The tubulature *E* serves to fill this jacket either in part or entirely

FIG. 37.



KOENIG'S GENERATOR.


with water whilst a glass tube *T* encircles the jacket obliquely, and is fused into the jacket at both ends. This small encircling tube is full of water, and, being heated by the small flame from a glass tube or a Bunsen burner, brings up a circulation and raises the temperature in *J* to any desired point. Upon the porcelain false bottom *D* rests the manganese superoxyd in small pieces, about pea-size. The column of  $\text{MnO}^2$  should not be over 3 inches high. The apparatus is held by the brackets *B*, *B*, *B*, against the wooden stand *Q*, the latter being supported by the bottom plate *N*. We start by filling water into *G* until the tube *R* is full and the water-level in the spout *P* is even with the level at the false bottom *D*. Then we heat the jacket to  $60^\circ \text{C}$ ., and thereupon drop the salt-gas solution at the rate of one drop a second. There will then be a steady current of chlorine gas issuing through, whilst an equally constant discharge of the by-product, *i. e.*,  $\text{MnCl}^2$  will discharge itself at *P* into the waste vessel *W*, provided the bowl of the funnel is kept replenished. If a stopping be desirable, simply turn the stop-cock in *F*. The gas will not be quite pure. It will be mixed with some liquid particles and also with some salt-gas. Hence we conduct it through a wash bottle containing some water, then through another such containing  $\text{H}^2\text{O}.\text{SO}^3$ ; in the latter the aqueous vapor will be removed, in the former the other admixtures. Of course, whenever *dry* gas is not required, the second wash bottle may be omitted. You will take notice that the rubber tube *R* forms



a trap against the escape of the gas downward, whilst the narrow funnel tube keeps the gas from escaping upwards. But it stands to reason that if a resistance be placed at *L* of greater weight than the column of liquid in either *F* or *R*, then the gas escapes through them, they forming the natural safety valve.

A generator may be rigged more simply and

FIG. 38.




cheaply. *A*, Fig. 38, is a small flask, *t* is a glass tube, *F* a funnel connected with *t* by a short rubber tube and clamp *C*, *t'* tube for escaping gas; *r* a  $\frac{1}{8}$ " rubber tube, *W* a beaker glass in which stands a large test-tube fitted with rubber stopper *S'*, and partly filled either with water or with  $\text{H}^2\text{O}.\text{SO}^3$  serving as wash bottle. The manganese superoxyd is placed in *A* at *M*, *F* is filled with salt-

gas solution. The apparatus is ready for use. In a measure, however, as  $\text{MnCl}^2$  accumulates in *A*, the evolution of chlorine becomes slower, a steady stream cannot be maintained by it, except for a very short time. I worked for twenty-five years with such an apparatus, until I made the more perfect one described above.


*Chemical properties of chlorine.* Chlorine is a most powerful agent at ordinary temperature, when oxygen is almost inert. It attacks *all* the metals, and in presence of water dissolves most of them: Gold, platinum, tin, copper, iron, zinc, but not so lead, silver, mercury.

*Upon oxyds, it acts thus:*



Chlorine acts upon all coloring matters taken from plants, such as litmus, indigo, the red cabbage, and many others, in such a way that the color vanishes, *bleaches*.

Upon the hydroxyds of potassium and calcium, chlorine acts as follows:




The milk of lime becomes dissolved to a clear liquid. If the dry slaked lime—flour of lime—be exposed to chlorine the same action takes place, but the resulting product is a slightly pasty solid, and is called *bleaching lime*; is soluble in water, and is used in large quantities by the bleachers and dyers

of yarn and cloth. The compound  $\text{CaCl}^2$  does not bleach, only the compound  $\text{CaO.Cl}^2\text{O}$ , and in this only the oxyd  $\text{Cl}^2\text{O}$  is the bleaching factor.

Upon potassium hydrate the action is parallel, thus:



You pass the chlorine gas into the solution so long as it is freely absorbed. The result is a *bleaching* solution. But if we boil this solution for some time, it begins to throw out scaly crystals, white or colorless. Let these crystals be separated from the liquid, then dried, and then heated in a closed glass tube, when they will be seen to melt easily with strong evolution of gas. The gas proves to be *pure oxygen*. (Explode it with 2 volumes of hydrogen.) When no more gas is given out the residue contains only potassium and chlorine, is KCl. The crystals themselves are produced from the boiling solution by the following reaction:



and when the crystals are decomposed by heat:



The crystals represent a compound of the metallic oxyd  $\text{K}^x\text{O}$  with the non-metallic *superoxyd*  $\text{Cl}^2\text{O}^5$ . The latter is unstable, overloaded, hence heat breaks it up easily, and as we have seen above that chlorine drives oxygen from the oxyds, the final result must be  $\text{KCl} + 6$  oxygen. The crystals shall be known as *potassium chlorate*. It is a very valuable compound

to us, because we can obtain by it at any time quantities of the purest oxygen.

*Manufacturing process for potassium chlorate.* The potassium hydrate is relatively costly, the calcium hydroxyd very cheap, KCl is also cheap (a natural mineral *sylvite*). On passing chlorine gas into water in which have been slaked 3 equivalents of calcium oxyd (burnt lime) at boiling heat until the liquid has become clear, we have a solution of calcium chlorate (easily soluble). We bring into it one equivalent of KCl and potassium chlorate falls out in crystals (because it is *not* readily soluble in water).

*Composition of salt gas.* We found the gas composed of hydrogen and chlorine. Now we have to establish their ratio in the compound. If we pass one volume of salt gas over heated zinc repeatedly, in the apparatus Fig. 36, we find that the volume of the gas becomes reduced to one-half. Chlorine unites with zinc, becomes solid so to speak, the remainder is pure hydrogen. Hence we deduce : Salt gas is composed of equal volumes of hydrogen and chlorine expressed by the symbol HCl. If we fill into a cylinder one volume of hydrogen and one volume of chlorine and let the mixture stand in the diffused daylight for some days, the volume does not change, but the two gases shall have become united to HCl. We can prove this by introducing a few c.c. of water ; the gas is all absorbed by the water ; chlorine would have only been slightly absorbed, hydrogen not at all. Should we expose the mixture

of  $H + Cl$  to the *direct* sunlight, the union would follow at once with explosive energy. It is important to note that the volume of the compound is equal to the sum of the volumes of the components; neither contraction nor expansion taking place. *This is a law for all unions of gaseous bodies in equal volumes.* The name for the compound  $HCl$  shall be *hydrogen chlorid*, and all combinations of metals with chlorine shall be named *chlorids*, as we name the oxygen compounds *oxyds*. Some chemists speak and write *chlorides*, *oxides*; it is quite immaterial which you use, but choosing one you should stick to it; the shorter sound would seem to be preferable.

*Properties of hydrogen chlorid*: A colorless gas at ordinary temperature, powerfully pungent odor, exciting the mucous membrane. Near the freezing-point of water at  $+4.4^{\circ} C.$  the gas becomes a liquid under a pressure of 30.67 atmospheres or 460 lbs. per square inch. At a temperature of  $-73.3^{\circ} C.$  only a pressure of 27 pounds is needed. Liquid  $HCl$  is mobile and colorless, heavier than water. No practical use has been found for it. The specific gravity of the hydrogen chlorid gas is 1.255 (air = 1); 1 c.c. of it weighs 0.00163 gram, just about one-half that of chlorine. By weight the gas contains 97.26 of chlorine, 2.74 of hydrogen. 1 volume of water can absorb 500 volumes of  $HCl$  gas at the freezing-point; at  $20^{\circ} C.$  (common temperature) water absorbs 440 volumes of the gas. The absorption of the gas produces heat. This would lead us to think that there must be a chemical affinity

between  $\text{HCl}$  and  $\text{H}^2\text{O}$ ; that there must be *hydrates*. In fact it is quite probable that two such exist. For, if a concentrated solution of  $\text{HCl}$  in water be heated (a thermometer registering the temperature), it will give out for some time only moist  $\text{HCl}$  gas. The temperature having risen to  $100^\circ \text{C}$ ., water passes over with  $\text{HCl}$  and a very concentrated solution condenses having specific gravity 1.19. As the temperature rises the distillate becomes more watery until the temperature reaches  $111^\circ \text{C}$ ., at which it remains constant whilst a solution distills over possessing specific gravity 1.104. This solution contains 21 per cent.  $\text{HCl}$  and 79 per cent. of water, nearly the hydrate  $2\text{HCl} + 15\text{H}^2\text{O}$ . A second hydrate is  $\text{HCl} + 6\text{H}^2\text{O}$ .

*Hydrochloric acid—Muriatic acid*— $\text{HCl} + \text{water}$ . These names are given to the water solution of  $\text{HCl}$ . We speak of highly concentrated acid, concentrated acid, dilute acid, very dilute acid.

	Sp. gr.	Per cent. $\text{HCl}$
Highest concentrated,	1.200	40.77
Highly concentrated,	1.1802	36.29
Strong acid,	1.151	30.58
Medium,	1.072	14.68
Dilute,	1.042	8.56
Very dilute,	1.006	1.12


Problem: Construct with these data a curve whose ordinates shall be the percentage and the abscissae, the specific gravities.

*Preparation and manufacture.* Small quantities are made by distillation in glass flasks or glass re-



torts. On a commercial or manufacturing scale, cast-iron vessels are used, either cylinders, or else flat pans standing within a brick furnace. Cast iron is not attacked by concentrated sulfuric acid nor by  $\text{HCl}$ , but is energetically attacked by a solution of  $\text{HCl}$  in water. Such properties render the iron vessels fit. Concentrated sulfuric acid acts violently upon salt even at ordinary temperature; the mass

FIG. 39.



threatens to froth over. An addition of water gives relief. Our prescription is: For every ten grams of salt take 9.6 c.c. of concentrated  $\text{H}^2\text{O}.\text{SO}^3$  and 5.5 c.c. of water. Mix the two liquids in a beaker-glass, fill from it the basin *B* (Fig. 39) of the funnel; place the salt in the flask *F*; make connection with the Wulf bottle *W* by tube *t* which passes

under the level of the wash water. The tube  $t'$  also passes under the surface level and is open at the top;  $t'$  is the safety valve of the apparatus, because air will pass into it whenever a partial vacuum is brought about;  $t''$  leads into the absorption flask  $A$  which contains as much water by weight as the salt taken. The student takes 50 grams of salt and places 50 c.c. of water in the absorption bottle. The stoppers  $S, S, S$  must be of rubber. The flask  $F$  must hold 500 c.c. All the acid may be put in at once, and then heat applied gently at first, and regulated to keep up a steady evolution. When the gas stops, finally, the operation is terminated and you let air in through the funnel tube. In removing the flame, it will be seen that the liquid in  $F$  solidifies, by degrees, as it cools down. Remelt it and pour it out on a clean stone or iron surface. Let it be named "crude salt cake." It must, evidently, be the vitriol of the unknown metal  $M$ , and will be taken up presently.

*Action of hydrochloric acid.* The water solution of  $\text{HCl}$  is nearly as powerful an agent as the gas itself, and for most purposes can be employed in its stead. Because the metallic chlorids are more readily soluble in water than the vitriols, excepting the chlorids of silver, lead, and mercury; therefore, we shall use it in preference to the sulfuric acid, whenever we desire to dissolve bodies which are not soluble in water. As, for instance, we wish to generate lime-gas. The calcium vitriol is very insoluble, and we have heretofore used acetic acid for the decomposi-

tion. Instead we shall use hydrochloric acid hereafter; the calcium chlorid is much more soluble than the vitriol, in fact, it will dissolve by merely allowing it to stand uncovered in ordinary air which is always more or less moist. Hence, we can employ, to advantage, this chlorid in place of concentrated sulfuric acid for the drying of gases. Being a *porous solid*, the calcium chlorid offers more surface to the gas than an equal volume of *liquid* sulfuric acid.

The chemical actions may be represented thus: If R stands for a metal, whose chlorid is soluble in water:

$R + \text{water} + \text{HCl solution} = \text{RCl solution} + \text{H};$   
and on oxyds

$\text{RO} + \text{water} + 2\text{HCl solution} = \text{RCl}^2 \text{ solution} + \text{H}^2\text{O};$

if, however, silver vitriol solution be brought together with hydrochloric acid, a white, curdy precipitate falls out at once, *AgCl*,

$\text{Ag}^2\text{O}.\text{SO}^3 + \text{water} + 2\text{HCl solution} = 2\text{AgCl} + \text{H}^2\text{O}.\text{SO}^3 \text{ solution}$

because the silver chlorid is insoluble in water. Thus we can prove the presence or absence of silver in any unknown solution by adding to it a drop of hydrochloric acid; if a curdy cloudiness follows, then silver is present.

Pure hydrochloric acid is quite colorless. The yellow color of the commercial muriatic acid is owing to some *iron chlorid* coming from the iron ves-

sels and iron oxyd in the crude salt. The strength of commercial acid is mostly indicated in degrees on the *Beaumé hydrometer*, instead of by the specific gravity. Such an instrument (Fig. 40) is correct if it sinks in pure water at  $17^{\circ}$  C. to the zero mark. Zero is therefore equal to specific gravity 1.000. In a 15 per cent. solution of common salt (salt = 15, water = 85) the spindle must sink to the mark 15; the degrees being equal divisions. Very concentrated

FIG. 40.





FIG. 41.



hydrochloric acid is said to be  $22^{\circ}$  Bé. (specific gravity 1.176). There is, of course, a second spindle for liquids lighter than water. In this instrument (Fig. 41) the zero point is near the bulb, the weight in the bulb, being so chosen that the spindle sinks in a solution of 10 per cent. salt (salt 1, water 9) to the zero point, whilst in pure water it sinks to

division 10; the same unit (found by dividing the space between 0 and 10 into 10 equal parts) is then drawn on the scale upward to the limit of the spindle. Kerosene, gasoline and other coal-oil products are gauged in the United States by degrees Bé. Alcohol is gauged by the hydrometer of *Tralles*. It is not sufficient to say that a liquid stands at so many degrees Bé., it must be said whether reference is had to a liquid lighter or heavier than water, which can be done by the signs +, —, or by letters: l. w; h. w. Thus the light gasoline for gas-making is  $-87^{\circ}$  Bé., and the concentrated sulfuric acid is  $+66^{\circ}$  Bé. Hydrochloric acid can be shipped only in glass vessels; the sulfuric acid can be transported in iron tanks. The glass vessels—large, spherical, holding 5 to 10 gallons, and being packed with straw into wooden boxes—are known as *carboys*. For manufacture on a large scale get information in Lunge's *Manufacture of Soda Ash* (recent) or in *Ure's Dictionary* (old).

*Crude salt cake, discovery of the metal sodium.*

Let some of the crude cake, obtained by action of  $\text{H}^2\text{SO}^4$  on salt, be heated in a porcelain crucible over an open flame. Dense white fumes are soon seen to arise. The fumes are now diagnosed by us as probably being sulfur trioxyd, oil of vitriol. As the vapors escape, the melting-point of the cake rises. Finally, when no further fumes come off, the stuff appears dry at a red heat. At yellow heat it melts again, and remains so without yielding any more fumes or gas. Reason for the fumes: The

crude salt cake is  $M^sO.SO^3.H^2O.SO^3$ , a so-called acid vitriol, quite constant at low heat, but breaking up at high heat into  $M^sO.SO^3 + H^2O + SO^3$ . The final remnant is  $M^sO.SO^3$  = salt cake—*refined*. We had added in the recipe given just twice as much sulfuric acid as is needed, and did this knowingly, for otherwise we could not have completely decomposed the salt in a glass flask ; the stuff would not have liquefied ; the flask probably broken. Salt cake dissolves in cold water easily ; this property distinguishes it from the potassium vitriol and the calcium vitriol. In order to isolate the metal let us make use of our experience with charcoal—carbon. We mix the powdered salt cake with powdered charcoal, cover the crucible, and expose it in a charcoal or gasoline furnace to strong red heat. We may do it in a hard glass tube first. Result is a dark-colored mass ; metallic particles are not visible, and if we bring it together with water there is no evolution of gas, such as potassium produced. However, it dissolves in water, leaving charcoal powder, which we separate by filtration, and test, after thorough washing, by burning ; as it quite disappears (leaving only trace of ash), we are correct in calling it charcoal. The filtered liquid has a brown or red-brown color ; it gives alkaline reaction to litmus ; has a strong taste. With hydrochloric acid, gives off gas smelling of rotten eggs, and if this gas be passed through a glass tube heated to redness a sublimate of sulfur forms. If bright copper or silver be brought together with the liquid,




the metal turns black, and if these strips of blackened metal be heated in an open tube we get the smell of  $\text{SO}^2$ . If boiled with finely-divided copper oxyd, the liquid becomes decolorized and does not any more blacken the metallic copper, whilst it gives still a strong alkaline reaction. From all these observed facts we draw the following deductions:

1. The action of charcoal must have been, symbolically, thus




the vitriol was converted into a *sulfid*.

2. The action of metallic copper and metallic silver was probably



3. The action of copper oxyd was probably



The hydroxyd of the unknown metal is easily soluble in water, and being strongly basic—alkaline—it must be akin to potassium. We evaporate the liquid from (3) to a white solid, which afterwards fuses and then goes off in white fumes, just like potassium hydroxyd. We act upon this material with zinc, and by evolution of hydrogen prove it to be hydroxyd. Next we act upon it with the electric current; gas at positive pole; metallic globules and gas at negative pole; and by using mercury as the negative pole we obtain a solid amalgam, from

which the metal M may be separated by distillation in a current of hydrogen gas. If we act upon the hydroxyd with charcoal-loaded iron chips at a yellow heat, the metal will distil over, same as potassium, but the flame issuing from the retort burns with a *yellow color* (distinction from potassium, *purple flame*). In the neck of the retort we find a black mass which acts like that found in the distillation of potassium. This mass is mixed with the condensed metal.

*Properties of the metal sodium.* The name *sodium* is given to the metal by English, French, and American chemists. Germans and all others give it the name *natrium*, using the first two letters as the symbol, Na, which stands for the chemical unit of mass. English, Americans, French use the same symbol, hence sodium, Na. Natrium is derived from Greek *nitron*; Egyptian and Hebrew *neter*, a name given by those people to the crusts forming around the desert lakelets, and which was found to have similar cleansing properties to potash. It is now known as soda ash. Whence the word soda comes is not known, nor what it means. Perhaps from the Latin *soldere*, English solder, since this material may have been confounded with borax, such mixtures being useful in joining metal pieces by heat. (Author's notion.)

The metal sodium is soft like wax at ordinary temperature, can be hammered at freezing-point, becomes liquid at about  $95^{\circ}$  C. (melting-point is higher than that of potassium  $65^{\circ}$  C.). In color

it is silver-white like potassium ; the fresh-cut surface becomes rapidly dull from oxydation. Sodium becomes vapor at red heat, and this vapor has a purplish color—potassium green. Specific gravity at  $10^{\circ}$  C. is 0.974 (water = 1). Coefficient of expansion 0.000073, larger than that of any other metal except potassium (0.000083). It conducts heat and electric waves well, about 37 (silver = 100). Its specific heat or heat capacity is 0.2934. Its heat of fusion is 0.73 Cal.

*Chemical properties of the metal sodium.* Thrown upon water it sets up an evolution of gas and melts into a globule which is covered by a gray film. The gas does not ignite, unless the water be heated to about  $60^{\circ}$  C., or unless the water be thickened with gum arabic or glycerine. We deduce from this action that sodium has not as much affinity or attractive tendency for oxygen, as potassium ; hence less heat is generated. If the globule be left on the water until the gas evolution stops, the globule flattens out suddenly and an explosion ensues. If the globule be taken from water when gas stops, it is found composed wholly of *oxyd*. (The explosion is explained by the sudden rise of steam when the oxyd becomes hydroxyd.)

If the metal be heated in oxygen gas, or in a mixture of air and oxygen, a yellow substance results which is the *superoxyd of sodium* ; on cooling it turns white. It dissolves in water without decomposition, and can be melted without decomposition. If heated with copper, lead, zinc or tin the sodium

superoxyd changes these metals into oxyds. With HCl it gives  $\text{NaCl} + \text{H}^2\text{O} + \text{O}$ . If this action be done in water solution, *oxygen does not escape*; the water then contains sodium chlorid, NaCl, and hydrogen superoxyd (also called peroxyd),  $\text{H}^2\text{O}^2$  or HO; we assume therefore that the sodium peroxyd must have a similar composition to the hydrogen peroxyd, namely,  $\text{Na}^2\text{O}^2$ , the reaction will be




This solution of hydrogen peroxyd is a powerful oxydizing agent and is much used both in laboratory work and on a large scale for manufacturing purposes.

If the superoxyd has the composition  $\text{Na}^2\text{O}^2$  then the *oxyd* must be  $\text{Na}^2\text{O}$ . We get this by fusing together the hydroxyd with the metal in proper proportion. It is a gray substance, and of no special application. It attracts moisture from the air and becomes hydroxyd.






## CHAPTER VII.

### THE STORY OF SODA ASH AND LEBLANC.

WHEN in the early years of the past century Napoleon closed all the harbors of Continental Europe to English and American ships, in order to destroy the commerce of England, as he could not destroy its navy, there arose in Continental Europe a scarcity of potash, since the supply had nearly all come from the American colonies. To ease the demand for this article, as well as to spite England, Napoleon offered a prize of 100,000 francs, 20,000 dollars, for the discovery of a substitute for potash. This prize was won and awarded to the French chemist Leblanc, who proposed a process by which common salt can be turned into a body which may be substituted for potash in most technical applications, a process which held its own for 80 years and is only now giving way slowly to better methods. (See Solvay process at end of chap. XI.) All the reactions were known to chemists at the time but one, the replacing of copper oxyd by a cheaper compound. Leblanc found that limestone, or oyster shells, or chalk could replace the copper oxyd, if the action takes place at red heat. To wit:



$\text{Na}^2\text{O}.\text{CO}^2$  is easily soluble in water;  $\text{CaO}.\text{CaS}$  (calcium oxysulfid) is insoluble in water, and  $\text{Na}^2\text{O}.\text{CO}^2$  is the practical substitute for  $\text{K}^2\text{O}.\text{CO}^2$  (potash). The entire run from salt to soda ash is represented in the following symbolic scheme:




The soda ash produced in this way is not pure: The carbonate predominates, but mixed with it we find  $\text{Na}^2\text{O}.\text{H}^2\text{O}$ ,  $\text{Na}^2\text{S}$  and other impurities. It can be purified; not necessary for most of the applications, such as soap-making. A factory in which soda ash is produced goes by the name of *alkali works*. It is usually divided into several departments, located in separate buildings. (a) Acid works where the sulfuric acid is made. (b) Salt-cake works. (c) Black ash smelter and extractor. (d) White ash works. (e) Bleaching-lime works, where the hydrochloric acid is converted into chlorine and the latter is absorbed by the dry slaked lime.



SODA ASH—SODIUM CARBONATE, SAL SODA,  
BAKING SODA.

When the liquid resulting from action (4) (see above) is evaporated to dryness and then fired to red heat, the product bears the name *crude white soda ash*. If the same liquid is however only evaporated or boiled down to a certain point and is then allowed to cool slowly, large crystals will form. These are  $\text{Na}^2\text{O}.\text{CO}^2 + 10\text{H}^2\text{O}$  and go by the name *sal soda* (salt of soda). The impurities remain in the mother liquor. If they be exposed on wicker hurdles in an exposed space into which lime gas is conducted from the top of a lime kiln, then they will pass into a fine granular white sandy material whilst water runs away from them. The white powder goes under name of *baking soda, bicarbonate of soda*. The action is thus :



The bicarbonate is slightly soluble in cold water, and at the boiling temperature it breaks up into  $\text{Na}^2\text{O}.\text{CO}^2 + \text{CO}^2 + \text{H}^2\text{O}$ . If therefore this material be mixed with flour and the resulting dough is put into an oven we will just get that same decomposition, the escaping lime gas causing the raising of the dough. But the bread must taste bitter from the sodium carbonate which remains. In the so-called *baking powders*, the baking soda forms only one ingredient, the other being an acid salt which not only decomposes the carbonate but also removes the bitter taste.


*Caustic soda or concentrated lye*, of the manufacturers is made by decomposing a 10 per cent. solution of soda ash with one equivalent of slaked lime, the same process we considered under caustic potash. The resulting solution of the sodium hydrate is evaporated, fused and cast into sheet-iron drums, or cans, for smaller quantities, or into one-pound balls, which latter are then dipped into molten rosin. The film of rosin keeps the material from attracting moisture from the air and thus liquefying. The farmers' wives now buy these *lye balls* for their soap-making instead of bothering with the wood ashes. The man who had the idea of the *rosin film* made a fortune from the patent rights.

Certain sea plants growing in the tide levels of France and England, called *kelp* in Wales, and *varec* in France leave an ash which is largely made up of sodium carbonate, the plant's energy transforming the salt into the soda.


## CHAPTER VIII.

### A CHAPTER ON THEORIES; ON COMBINING WEIGHTS, ATOMIC WEIGHTS AND VALENCES.


THE existence of such bodies as *chlorids*, especially sodium chlorid, containing no oxygen, and yet so much alike to oxygen salts, leads us to thinking. Let us compare hydrogen chlorid,  $\text{HCl}$  and sulfuric acid,  $\text{H}^2\text{O}.\text{SO}^3$ . As the symbols stand written, two do not seem at all comparable. But supposing we change the grouping of the elements in the sulfuric acid thus:



then at first glance we notice the resemblance,



and more so still if we place the  $\text{SO}^4$  into a bracket,



We have no knowledge of an oxyd,  $\text{SO}^4$ ; we only know  $\text{SO}^2$ ,  $\text{SO}^3$ ,  $\text{S}^2\text{O}^3$ ,  $\text{S}^2\text{O}^4$ ; yet without a great wrench we can imagine that the very moment when the two oxyds,  $\text{H}^2\text{O}$  and  $\text{SO}^3$ , are brought together, a rearrangement of the elementary particles comes about harmoniously, quite imperceptible to the eye. As soon as we disturb the harmony by trying to remove the hydrogen, then disarrangement, a break-up takes place. But if we accept the reality of

this unprovable state of things, then the schism in the fundamental chemical phenomena gives way to resolved uniformity. *The acids become then combinations of hydrogen with a non-metallic radical*; the radical can be either one non-metallic element or a group of such elements. In hydrochloric acid, chlorine is the radical, in sulfuric acid the group ( $\text{SO}^4$ ) is the radical. The word *radical* is the adjective of the Latin noun *radix* = root; from the radical, root, arises the stem—the sourness, the acidity. When a metal acts upon an acid, hydrogen escapes. According to the new light we circumscribe this by saying the metal takes the place of the hydrogen :



Why does it do so? Because potassium has a stronger affinity for the radical chlorine than hydrogen, and this stronger affinity is made perceptible by the greater quantity of heat which is set free by the union of the two. If Cal. = the unit of heat, then



$$n' > n.$$

Metals such as lead, copper, silver, gold do not act upon the dilute acids; they are not dissolved by them, because their heat of formation is very small and requires a heat-addition from the external conditions. Thus we can establish a series of the metals in which potassium will occupy the one flank and

gold the other: K., Na., Ca., Mn., Zn., Fe., Sn., Pb., Cu., H., Ag., Au.

If we designate K as +, then Na will be negative in regard to K, but positive in regard to Ca, and each metal in the same way positive towards its neighbor on the right, negative towards the one on the left. Equally if we place two pieces of sheet metal upon one another with a moist piece of felt between, for example, zinc and copper, an electric tension will show itself between them. Galvani observed this 120 years ago, and his name is attached to this electric current even now—galvanic electricity as against frictional or static electricity. The series is therefore usually alluded to as the electro-chemical series of metals.

Of the non-metals, oxygen occupies the extreme flank, with chlorine next, as sodium stands alongside of potassium: O., Cl., Br., S., N., C.

*Valence.* Glancing at the symbols  $H(Cl)$ ,  $H^2(SO^4)$  we cannot help but being struck by the fact that in one symbol there is but one hydrogen, whereas there are two of hydrogen in the other. We cannot remove one of these hydrogens without destroying the harmony, without breaking up the body completely. To give expression to this fact we are led to the term *valence*, a slight difference of meaning from *equivalence*. We say the elementary group (Cl) is *monovalent*, because it finds its satisfaction with one volume-unit of hydrogen, *monos* = once; the compound group ( $SO^4$ ) is *divalent*, *dis* = twice, because it must have two hydrogens to exist. We

shall find later on elements as well as complex radicals whose valence are 3, 4, 5, 6 designated respectively as *tri*-, *tetra*-, *penta*-, *hexavalent*. I warn you not to be dazzled by these full, sonorous words; they are but a short cut of speech. You can prove nothing by means of valence, because the expression valence only states a fact, yet is convenient as an expression.

*The atomic weights of elements, molecular weights of compounds, volume weights.* We find that one volume of chlorine unites with one volume of hydrogen; in symbols  $\text{HCl}$ . Chlorine gas is 35.5 times heavier than hydrogen gas hence the symbol stands


$$\text{HCl} = 1 + 35.5 = 36.5$$

Also we find that 35.5 grams of chlorine gas combine with 23 grams of sodium and thus produce  $35.5 + 23 = 58.5$  grams of  $\text{NaCl}$  (common salt). Only one compound between  $\text{Cl}$  and  $\text{Na}$  has been observed with certainty and this is a very *stable* one. We *assume* that the vapor of sodium, if it could be produced and weighed, would be 23 times heavier than hydrogen, under the same conditions of temperature and pressure. Though the metal volatilizes at red heat yet have all experimental attempts thus far failed, because the sodium vapor attacks all the vessels which are available: platinum, gold, silver, nickel, porcelain; hence we do know only by *inference* that the volume of the mass unit weighs 23.


Both by composition and decomposition (synthesis and analysis) we know that 35.5 grams of chlorine gas combine with 39 grams of potassium to form





35.5 + 39 grams of KCl. That 39 is the weight of one volume of potassium vapor we do not know any better than in the case of sodium, for the same difficulties. The symbol KCl or NaCl is not a certainty in other words. Its strong probability follows from the following consideration. By acting upon sulphuric acid  $H^2(SO^4)$  with either of the two metals we can produce well crystallized salts or vitriols to wit :



either of which possesses strong acid reaction, and



salts which are neutral towards litmus paper. The former salts convert into the latter thus :




Transposed into numbers this means that we can combine with one S (32 parts) either 23 or 46 of sodium, either 39 or 78 of potassium ; but with 23 and 39 only when in each instance there is also one hydrogen present. Hence it follows that 23 of sodium, 39 of potassium, can take the place of—are equivalent to—one hydrogen in the two acids



Potassium, sodium, hydrogen, on these given terms are monovalent metals, or *monads* (a still shorter expression), It is self-evident then that the oxyd of these metals must be  $\text{Na}^2\text{O}$ ,  $\text{K}^2\text{O}$  if the oxyd of hydrogen is  $\text{H}^2\text{O}$ . The hydroxyds of the metals sodium and potassium become  $\text{Na}^2\text{O}.\text{H}^2\text{O}$ ;  $\text{K}^2\text{O}.\text{H}^2\text{O}$ . But since we saw in the vitriols one hydrogen being replaced by one Na, or one K, the same must be possible in water, to wit :



Hence it follows that the symbol  $\text{Na}^2\text{O}.\text{H}^2\text{O} = 2 (\text{NaHO})$  expresses *two* units or molecules of sodium hydroxyd, that *NaHO* is the true representation of sodium hydroxyd. In the metallic hydroxyd we have, therefore, a combination in which the metal is united to the group (HO). A group which acts as a non-metallic radical of the value (valence) one. This group contains the metal H and the non-metal O, whilst the negative or non-metallic group ( $\text{SO}^4$ ) contains the two non-metals. Perhaps in this hybrid nature of the group (HO) lies part of the reason for the action of the hydroxyd towards litmus and other actions totally opposed to the acids. The relative action, then, of the hydroxyds and acids is this :



Two attractions exist to account for the powerful action : First the greater attraction of the metal to the non-metallic radical and the tendency of (HO)

to become  $\text{H}^2\text{O}$  by taking another H. Owing to the important role of the group (HO) the name *hydroxyl* (*ulé* = matter) is given to it. Hydrogen = the generator of water, hydroxyl = the matter from which water is made. \* In the light of all these considerations and speculative deductions, the previous definitions of base, acid, salt shall be changed to read as follows:

*Base* = Combination of metal with one or more hydroxyl groups.

*Acid* = Combination of a non-metallic group or radical with one or more hydrogens.

*Salt* = Combination of a metal with a non-metallic radical.

The definitions of acid and salt are identical. The two things are of *one kind*. *Sulfuric acid is hydrogen vitriol*.

The term vitriol has been abandoned for the sake of greater uniformity of chemical expressions. Its place is taken by the word sulfate, hence the following:

$\text{H}^2(\text{SO}^4) = \text{Hydrogen sulfate} = \text{sulfuric acid.}$

$\text{Na}^2(\text{SO}^4) = \text{Sodium sulfate.}$

$\text{NaH}(\text{SO}^4) = \text{Sodium, hydrogen sulfate.}$

$\text{K}^2(\text{SO}^4) = \text{Potassium sulfate.}$

$\text{KH}(\text{SO}^4) = \text{Potassium, hydrogen sulfate.}$

$\text{Ca}(\text{SO}^4) = \text{Calcium sulfate.}$

$\text{Fe}(\text{SO}^4) = \text{Ferro sulfate (iron sulfate).}$


$\text{Cu}(\text{SO}^4) = \text{Copper sulfate.}$

$\text{Pb}(\text{SO}^4) = \text{Lead sulfate.}$

$\text{Zn}(\text{SO}^4) = \text{Zinc sulfate.}$


You should accustom yourself to use the expression: hydrogen sulfate in place of sulfuric acid, though no harm is done by the latter. Logical consequential speech leads to logical thought and work.

*Atomic weight of calcium, copper, lead, zinc.* By analysis we find that 35.5 grams of chlorine unite with 20 grams of calcium to a stable chlorid. If 20 were the representative of one volume of calcium vapor then the symbol of the chlorid would be  $\text{CaCl}$ , as  $\text{NaCl}$ . But we find that we can combine the calcium with the hydrogen sulfate,  $\text{H}^2(\text{SO}^4)$ , only in *one way*, not in two ways as with potassium and sodium, namely, so that 40 of calcium correspond to one  $(\text{SO}^4)$  or one S, that therefore the number 40 must stand for the atomic weight of calcium, that one calcium is equivalent to two hydrogens and therefore the symbol of the chlorid must be written  $\text{CaCl}^2$ .




Copper, zinc, lead, also form only one kind of sulfate,  $\text{Cu}(\text{SO}^4)$ ,  $\text{Zn}(\text{SO}^4)$ ,  $\text{Pb}(\text{SO}^4)$ , therefore, we say these metals are *divalent* like calcium, their unit weight stands for *two* hydrogens, and in each case 35.5 chlorine combine exactly with one-half as much metal as 32 sulfur, hence their chlorids are  $\text{CuCl}^2$ ,  $\text{ZnCl}^2$ ,  $\text{PbCl}^2$ . Copper makes an exception in so far as it can unite with chlorine in two ways, to wit:  $35.5\text{Cl} + 31.5\text{Cu}$  and  $35.5\text{Cl} + 63\text{Cu}$ . The compound whose ratio of Cl : Cu is 35.5 : 31.5 is the stable compound, permanent at ordinary heat and even

up to red heat. But since in the sulfate 32S correspond to 63Cu, therefore we take the number 63 as the representation of one divalent volume of copper and write



In the first, the stable compound, *cupric* chlorid,  $\text{CuCl}^2$ , the metal is normal, divalent. In the second, the unstable compound, *cuprous* chlorid,  $\text{Cu}^2\text{Cl}^2$ , the metal is abnormal, monovalent. Silver acts like potassium. 107.6Ag combine with one chlorine; but  $2 \times 107.6$  combine with one sulfur, 32. Hence *silver is monovalent*,  $107.6\text{Ag} = \text{one H}$ . The chlorid is AgCl, the sulfate is  $\text{Ag}^2(\text{SO}^4)$ .


Gold unites with chlorine in two ways. In one compound, which is a white, powdery substance, we find 196Au with 35.5Cl, in the other 65.33Au with 35.5Cl. The first compound is so unstable that it falls to pieces upon the addition of water. In the other compound the chlorine acts upon certain substances as free chlorine. Besides, the specific weight of gold—19.5—is so high that necessarily the weight of its vapor must be very high (though we cannot make this vapor). We take, therefore, 196 to represent the weight of one volume, and write the symbols of the two chlorids:



Auric chlorid is a deep yellow substance, easily soluble in water, and fairly stable. In it gold =

Au has the valence 3. In aurous chlorid Au has the valence 1.

*Molecular weight.* Two or more simple bodies united into a chemical union form a *molecule*. By some it is contended that in the free state even the simplest bodies—the elements—form molecules, that in the free state hydrogen is  $(\text{H.H}) = 2$ , chlorine  $(\text{Cl.Cl}) = 71$ ; oxygen  $(\text{O.O}) = 32$ , and  $\text{H}^2(\text{SO}^4) = 98$  of course. When chlorine acts upon hydrogen the action must, according to this view, be represented by :



Nothing is changed, in reality, by adopting this view, or by rejecting it. If we speak of molecular weight we shall invariably mean that the molecule is composed of several elements.

*Relation between atomic weights and specific heat of the elements.* By *heat capacity* the physicists understand the quantity of heat energy expressed in *calories* (heat units), which is necessary to raise the temperature of a mass equal to one gram of a substance by one degree of the centigrade thermometer. The *specific heat* of a body (solid or liquid) means its heat capacity referred to that of water as the unit. The specific heat of gases is referred to that of air or also to that of water as units. The variation of values thus obtained is highly astonishing. In general, the heat capacity of metals is very low, that is, a metal shows the heat very quickly, water very slowly. The numbers representing the specific



heats are therefore always true decimal fractions. *When the atomic weights of the metals are multiplied by their specific heats the product is a constant.* Reason therefore demands that whenever the product is not equal to the constant, there must be something wrong, that we stand before a riddle. The following numbers show this relation, which is also known as the "Law of Dulong and Petit:"

## A. W. Spec. Heat. Product.

Silver	Ag	$108 \times 0.0570 = 6.156$
Iron	Fe	$56 \times 0.1138 = 6.375$
Copper	Cu	$63 \times 0.0952 = 5.99$
Zinc	Zn	$65 \times 0.0955 = 6.207$
Calcium	Ca	$40 \times 0.167 = 6.68$
Sodium	Na	$23 \times 0.2930 = 6.73$
Potassium	K	$39 \times 0.1655 = 6.45$
Lead	Pb	$207 \times 0.0314 = 6.499$
Tin	Sn	$118 \times 0.0562 = 6.631$

These numbers do not show exactly the same product for all the metals, but the constant appears to be about 6.5.

Inversely it would follow that the *specific heat of the atomic volume is the same for all the metals*; the specific heat is an inverse function of the mass; the greater the specific gravity, the smaller the specific heat.

## REVIEW OF THE ACTION OF CHLORINE ON THE ALKALINE HYDROXYDS.

The action of chlorine upon the alkaline hydroxyds is so important, theoretically and practi-

cally, that we must now transcribe the symbols for those reactions according to the notion of valence :




1.  $2K(HO) + 2Cl = K(ClO) + H_2O$ . The radical  $(ClO)$  should be called *chloryl* like hydroxyl, but this name is rarely met with. It cannot be isolated, it has no real existence, it is unbalanced ;  $Cl^2O$  is the balanced or saturated molecule.

$Cl^2O$ —*Dichloroxyd* is at ordinary temperature a reddish-yellow gas of penetrating odor. Specific gravity = 2.977. 1 cubic centimeter weighs 0.0039 grams, nearly 4 mgs.; 1 c.c. of chlorine weighs 0.00317; 1 c.c. of oxygen weighs 0.00143. 2 volumes  $Cl + 1$  vol.  $O = 2 \times 0.00317 + 0.00143 = 0.00777$ . If the latter sum be divided by two we get 0.00388 which is equivalent to the experimental weight 0.0039. Hence it follows that  $2Cl + 1O = 3$  vols., in combining to  $Cl^2O$  contract one-third. We found this to be so for  $SO^2$  and for  $H^2O$ . We may deduce the general law that *two volumes of one element combining with one volume of another element always produce two volumes of the combination*. Thus is explained why the unit weight of a compound can be greater than the sum of the unit weight of its composing elements.

$Cl^2O$  becomes a blood-red liquid, when the gas is conducted into a tube which stands in a freezing mixture. The liquid is terribly explosive. A scratch with the file on the glass tube may cause an explosion; the  $Cl^2O$  just breaking up into  $Cl^2 + O$ . This is an interesting fact. For in spite of  $Cl^2O$  being a compound like water  $H^2O$ , yet whilst the


latter is strongly cohering, the former has little coherence, because in  $\text{Cl}^2\text{O}$  we have *two non-metals*, whilst in  $\text{H}^2\text{O}$  we have *metal and non-metal*. The compound  $\text{Cl}^2\text{O}$  is made by acting with chlorine upon the oxyd of mercury, thus:  $\text{HgO} + 4\text{Cl} = \text{HgCl}^2 + \text{Cl}^2\text{O}$  (reddish-yellow gas). In its actions upon metals this body is more energetic even than chlorine itself.

When chlorine acts upon  $\text{K}(\text{HO})$  or  $\text{Na}(\text{HO})$  or  $\text{Ca}(\text{HO})^2$  at *boiling heat* the following actions occur :




The important products are  $\text{K}(\text{ClO}^3)$ ,  $\text{Na}(\text{ClO}^3)$ ,  $\text{Ca}(\text{ClO}^3)^2$ . These bodies we will designate *chlorates*. The group  $(\text{ClO}^3)$  is a monovalent radical but has no real existence; neither do we know the corresponding chloroxyd  $\text{Cl}^2\text{O}^5$ . But we can prepare the *hydrogen chlorate*,  $\text{H}(\text{ClO}^3)$ . It forms a thick, syrupy liquid at ordinary temperature, has strong acid taste, no odor. Above  $40^\circ \text{C}$ . it begins to give out chlorine and oxygen.

*Potassium chlorate*, is as above stated, the most important of the chlorates, because with it we can generate chlorine gas easily in immediate contact with the bodies to be acted upon :



*Generation of pure oxygen gas by means of potassium chlorate.*



Converted into figures this means: Molecular weight of  $\text{KClO}_3 = 39 + 35.5 + 3 \times 16 = 122.5$  grams, give 48 grams oxygen gas. One cubic centimeter of oxygen weighs 0.00143 grams, hence 48 grams =  $\frac{48}{0.00143}$  c.c. = 33636 c.c. = 33.636 litres = 0.033636 cubic meter.

*Problem.* Let a gas holder be a sheet-iron cylinder with the dimensions: Diameter = 13.5 inches, height = 27 inches. How many grams of potassium chlorate will be required to fill this holder with oxygen? 1 inch equals 2.5 centimeters = 0.025 m.

## CHAPTER IX.

### BROMINE, IODINE, FLUORINE.



#### BROMINE.

IN the process of salt-making from natural and artificial salt wells, the brine (salt solution) is evaporated. At a certain concentration of the boiling liquid, salt crystals fall out and keep on precipitating up to a given point. The crystals are steadily removed by means of a sieve-ladle. Finally a heavy solution remains from which no crystals of salt fall. This solution is the *mother liquor*. It contains the chlorids of calcium and magnesium,  $\text{CaCl}^2 + \text{MgCl}^2 + x$ ,  $x$  being the combination of the new element *bromine*, from Greek *bromos* = stench, with *magnesium*. If the mother liquor be heated with  $\text{H}^2\text{SO}^4$  and  $\text{MnO}^2$  the liquid becomes dark red-brown and heavy red-brown vapors appear above it. The vapors condense in a water-cooled receiver to a deep red-brown liquid, almost black, and emit a strong, irritating, suffocating odor (indicated in name). Specific gravity = 3.187 at  $0^\circ \text{C}$ ., 2.97 at  $15^\circ \text{C}$ . It boils at  $63^\circ \text{C}$ . and 760 mm. It becomes a brown-red, crystalline solid at  $-24^\circ \text{C}$ . The symbol for bromine is Br. Bromine gas is 80 times heavier than hydrogen. In most of



its chemical actions it is similar to chlorine. It is soluble in water; 1 part of bromine dissolves in 33.3 parts of water at  $15^{\circ}\text{C}$ . The solution is blood-red in color, and gives off bromine vapors. It is soluble in ether, alcohol, chloroform and carbon disulphid. If a water solution, containing little bromine, be shaken with carbon disulphid, the bromine will leave the water and pass into the carbon disulphid.

1 vol. Br. + 1 vol. H + heat = 2 vols. HBr.

Hence Br is monovalent like chlorine. The resulting HBr, *hydrogen bromid* is a colorless pungent gas like HCl. Liquefies at  $-73^{\circ}\text{C}$ . into a colorless liquid, which becomes solid when the liquid HBr is allowed to evaporate in the air. One c.c. weighs 0.003616 gr. HBr is eagerly absorbed by water, and is then named *hydrobromic acid*. The highest concentration is 82 per cent. HBr, corresponding to  $\text{HBr} + \text{H}^2\text{O}$ . HBr dissolves metals except Ag, Cu, Hg, Pb, because the resulting bromids  $\text{AgBr}$ ,  $\text{Cu}^2\text{Br}^2$ ,  $\text{Hg}^2\text{Br}^2$ ,  $\text{PbBr}^2$ , are not soluble. HBr combines with the hydroxyds, as HCl does thus



Bromids and oxybromids form when bromine acts upon the alkaline hydroxyds, thus:






*Potassium bromate*,  $KBrO^3$ , is even more insoluble than  $KClO^3$ . If therefore bromine be added to concentrated solution  $KOH$  (1:3) until the liquid retains a permanent yellow color,  $KBrO^4$  will fall out as a crystalline, colorless powder;  $KBr$  remains in solution.  $KBrO^3$  can be made pure by dissolving the powder in boiling water, when the salt will crystallize on cooling.

The most important salt is *sodium bromid*,  $NaBr$ . It forms a part of *bromo seltzer*; is much prescribed by physicians against headache. Tons of it are consumed annually.

Bromine itself and bromine water are valuable oxidizing agents in the hands of the chemist. Thus,



Upon heating the solution,  $HBr$  escapes with aqueous vapor and leaves hydrogen sulphate. There are many similar actions, with which we shall meet hereafter.

#### IODINE.

Before the discovery of bromine a French chemist, Courtois, had found a strange action in the mother liquor of the varec. By this name the peasants of the Channel Coast in northern France designate the extract from the ashes of the sea weeds which are thrown ashore by the storm. These ashes show alkaline reaction like the ashes of land plants, but it was recognized that the *sodium carbonate* is the principal component, not potassium carbonate, as in wood ashes. When the strained lye

from the water extraction is boiled down, potassium sulfate falls out first (being least soluble), then falls  $\text{Na}^2\text{SO}^4 + 2\text{H}^2\text{O}$ , then  $\text{NaCl}$ , then  $\text{Na}^2\text{CO}^3 + 3\text{H}^2\text{O}$ , finally leaving a mother liquor quite strongly alkaline, but containing still  $\text{NaCl}$ , together with  $\text{Na}^2\text{CO}^3$  and small quantities of sulfur-compounds of sodium. To this mother liquor  $\text{H}^2\text{SO}^4 + \text{water}$  (1.7 sp. g.), *pan acid*, is gradually added until the solution is decidedly acid.  $\text{CO}^2$  escapes and hydrogen sulfid, while a scum forms, chiefly consisting of sulfur. After this scum has been dipped out and the solution has become quite clear, it is transferred into a retort (cast iron with a leaden alembic or cap), manganese dioxyd is added and heat applied. Iodine vapors are given off which become a black crystallized sublimate in the receiver, made of earthenware. It is, however, impure with salts and water. The water is allowed to drain off, the residue redistilled with quicklime, which absorbs the remaining moisture. The nitre works of Chili furnish a mother liquor from which large quantities of iodine are manufactured. So also from the mother liquor of the chemical works at Stassfurt, Germany. In fact iodine or rather an iodid (either  $\text{NaI}$  or  $\text{MgI}^2$ ) is contained in the sea water, thence it gets into the algae (sea weed); also into the salt beds of the earth; thence into all salt springs and wells. More in some places than in others. The sea weed contains up to 0.6 p. c. of iodine, but most of it gets lost in the drying and the burning. All iodine producers are in a big trust, maintaining high prices. Probably a million pounds are consumed annually.

*Properties.* At ordinary temperature a grey-black solid, *always crystals* or crystal fragments; metallic lustre; emits a strong, unpleasant odor. It is very soft. Sp. G. 4.958. Melts at  $107^{\circ}$  C.; boils at  $180^{\circ}$  C. The vapor of iodine is of a beautiful *violet* color. The name iodine from *iodos* — similar to violets, was given for this color, which is so very characteristic and even unique. Water does not dissolve iodine freely. One gram I dissolves at  $10-12^{\circ}$  C. in 5524 grams of water. This solution is known as iodine water. It *bleaches* the same as chlorine water. Much more soluble in alcohol. This solution is known as *tincture of iodine*, much used by physicians, to relieve swellings of the skin. Makes a dark-brown stain on the skin, on wool or silk. Iodine dissolves readily in a water solution of potassium iodid, KI, giving a yellow, brown or blood-red liquid. Soluble in ether, in carbon disulfid. If a trace of iodine be contained in much water, or salt solutions, a few drops of carbon disulfid, shaken with the water, will absorb all the iodine and assume a rose color or purple color. The vapor of iodine is 127 times as heavy as an equal volume of hydrogen at the same temperature. Hence 127 is the atomic weight; the symbol is I.

*Starch and iodine.* If starch be boiled to thin paste, and the paste filtered, making a clear solution, then this colorless solution will become intensely *blue* if a small quantity of iodine solution be added. Starch + iodine equals *blue body*. Thus we can recognize starch from other parts of a plant

or seed by means of iodine, and detect iodine in solution with other bodies.

*Chemical properties.* Iodine combines with the metals directly forming *iodids*.  $K + I = KI$  (with explosive energy),  $Na + I = NaI$ , the two elements melt together without explosive display.  $Hg + 2I + \text{heat} = HgI^2$  (producing light) a scarlet-red body.  $Hg + I = HgI$ , a green-yellow body. These iodids are decomposed by bromine ;




Then in its turn  $KBr + Cl$  give  $KCl + Br$ .

Chlorine, bromine, iodine form a series whose chemical affinity is *inversely* as their atomic weights which are 35.5, 80, 127. The greater the mass, the more sluggish the activity.

Iodine does not readily combine with hydrogen, as chlorine does. It requires a high temperature. (6030 calories.)


*Hydroiodic acid*,  $HI$  + water is best prepared by working along the equation



We keep the finely powdered iodine stirred up in water whilst the gas  $H^2S$  is passed into the water ; the color of the solution disappears.  $HI$  is then dissolved in the water, the sulfur is to be removed by filtration. The liquid is strongly acid, smells pungent like  $HCl$  and acts upon metals and hydroxyds in a general way the same as  $HCl$ . A solution of  $Ag^2SO^4 + HI + \text{water}$  gives a *yellow precipitate of  $AgI$* , while  $HCl$  produces a *white precipitate of  $AgCl$* .

One does not often have occasion to prepare and use HI.

*Oxyiodids, iodates, hydrogen iodate, HIO<sup>3</sup>.* By the action of iodine upon KHO we obtain KI and K(IO<sup>3</sup>).  $6\text{KHO} + 6\text{I} = 5\text{KI} + \text{K}(\text{IO}^3) + 3\text{H}^2\text{O}$ . Potassium iodate is slightly soluble in water. If K(ClO<sup>3</sup>) be dissolved in water, to the liquid finely powdered iodine added and the solution boiled, then the iodine will displace the chlorine.



From an iodid as KI—chlorine displaces iodine; but from a chlorate iodine displaces chlorine, we say: Iodine has a stronger affinity for oxygen than chlorine. We make use of this property in quantitative analysis.

*H(IO<sup>3</sup>), Hydrogen iodate, iodic acid* is formed by acting with chlorine gas upon water in which finely ground iodine is suspended—




If a dilute solution of NaIO<sup>3</sup> be heated and chlorine be passed through until no further precipitation of salt be noticed, then the precipitate is Na<sup>2</sup>(IO<sup>6</sup>)H<sup>2</sup> or Na<sup>2</sup>O.H<sup>2</sup>O(IO<sup>4</sup>) *sodium hydrogen periodate*, and from this can be made the silver salt Ag(IO<sup>4</sup>), *silver periodate*.

*General remarks.* The most important compound of iodine is KI, potassium iodid, which forms white or colorless cubic crystals, like KBr, KCl; the three salts are *isomorphous*, have equal form and can re-



place each other in any crystal. We can have a crystal, any particle of which contains K, Cl, Br, I, even the very smallest. This leads us to the conclusion that Cl or Br do not in reality stand for one smallest unit, but that they represent a vast number each of smallest units; roughly, the circle represents the *active unit*, but the dots mean smallest particles so that one active unit may contain any number of isomorphous particles as Cl, Br, I. The two spheres, Fig. 42, mean the molecule K(Cl, Br, I).

FIG. 42.



The sphere representing potassium contains also a multitude of smallest particles, which in their turn may be a mixture of any number of isomorphous metals, such as Na, Ag. That is, the smallest fragment of a microscopic cube might contain (K, Na, Co, Rb, Tl), (Cl, Br, I). Minute quantities of iodine are found in the blood of man and animals as well as in plants, more particularly in the bones of the animal. We find iodine in the mineral *iodyrite*,  $AgI$ , among silver ores in yellow hexagonal crystals.

#### FLUORINE.

The mineral *fluorite*, *fluorspar* from German *flussspath*, occurs widely as *gangue* (German *gang* = vein)



or vein-matter with silver-lead ores, sometimes filling considerable fissure veins all by itself. The German miners call all minerals which are transparent or translucent and possess strong cleavage, *spath*, thus calcite is *kalkspath*, iron carbonate is *eisenspath*, orthoclase is *feldspath*, and our present mineral was named flussspath because the smelters noticed that it melts not only by itself, but causes other gangue minerals to become fluid, in other words, to act as a *flux* in smelting. *Fluss* = flux, whilst *fluere* is Latin for *to flow*.

Fluorite is characterized by its isometric crystals, cubes, octahedrons, tetrahexahedrons, hexoctahedrons, and its strong cleavage parallel to the faces of the octohedron. It scratches calcite, is therefore harder. Mostly colored green, purple, pink, blue, black, yellow, yet these colors are accidental, do not belong to the substance of the mineral itself, which is colorless or white. Shows no taste, therefore insoluble in water.


Fluorite melts just at the temperature glass melts ; experiment to be made in a small platinum spoon or crucible ; no gas will be given out. If, however, we put a piece of the spar upon charcoal and direct a strong oxydyzing flame upon it, the spar will melt at first, but after some time will solidify. Why ? Because the conditions are different from those in the crucible. The flame itself is not a mere source of heat, but a mixture of gases at high temperature. Among the gases are aqueous vapor and oxygen (from the air). We know both to be powerful

agents when assisted by heat. Indeed, by bringing the nose near the charcoal, a pungent odor becomes noticeable ; the spar decomposes under the influence of aqueous vapor and heat and oxygen. The residue assumes more and more the appearance of lime, glowing incandescence. By acting upon the finely-powdered spar with  $\text{H}^2\text{SO}^4$ , in a test-tube, there appears no action at ordinary temperature. Upon applying heat, gas bubbles arise, slowly at first, finally in large number, so that the liquid froths. A gas evidently escapes. To the nose the gas is pungent, acrid, recalling  $\text{HCl}$ , but more repelling. Litmus turns red at once in the gas.

*Investigation of the gas.* Hypothesis: Having an odor like  $\text{HCl}$ , we may infer that the gas is a compound  $\text{H}^n\text{X}^m$  or perhaps simply  $\text{HX}$  ; and proceeding a little further with the assumption that  $\text{X}$  is a new element, let  $\text{X}$  at once be represented by the letter  $\text{F}$  (first of fluor spar) ; hence the gas may be  $\text{H}^n\text{F}^m$  or perhaps  $\text{HF}$ . At the outset in the investigation, we find ourselves confronted with a serious difficulty. Namely we find the test-tube, which was used in the first experiment, strongly *corroded*, after it is washed out with water: The gas evidently decomposes glass. This unwelcome fact it is true puts a sure stamp of originality upon the new element, quite unlike any other thus far met with, but at the same time excludes the use of glass tubing and all other glassware. On porcelain it acts as on glass ; on iron and zinc it acts strongly, also on copper and silver. Not so on lead, gold and

platinum, nor on wax, paraffine or rubber. The latter materials do not stand heat. We are thrown then upon the three metals, of which lead is the cheapest. The gas acts somewhat upon the lead surface, but the product of the action being nearly insoluble, the metal answers well enough for all ordinary purposes. We construct a distilling apparatus of which Fig. 43 gives a sectional represen-

FIG. 43



tation. *C* is a cup of sheet lead with a ring-shaped groove *G* at the rim. The cup sits in an iron cup forming a sand bath, this resting on the tripod *T*. Into the groove *G* fits the alembic or helmet *A*, with the tube neck *N*. After filling the mixture of fluorite powder and  $\text{H}^2\text{SO}^4$  (so much acid that a thin mush forms) at *M* into the cup *C*, the alembic *A* is set into the groove *G*, and the latter filled with plaster of paris and water. The plaster *sets* and forms a gas-tight joint. The platinum dish *D* is partly filled with pure water and so placed under *N* that the water level *L* just covers the mouth of

*N.* *D* may be set into another dish and surrounded with snow and ice or salt. When the lamp is put under the sand bath, the air is first driven out, then the gas appears and dissolves in the cold water. If we substitute a cylindrical bottle of lead or platinum for the dish, the gas itself will condense into a liquid; specific gravity at 12° C. 0.988, nearly equal to water. This liquid boils at 19.5° C. It fumes in the air; the gas combining with the moisture of the air gives the visible fume. Fumes cause violent coughing, and may produce death if taken into the lungs. A drop of the liquid will produce on the skin a white spot, a blister forms, bursts and a painful, slowly healing ulcer forms. Be very careful with this concentrated liquid; but even the dilute water solution has given one much discomfort, when some of it got under the finger nails. The perfectly dry gas does not attack glass; the least moisture causes an immediate attack; the glass becomes *opaque*, we say the fluorspar gas, or liquid, *etches* glass; German *aetzen*, French *mordre* = to bite. To etch means to bite out something.

*Composition of the gas.* When the gas acts upon Na, K, Zn, Fe, two products arise: Hydrogen +  $\text{Me}^n\text{F}^m$  (metallic fluorid). When it acts upon hydroxyd:  $\text{H}^2\text{O} + \text{Me}^n\text{F}^m$  result.



That the gas is a hydrogen compound of the radical F there can be no doubt. But there is a difference about the valence of F. Because there is, as with

the molecule  $\text{H}^2\text{SO}^4$  (see above) the existence of two salts to wit  $\text{NaHF}^x$  and  $\text{NaF}^x$ . In the latter 23 Na (sodium) are combined with 19 fluorine (F), but in the former 23 Na are combined with one H and 38 F. If we admit the proof power of this acid salt (as with the sulfate) then fluorine (F) is divalent, the sodium fluorid is  $\text{Na}^2\text{F}$ , the atomic weight of  $\text{F} = 38$ . There are, on the other hand, quite weighty reasons why we should assume F to be monovalent. Chiefly the closeness between its action and that of chlorine is convincing to the large majority of chemists to declare fluorine a monad and its atomic weight = 19.

Sodium fluorid is thus made  $\text{NaF}$ . The acid salt is  $\text{NaH}(\text{F}^2)$ . The solution of HF in water is hydrofluoric acid. The acid is in all actions closely like HCl, in some instances however fluorids are unlike chlorids. When a water solution of silver sulfate is added to  $\text{HF} + \text{water}$ , no precipitate falls, the  $\text{AgF}$  being easily soluble in water, whilst  $\text{AgCl}$  is insoluble in water. On the other hand, calcium chlorid  $\text{CaCl}^2$  is very soluble in water, calcium fluorid  $\text{CaF}^2$  is insoluble in water.

*The etching process.* We need in the laboratory graduated glass vessels, tubes and flasks; or we want to mark and number the vessels. Hydrogen fluorid, or the acid salt  $\text{NaHF}^2$  is invaluable for this purpose; they can be bought ready for use, being sold either in caoutchouc or *ceresine* bottles (*ceresine* is mineral wax). First cover the glass with a thin film of paraffine or of wax. Melt these materials,



warm the glass and apply the liquid material with a brush. After cooling draw the mark or number upon the wax or paraffine film ; then scratch away the film (with a steel point) all the lines or dots which are to appear in the etching. After this two ways are open : (1) to expose the engraved spot to the slow vapors of HF (this gives the best result); or, (2) to apply the liquid HF by means of a camel's hair brush repeatedly, according to the desired depth of the etching. There is a so-called *glass ink* on the market which appears as a milky white liquid. It is a solution of the acid salt  $\text{NaHF}^2$  mixed with plaster of paris.

*The nature of fluorine.* When HF is made to act upon metallic peroxyds as  $\text{MnO}^2$ , water is produced,  $\text{MnF}^2$  and a mixture of oxygen with fluorine. Many statements by experimenters are on record contradicting each other. The work is exceedingly difficult and expensive. No pure fluorine has as yet been obtained by the above method. This much seems certain, that fluorine possesses a color similar to that of chlorine, *i. e.* yellowish-green, that its odor is similar to that of chlorine, that it attacks all substances except fluorspar, especially platinum and glass, and bleaches indigo. This state of affairs explains why we are uncertain about the valence and the atomic weight.

*The metal contained in fluorspar.* After the mixture of spar powder with  $\text{H}^2\text{SO}^4$  has been heated until HF no longer escapes, but the thick white fumes of  $\text{H}^2\text{SO}^4$ , that means when complete decom-



position has taken place, we find the residue to be a semi-solid which is soluble in a large quantity of water. From the solution precipitate characteristic monoclinic crystals of calcium vitriol = calcium sulfate or gypsum. There may be minute quantities of other metals, which do not now concern us. Calcium is therefore the metal of fluorite, it is *calcium fluorid*,  $\text{CaF}^2$ .

## CHAPTER X.

### LECTURE ON NITER OR SALTPETER.

SALTPETER or niter is the name at present given to a peculiar salt of immense technical or industrial importance. The word saltpeter is the corruption of Latin = *sal petrae* = salt of the rock ; niter is derived from Hebrew-Egyptian = *neter* thence Greek = *natron-nitron*, thence Latin *nitrum*, the meaning of which has been explained above as original for natrium = sodium. At the present time the name applies to something very different from either rock salt or soda. It applies to two salts, *one* forming always prismatic crystals of the orthorhombic system, and *the other* appearing in grains, roughly resembling common salt. On closer examination the granular crystals are seen to possess rhomboid faces not rectangular as in the cube. Geometrically a cube and a rhombohedron are not different except as to the position of the faces in regard to a system of arbitrary axes. It is quite possible that a crystallographic rhombohedron has rectangular faces, and a crystallographic cube has facial angles slightly deviating from  $90^{\circ}$ . The action of a crystal towards the polarized light alone determines the system of crystallization. The angle of the pole edges is  $106^{\circ} 30'$  very near the angle of the cleavage rhom-

bohedron of calcite, which is  $105^{\circ} 30'$ . At all events the obliquity of the angle is large enough to exclude the idea of the cube.

The prismatic niter is known as *potash niter*, the granular rhombohedral variety is known as *Chili niter* or *soda niter*. Both varieties are easily soluble in water, show a *cooling* taste; nevertheless, there is a marked difference in the solubility of the two forms of niter. 100 grams of water dissolve

Soda Niter.		Potash Niter.	
at— $6^{\circ}$ C.	68.8	at $0^{\circ}$ C.	13.3
+ $10^{\circ}$ C.	84.3	+ $18^{\circ}$ C.	29.0
+ $20^{\circ}$ C.	89.5	+ $45^{\circ}$ C.	74.6
+ $100^{\circ}$ C.	168.2	+ $97^{\circ}$ C.	236.0

Between  $0^{\circ}$  and  $100^{\circ}$  C. the ratio of solubility is for soda niter  $\frac{168}{13} = \frac{2}{1}$ ; for potash niter  $\frac{236}{13} = \frac{18}{1}$ . Both niters melt easily. Potash niter, when held in the flame, gives a purple color, soda niter a yellow color to it. About the localities and conditions where and under which the niters are found, we will speak at the end of this investigation, as you will be better able to understand several of the intricate questions which arise in connection with these bodies.

*Investigation of the soda niter.* Let first a crystal fragment be heated in a closed tube. When the heat rises to redness, we notice gas bubbles. On trying the gas we find it to act like oxygen, being without odor, and being able to fan a dark red glowing taper into bright incandescence. Now let

us drop a small piece of feathered tin into the molten niter. Notice the intense action, emission of light and conversion of the tin into white oxyd. Repeat these actions with sulfur, with antimony, with charcoal; in all these cases there is displayed the phenomenon of burning, of combustion. Similar action is displayed by melting potassium chlorate. We may then rightly infer that niter is a salt similar to  $K(\text{ClO}^3)$ , or  $\text{Na}(\text{ClO}^3)$ , but that the salts are not identical follows from several reasons: (1) unlike form of crystals, (2) great difference in solubility, (3) that the chlorates part with their oxygen easily at low temperature, whilst niter only parts with it at red heat, (4) that a brown gas arises from the niter when it burns up a piece of metal, of sulfur or wood, (5) that this brown gas is suffocating. The unavoidable conclusion points to the existence in niter besides sodium and oxygen, of another unknown element. Let this element be designated by N the first letter of niter, and let it be pronounced *nitrogen* = generator of niter; then soda niter will most probably be  $\text{Na}^n\text{N}^m\text{O}^p$ , and of potash niter,  $\text{K}^n\text{N}^m\text{O}^p$ . What is the nature of N? What is the ratio of its combination, what are the numerical values of n, m, p?

(1) *What is the nature of N— of nitrogen?* Let us return first to that experiment in which the niter was heated by itself, yielding oxygen. On acting upon it with water we will get a solution which shows strong alkaline reaction; the niter itself has a neutral reaction—neither acid nor alkaline. This

may mean that  $\text{Na}^2\text{O}$  has formed, and likewise that a compound has formed with less oxygen than the original niter, *i. e.*,  $\text{Na}^n\text{N}^m\text{O}^{p-q}$ . Incidentally we noticed a strong corrosion of the glass tube at the places where the niter had been longest exposed to the flame, which suggests  $\text{Na}^2\text{O}$ , because we know from handling  $\text{NaOH}$  and  $\text{Na}^2\text{CO}^3$  in the glass tubes that these bodies attack the glass at high heat. In the water solution would either be  $\text{Na}(\text{HO}) + \text{Na}^n\text{N}^m\text{O}^{p-q}$  or only one of the two. Addition of dilute  $\text{H}^2(\text{SO}^4)$  will neutralize  $\text{Na}(\text{HO})$ .

$2\text{NaHO} + \text{H}^2(\text{SO}^4) = \text{Na}^2(\text{SO}^4) + \text{H}^2\text{O} = \text{neutral}$ ; a further addition of  $\text{H}^2(\text{SO}^4)$  will cause the evolution of a gas of peculiar odor, rather aromatic. The gas *may* be  $\text{H}^2\text{N}^m\text{O}^{p-q}$  or not; at any rate it is a peculiar body. Now let us act with concentrated  $\text{H}^2(\text{SO}^4)$  upon the original niter. At ordinary temperature there is but little if any action, except that the niter seems to dissolve, at least partly, and but a faint odor is noticed. As heat is applied, effervescence ensues. A sour, pungent gas appears, which condenses in a sufficiently cold receiver into a liquid, or else is energetically absorbed in water. In the residue we have sodium hydrogen sulfate + hydrogen sulfate. We pour it into a porcelain dish, and may convert it into salt cake by means of heat; prove it to be  $\text{Na}^2\text{SO}^4$  by means of its easy solubility in water and its resistance to crystallization. Only low temperature will induce crystals. The liquid distillate we will name *spirits of niter*. We study its action upon the metals, upon paper, wood,

the skin, wool, in fact upon all bodies known to us and handy to procure. For remember always that chemistry means try anything upon everthing else. All the actions will be remarkable.

Lead (Pb) + sp. niter + heat = white salt + brown fumes.

Copper (Cu) + sp. niter = blue salt + brown fumes.

Silver (Ag) + sp. niter = white salt + brown fumes.

Filter paper + sp. niter + heat = solution + brown fumes.

Gold (Au) + sp. niter + heat = no action.

Platinum (Pt) + sp. niter + heat = no action.

Tin (Sn) + sp. niter = white oxyd + brown fumes.

Hydrochloric acid (HCl) + sp. niter + heat = brown liquid + brown fumes.

HCl + sp. niter + gold + warm = yellow solution,  $\text{AuCl}^3$  + brown fumes.


HCl + sp. niter + platinum = yellow-brown solution + brown fumes.

The spirits of niter proves itself thus one of the most powerful agents. The Arab chemist Geber was the first to mention this body. The Latin translation of his works speaks of it as *aqua fortis* (the strong water) or *aqua dissolutiva* (the dissolving water) because it dissolved both silver and lead. But the combination of the spirits of salt (HCl) with the spirits of niter went, and still goes, by the name *aqua regia* (the kingly water), because it dissolves gold, the very king of the metals.



On the other hand, if we dilute first the spirits of niter with water, very considerably, then we get *hydrogen* when acting upon either iron or zinc, but not with lead, copper or any other metal; with all of these it is either brown fumes or nothing. Remember the similarity with the actions of oil of vitriol or of the concentrated sulfuric acid. Concentrated acid on the metals gave vitriols and  $\text{SO}^2$ ; diluted acid on iron or zinc gave vitriols and  $\text{H}$ . Just as  $\text{SO}^2$  was demonstrated as an oxyd with less oxygen than the sulfur oxyd which constitutes the sulfuric acid, so it follows logically that in the action of the strong spirits of niter, the brown fumes must

FIG. 44.



constitute a lower oxyd of the nitrogen, the element whose properties we are after. But if thus the metals can take away oxygen from the nitrogen, we argue, at a temperature below even the boiling-point of water, will it not seem probable that at a still higher heat more will be taken, or perhaps even all?

We set up an apparatus as shown in Fig. 44. *F* is a small flask with twice perforated stopper.

Through the latter pass the stem of a funnel 2 and the delivery tube 3. In *F* we place finely divided copper, *K* (gauze, granules, chips). The funnels hold the diluted spirits of niter. 3 connects with U-tube 4. The latter is partly filled with concentrated  $\text{H}^2\text{SO}^4$  forming a trap to dry and control the escaping gas. 5 is a tube filled with quick-lime ( $\text{CaO}$ ) between two cotton plugs. In the charcoal furnaces *F'*, *F'*, lie the hard glass tubes *T*, *T'*, each charged with rolls of copper wire-gauze. By means of rubber tube 6, *T'* connects with the bell jar *B*, which is filled with boiled water (in order to expel any absorbed air). Before connecting *F* with tube 7 we heat up the furnaces and pass hydrogen through *T*, *T'*, in order to have perfect metallic surface on the copper. The tube 5 will act as dryer and will also retain any gas of an acid nature. While the action of hydrogen was going on in *T*, *T'*, we have utilized time by starting the action in *F* so that all the air is driven out by the gas. We regulate the flow of liquid from the funnel, so that, if possible, the sulfuric acid trap in 4 will indicate the passing of a slow current of gas. The slower and steadier (not in gulps) the current, the better will be the chances of a perfect deoxydizing action. Before connecting *T* with tube 5 by a rubber tube 7 we remove most of the charcoal from *F*, *F'*, let the tubes *T*, *T'* come down below red heat, for the unknown niter gas, being an oxyd might produce with the hydrogen an explosive mixture. When we connect 7 with *T'* we wait a sufficient time to let the hydro-

gen be displaced by the niter gas, then we replace the charcoal, get a good heat, cherry red, being always careful to protect the rubber stoppers in  $T$  and  $T'$  by guarding shields and dropping water, and again after some minutes' wait, we connect the rubber tube  $C$  with the bell jar  $B$ . In the position as shown in the figure there will be suction through the chain of apparatus as soon as the stopper  $\delta$  is opened, owing to the difference of level,  $h$ , between the water inside and outside the bell. Therefore all stoppers and connections must have been made airtight, otherwise air will be sucked into the apparatus: The *true nature* of nitrogen will be masked. We maintain the action until  $B$  is filled with gas, or until several holders shall have been filled.


*Properties of nitrogen.* After having been produced, as just stated, the gas possesses the characters of an element. With present means, it cannot be further split. A gas devoid of color, odor, taste. Specific gravity 14 ( $H=1$ ); 0.9674 ( $Air=1$ ). This specific gravity is so nearly the same as that of the *azote*, the nonrespirable part of the air, this being 0.9713, that we are justified in declaring *azote* = nitrogen, because the nitrogen also is *nonrespirable*, it causing death by suffocation. The gas is but very slightly soluble in water, it is not absorbed by the alkalies. It is very indifferent towards all agents, and yet it is evident that under certain conditions it may be made to unite with oxygen in several ratios, and also with hydrogen giving rise to a most interesting body. Nitrogen we shall find in

all animal and plant bodies, constituting the essential constituent of *protoplasm*, the body which is at present taken to be the basis, the substratum of all life. It is important to note that nitrogen has no property by which we can at once identify it, except the specific gravity; all other properties are mere negations of the properties possessed by other gases.

*Properties of the spirits of niter and its quantitative composition.* Acting with the spirits of niter upon a metallic oxyd (MeO) we get water + niter. The most suitable metallic oxyd for our present purpose is lead oxyd PbO which has the pale yellow color; the red oxyd is not equally suitable. Acting upon Na(OH) or K(OH) we restore the original niter, either soda niter, or potash niter and water. There is only one salt formed, no acid salt having been obtainable. Hence the radical contained in the spirits is there combined with *one* H, the radical is a monad, and will, therefore, be represented by the symbol



Hence, also, if we act with the spirits upon lead oxyd, the reaction must be




Like the soda and potash niter, the lead niter contains no water, except a trifle with the mother liquor. The white or colorless crystals of this lead niter are rhombohedral, hence isomorphous with the soda niter. It is best, because most rapid, to

use the spirits diluted, because the lead niter is not soluble in the concentrated spirits. When all of the lead oxyd is thus dissolved, or when the liquid will not further dissolve the oxyd, filter and evaporate the filtrate to complete dryness. The residue is then pure lead niter. Heated in a crucible or glass tube, the lead niter breaks up into brown gas and yellow lead oxyd. The brown gas is very acrid and suffocating, yet it will act upon a glowing taper like oxygen. In fact we can readily prove the brown fumes to have an admixture of oxygen; therefore



This action opens the way for a quantitative determination of the ratio existing between Pb, N and

FIG. 45.



O, if we arrange the conditions of the experiments in such a way that the volume of nitrogen can be accurately measured, which results from the decomposition of say one gram of the lead niter.

Let *T*, Fig. 45, be an ample hard glass tube fitted



with stoppers. 8 is a porcelain boat containing 0.5 gram of lead niter. 9 is a clean roll of copper gauze. 7 is a U-tube filled with pieces of pumice and  $\text{H}^2\text{SO}^4$  (to retain moisture). 6 is a smaller U-tube with enough  $\text{H}^2\text{SO}^4$  to form a trap. *H* is the holder filled with lime gas and the water in *B* furnishes the pressure to drive out the gas from *H*. 10 is a gas burette and 11 the cylinder to regulate the pressure. Both cylinders are filled with solution of sodium hydrate. All stoppers and connections being tight, we first displace all air from the apparatus, because four-fifths of it are nitrogen; then heat the copper gauze to redness, while we protect the boat from the heat by the shield 12. When the gauze is glowing we move shield 12 to the left, from time to time, so that the decomposition of the lead niter shall be slow and gradual; but at length the entire tube is at redness up to the shield. As the niter decomposes the oxygen goes to the copper and the nitrogen passes into the burette, pushing before it the lime gas. As soon as the contents of the boat are pure dark yellow or red, we open the stop cock 4 and drive all niter gas into the burette. Then, closing the latter's stopcock, we shake the gas with the liquid, thus absorbing all the lime gas into the  $\text{Na}(\text{OH})$  solution, and then read the volume of gas. Let  $V'$  cubic centimeters be the volume of the gas, measured under the pressure of the atmosphere at  $20^\circ \text{C}$ , the pressure of the atmosphere, measured by the mercury column of the barometer, be *B* millimeters. As the gas is satur-



ated with aqueous vapor, being over water, *i. e.*, a dilute solution of  $\text{Na}(\text{HO})$ , the tension  $T$  of the aqueous vapor increases the pressure  $B$  and must therefore be subtracted, because we wish to get at the volume of the *dry gas*. The expansion of the air for one degree C. is 0.00367 of its volume, hence the volume  $V^\circ$  of dry gas at  $0^\circ$  C. and sea level barometer, 760 mm., will be

$$V^\circ = \frac{V'(B - T)}{(1 + 0.00367t) \times 760} = 33.67 \text{ c.c.}$$

(The tables calculated by Prof. Leo Liebermann are most convenient in such calculations.) If the weight of one c.c. of dry nitrogen at  $0^\circ$  C. and 760 mm. be 0.001256 gram, then

$$V^\circ \times 0.001256 = G,$$

will be the weight of the nitrogen contained in 0.5 gram of lead niter,  $\text{Pb}(\text{N}^\text{m}\text{O}^\text{p})^2$ .  $G$  equals 0.0423 gram and the weight of lead oxyd,  $\text{PbO}$ , is 0.337 gram (found by weighing boat after operation). Then we will have

$$\text{PbO} = 0.3370$$

$$\left. \begin{array}{l} \text{N} = 0.0423 \\ \text{O} = 0.1207 \end{array} \right\} = 0.163 \text{ gram} = \text{wt. of the N. oxyd.}$$

---


$$0.5000$$

The weight of oxygen if found by difference. The volume weights of oxygen and nitrogen (found by direct weighing) are 16 and 14. Hence we will obtain the atomic ratio of the two elements by dividing gram weights of N and O by 14 and 16 respectively.

$$\frac{0.0423}{14} = 0.00302; \frac{0.1207}{16} = 0.00754$$

$$\frac{302}{754} = \frac{2}{5}, \text{ hence } \text{N}^2\text{O}^5$$

We have directly proved that lead niter is a combination of the oxyd PbO with the oxyd  $\text{N}^2\text{O}^5$ . Above we showed the probability of the radical  $\text{N}^m\text{O}^p$  being a monad. In the lead niter there are then two molecules of the nitric radical.

$\text{PbO}.\text{N}^2\text{O}^5$  becomes  $\text{Pb}(\text{N}^2\text{O}^6)$  or  $\text{Pb}(\text{NO}^3)^2$ .

For the sake of uniformity we will designate hereafter the niters by the word *nitrate*, thus:

Hydrogen nitrate =  $\text{H}(\text{NO}^3)$  = nitric acid = aqua fortis.

Sodium nitrate =  $\text{Na}(\text{NO}^3)$  = soda niter = Chili niter.

Potassium nitrate =  $\text{K}(\text{NO}^3)$  = potash niter = common niter.

Calcium nitrate =  $\text{Ca}(\text{NO}^3)^2$ .

Silver nitrate =  $\text{Ag}(\text{NO}^3)$ .


Lead nitrate =  $\text{Pb}(\text{NO}^3)^2$ .

Copper nitrate =  $\text{Cu}(\text{NO}^3)^2$ .

Ferric nitrate =  $\text{Fe}(\text{NO}^3)^3$ .

All normal nitrates are *soluble* in water, whilst some sulfates, some chlorids, bromids, iodids, fluorids are *insoluble* in water.

Sol.  $\text{Ag}(\text{NO}^3) + \text{sol. NaCl} = \text{insol. AgCl} + \text{sol. Na}(\text{NO}^3)$ .




These two reactions serve us as tests for *soluble* chlorids and sulfates respectively.

Of all bodies we have investigated, the nitrates appear to me the most wonderful. The very same elements in which we breathe and lead a more or less harmless life, the existence of which elements we are not even ordinarily aware of, become violently active in the form of nitrates. The chlorates, though acting similarly, are not so astonishing, because in them we find chlorine, a violently offensive body by itself. A rather rough simile may bring nearer to your grip of imagination this action of the nitrates. Let the atoms be imagined as spiral springs (watch spirals). In the atmosphere the nitrogen molecules lie alongside of the oxygen molecules as uncoiled springs, inert, inoffensive things. A powerful shock strikes the inert, uncoiled bodies, say the electric spark of a thunder storm; the shock causes the springs to coil up, and the affinity of a strong basic oxyd, as  $\text{K}^2\text{O}$ ,  $\text{Na}^2\text{O}$ , lies handy as a binding rope of the springs, as shown in Fig. 46; the nitrate molecule is achieved; the potential energy of the coils is restrained by the thin band of affinity. Now let the nitrate molecules be brought into intimate contact with other molecules which possess a stronger affinity for oxygen than the nitrogen, for instance, carbon molecules, at a red heat. We may even carry the picture further and say the thermic energy expands the springs,

straining them against the restraining bond until at red heat this bond snaps, giving way to carbon—oxygen attraction. With the breaking of the

FIG. 46.




bond, the oxygen springs uncoil and display an extraordinary energy, such as we are forced to admire in gunpowder.

#### COMPOSITION OF GUNPOWDER OR BLACK POWDER.

We saw in a small experiment how the mixture of niter and charcoal powder flew out of the glass tube with a flash. The products of that action were  $\text{Na}^2(\text{CO}^3)$ , sodium carbonate, N, nitrogen and  $\text{CO}^2$ . The two latter gases, through their expansion, impart to the explosion its propelling or its tearing, splitting effect. The more gas, the greater the effect from a given mixture. But in forming  $\text{Na}^2\text{CO}^3$  much gas passes into the solid state and lessens the effect of the powder. It was soon found that a greater effect could be obtained by mixing with niter and charcoal a certain quantity of sulfur. This was all arrived at by those patient experimenters without knowing even that the production of gas was the chief object. They mixed sulfur

with the powder on general principles that it would be a good thing, because sulfur was a very peculiar and mysterious body. We modern chemists, who experiment less and think more, know why the sulfur increases the effect. This is the theoretical picture of the explosion :



By weight

$$\begin{array}{rccccccc} 2(39 + 14 + 48) & + & 32 & + & 3 \times 12 & = & 2 \times 39 + 32(\text{solid}) \\ \underbrace{\hspace{1.5cm}} & & & & \underbrace{\hspace{1.5cm}} & & \underbrace{\hspace{1.5cm}} \\ 202 & + & 32 & + & 36 & = & 110 \\ & + & 2 \times 14 & + & 3(12 + 32) & & \\ & \underbrace{\hspace{1.5cm}} & & \underbrace{\hspace{1.5cm}} & & & \\ & + & 28 & + & 132 & & \\ \text{KNO}^3 & = & 202 & = & 74.81 \% & 75 \text{ potassium niter} \\ \text{S} & = & 32 & = & 11.85 & 12 \text{ sulfur} \\ \text{C} & = & 36 & = & 13.34 & 13 \text{ fine charcoal} \\ \hline & & 270 & & 100.00 & & \end{array}$$

On the second side of the equation stand

$$\begin{array}{rccccccc} \text{K}^2\text{S} & = & 110 & = & 40.70 \% & (\text{solid}) & 41 \text{ solid} \\ 2\text{N} & = & 28 & = & 10.40 \% & \text{gas} & 10 \} \text{gas} \\ 3\text{CO}^2 & = & 132 & = & 48.90 \% & \text{gas} & 49 \} \\ \hline & & 270 & & 100.00. & & \end{array}$$

59 per cent. of the powder is converted into gas. One gram of the powder gives by the explosion 0.1 gram of nitrogen, 0.49 gram of carbon dioxyd. At 0° C. and 760 mm., 0.001256 gram of nitrogen occu-

pies the space of one c.c.; 0.1 gram of nitrogen occupies the space of 79.6 c.c. 0.001977 gram  $\text{CO}^2$  occupies the space of 1 c.c.; 0.49 gram  $\text{CO}^2$  the space of 247.3 c.c. The volume of gas produced by the explosion of one gram of perfect black powder is at  $0^\circ \text{C}$ ,  $247.3 + 79.6 = 326.9$  c.c. One gram of best powder, in small but perfect angular grains, occupies the space of 0.9 c.c.; the surface of 1 c.c. being six square centimeters. Hence if one gram of such powder be filled into a cartridge and a bullet be pressed tightly upon the powder, a space will be filled possessing six times  $0.9 = 5.4$  square centimeters. After explosion this same space will be filled with 326.9 c.c. of gas which will press upon the enclosing surface with  $\frac{326.9}{0.9} = 363.2$  times the pressure of the atmosphere or upon 1 square centimeter with  $\frac{363.2}{5.4} = 67.26$  atmospheres. But there is another very important factor; the heat generated by the explosion, which expands the gas mercilessly, and if the enclosure be rigid, exerting an ever-increasing pressure. The gases expand, within certain limits so nearly alike, that one coefficient answers for all. This coefficient is  $\frac{1}{273}$  or 0.00367 volume for  $1^\circ \text{C}$ . By the following reasoning we arrive at the theoretical temperature produced by the explosion of one gram of powder: 1 gram carbon through the oxydation into  $\text{CO}^2$  produces heat equal to 8050 calories (the heat would raise the temperature  $t$  of 8050 grams of water by one de-



gree C. Hence 0.13 gram of carbon will produce  $8050 \times 0.13 = 1046$  calories. By the burning have been produced 0.49 gram of  $\text{CO}^2$ , 0.1 gram of N and 0.41 of  $\text{K}^2\text{S}$ . Each of these bodies has a capacity for heat to be swallowed up before the heat can be felt. The temperature, the sensible heat, must therefore be directly proportional to the absolute heat—the 1046 calories, and inversely to the absorbed heat. In a general way

$$T^{\circ} = \frac{A}{a}$$

in which  $A$  = absolute heat,  $a$  = weight of products into their respective specific heats.

In our special case

$$T^{\circ} = \frac{1046}{0.499 \times 0.22 + 0.1 \times 0.24 + 0.41 \times 0.4}$$

$$T^{\circ} = \frac{1046}{0.108 + 0.024 + 0.164} = \frac{1046}{0.296} = 3534^{\circ}\text{C}.$$

The  $a$  in the denominator is represented by the sum of the products of the combustion of the powder; each member multiplied by its factor representing the unit of heat capacity or specific heat.

Thus it is seen that by the combustion of one gram of powder a temperature is generated equal to  $3534^{\circ}\text{C}$ ., higher than that of an intense coal fire. At this temperature the 326.9 c.c. of gas must have expanded to  $326.9 \times 0.00367 \times 3535 = 4236.6$  c.c., to 12.9 times their volume at  $0^{\circ}\text{C}$ ., or 871.7 atmospheres pressure per sq. cm., provided the law of expansion holds good at such high temperature, but

this is by no means certain. At any rate, the theoretical picture gives a measure for the actual phenomenon. We see that the pressure of the gas must be enormous, though less than the theoretical, in fact only about  $\frac{1}{2}$  as shown by actual experiment, made for the military departments of several governments. Such experiments are known by the adjective *ballistic* (made with a ball). The reasons are several. In the first place the burning of the powder is never complete. 2. The reactions are not quite like the formula indicates. A part of the oxygen remains fixed by forming  $K^2SO^4$ , hence some of the carbon only burns to CO, whilst some of the nitrogen remains as NO. 3. The material of the apparatus—a gun for instance, or a mortar—is somewhat elastic.

As engineers we should know all that has here been given, to understand the effects of the blasting powder we use. I give you only the chemical facts connected with the matter.

*Other powders.* The white and smokeless powders all contain a nitric radical, either  $NO^2$  or  $NO^3$ . In principle there is no difference between them and black powder. Their special compositions will be dealt with under the subject of carbon compounds.

#### INVESTIGATION OF THE BROWN FUMES OR NITROUS FUMES.

These always form, when  $H(NO^3)$  acts upon an oxydizable substance, as copper for instance, or  $SO^2$ .


Acting with  $H(NO^3)$  upon copper in a test-tube

we observe the tube filled with the brown fumes so long as the action continues. But if the tube be stoppered with a narrow tube for the escape of the gases, a different action ensues. We notice that the gas inside the tube turns lighter by degrees, at last being quite colorless, but at the mouth of the escape tube a steady cloud of dense, brown-colored gas is visible all the time. This at once suggests the presence of two different gases within the brown fumes. More precisely we would say: Through the action of  $\text{HNO}^3$  upon an oxydizable body is generated a colorless gas—an oxyd of nitrogen. When this gaseous oxyd comes together with air the *brown gas* forms by combination of the air-oxygen with the colorless gas; but the brown gas itself must possess a ratio of oxygen to nitrogen *less* than  $5/2$ , for if it is absorbed in ice water we find produced a mixture of *spirits of niter* and the *same* acid that was made by acting with  $\text{H}^2\text{SO}^4$  on the residue left after heating niter. All this will be proved presently. The brown gas becomes colorless if mixed with an oxydizable body, as  $\text{SO}^2$  for instance, evidently by loss of oxygen. Hence the *colorless gas* must have an oxygen to nitrogen ratio *less* than the brown gas.


*Nitrogen dioxyd*,  $\text{NO}^2$  or  $\text{N}^2\text{O}^4$ , *hyponitric anhydride* the *nitrous fumes*. Let the tube *T*, Fig. 47, be partly filled with dry lead nitrate and placed in furnace *F*. Make the U-tube *U* from a  $\frac{1}{4}$ " tube, draw it out into a capillary at *C*, *a* and *b*, then surround it in the basin *B* with a mixture of ice and coarse common salt; stick a thermometer *t* into the

mixture. If the salt be kept on snow or ice until its temperature falls to  $0^{\circ}\text{C}$ ., and if it be then mixed with one volume of snow or fine-cut ice, the temperature of the basin will drop to  $-20^{\circ}\text{C}$ . Make

FIG. 47.



connection at *C* with *T*, and heat *T* to redness gradually. We know from previous experiment (composition of  $\text{N}^2\text{O}^5$ ) that the  $\text{Pb}(\text{NO}^3)^2$  breaks up into  $\text{PbO} +$  brown fumes. Thus




Passing into *U* the fumes condense into a brown liquid and at *6* issues a gas which sustains combustion with energy—(oxygen). At end of decomposition close *U* at *C* and *a* with the blow-pipe. If the U-tube be changed twice, then you find in the third tube colorless crystals. These represent the true substance. The crystals melt at  $-12^{\circ}\text{C}$ . to a colorless or slightly yellow liquid. It follows that no crystals result if the temperature of the freezing mixture be not at the least  $-15^{\circ}\text{C}$ . As the liquid is heated by the warmth of the hand it becomes more and more highly colored, giving out dense brown fumes and reaches a constant boiling-point

at  $+22^{\circ}$  C. (in the hand, because the temperature of the blood is  $33^{\circ}$  C.). On the addition of ice water drop by drop, the liquid turns first green, then blue, then colorless. It acts upon oxydizable bodies more energetically than  $\text{HNO}^3$ , because the restraining bond of the hydrogen is removed. Mention has already been made of the suffocating action of the brown fumes. The living organism tries to avoid the danger lurking in the breathing of the gas, for the muscles of the epiglottis contract instantly when the gas comes in contact with their nerve ends. At Berlin, some years ago, fire broke out in the yard of a large chemical works. Sixty carboys of aqua fortis were stored under one shed. They exploded, one by one. The acid flowing into the straw and wood of the packing let out an immense volume of brown fumes. Five of the firemen who had been endeavoring to save the carboys, went back to the station with the others, smoked a pipe and went to sleep in their bunks. Within a few hours they woke in convulsions and died shortly after, in great agony. Let this be a warning to be careful. Do not act upon metals, or sulfids with  $\text{HNO}^3$ , except in a well-drawing hood; for even if death does not ensue, there may be permanent injury to the bronchiæ and their capillary ramification in the lungs, from ulceration of the mucous membrane. The composition of the brown gas we find similarly to that of the lead niter. We make a small U-tube *U*, Fig. 48, drawn out into capillary thread at *a*, *b*. We take the weight and then fill into it a



few drops of the liquid, by pressure or by suction. In *T* there is the coil of copper wire. At *B* is the burette for measuring the gas, filled with solution  $\text{Na}(\text{HO})$ , and which is immersed in a dish of water

FIG. 48.




when in use. We fill the tube *T* with lime gas, insert *U* at *b* into the stopper and *a* into rubber tube leading to lime gas holder and furnished with clamps *C*; then connect the burette *B*. We bring *T* up to redness and turn on the lime gas in a gentle current. The radiant heat will suffice to volitalize the liquid oxyd of nitrogen and the lime gas will carry it over the copper gauze. Nitrogen and lime gas pass into *B* and the  $\text{NaHO}$  solution will absorb the lime gas. Thus we get the nitrogen volume which we deal with as in the previous experiment; *w* being weight of oxyd, *n* being the weight of nitrogen,  $w - n = \text{O}$ , the weight of the oxygen. Dividing *n* by 14 and  $w - n$  by 16, we get the ratio

$$\frac{n}{14} : \frac{w-n}{16} = 1 : 2 = \text{NO}^2 \text{ or } \text{N}^2\text{O}^4 \text{ or } \text{N}^4\text{O}^8$$

Some investigators claim the ratio  $\text{N}^4\text{O}^8$  which can be interpreted as  $\text{N}^2\text{O}^3.\text{N}^2\text{O}^5$  a combination of the two oxyds, on the ground that when ice water be



mixed with the liquid oxyd it breaks up into the two acid bodies:  $\text{H}(\text{NO}^2)$ , *hydrogen nitrite* and  $\text{H}(\text{NO}^3)$ , *hydrogen nitrate*: to wit



However, this view, which looks at the molecule as a *polymeric* molecule, *i. e.*, 4 times  $\text{NO}^2$ , only applies at low temperatures, for Dulong found for the brown gas the specific gravity 1.62, and this corresponds to

$$\frac{1}{2} \text{ vol. nitrogen} = 0.4856$$

$$1 \text{ vol. oxygen} = 1.105$$

$$\hline 1.5906$$


so nearly 1.62 that we can have no doubt left. The new gaseous molecule is surely  $\text{NO}^2 = 2(\frac{1}{2}\text{N} + \text{O})$ . 1 vol. N + 2 vols. O = 3 vols., become 2 vols.  $\text{NO}^2$ ; there is a condensation of 3:2 of  $1\frac{1}{2}:1$ .

When the brown fumes are passed into  $\text{H}^2(\text{SO}^4)$ , the fumes become absorbed, and if the operation be continued for a time, colorless crystals form in the acid. The crystals are obtained more readily by pouring a few cubic centimeters of  $\text{H}^2(\text{SO}^4)$  into a small flask and by spreading the liquid, through rotation, upon the entire glass surface. If now the brown fumes are brought into the bottom of the flask, under the rotation the brown fumes become absorbed, a crystalline crust resembling ice forming all over the flask (inside). The composition is not definitely settled, probably  $2\text{H}^2(\text{SO}^4).\text{NO}^2$ . The reaction is of much economical importance in the manufacture of  $\text{H}^2(\text{SO}^4)$  on a large scale and will be brought forward when we shall arrive at that process.

The colorless gas, nitrogen monoxyl, nitric oxyd,  $NO$ . This body becomes generated whenever  $H(NO^3)$  acts upon oxydizable bodies, metals, metallic sulfids, paper, wood, sulfur dioxyd. As soon as it comes into contact with the air it changes into nitrogen dioxyd  $NO^2$  (brown nitric fumes). We obtain this gas very pure by passing  $SO^2$  into warm  $HNO^3 +$  water.

In flask  $F$ , Fig. 49, we generate  $SO^2$  from the

FIG. 49.




mixture  $M$ , which is copper gauze and  $H^2SO^4$ , heated by a flame, according to equation



Through tube  $1$  gas passes into the washing tube  $2$ , partly filled with  $H^2(SO^4)$  (moisture is retained and flow of gas can be regulated).  $W$  is a so-called Will's condenser, possessing the three bulbs,  $3$ ,  $4$ ,  $5$ , and containing  $2H(NO^3) + 2H^2O$ .  $B$  is a basin holding warm water to heat the  $H(NO^3)$ ; through tube  $6$  the gas can be delivered into any suitable vessel. In the figure this vessel is a knee-tube  $T$  filled with mercury and standing in mercury trough  $Q$ .  $T$  is held by stand  $S$ . When the gas  $SO^2$  bubbles into the  $H(NO^3)$  it becomes oxydized into

$\text{H}^2(\text{SO}^4)$  and  $\text{H}(\text{NO}^3)$  becomes changed into the nitrogen monoxyd.



Do not think the composition of the gas must be NO, because the equation demands it, many students, and even some professors, have that belief. I wrote the equation because I know the gas to be NO. I could have balanced the equation in several other ways, merely by changing the coefficients of  $\text{SO}^2$  and of  $\text{HNO}^3$ .

*Proof of the composition of nitrogen monoxyd.* Suppose we have allowed to enter into the knee-tube T, Fig. 49, about 10 c.c. of the gas and have marked this volume by a sticker. Banking upon the known maximum of affinity of potassium for oxygen, we will introduce, by means of a thin copper wire, a piece of metallic potassium at P, and heat the metal with a burner. A flash of light and a violent commotion of the gas accompany the act of deoxydation of the gas. I must hold the tube firmly, with the left hand, to prevent its being raised above the mercury level in Q. When the tube has resumed the temperature of surrounding air we find the volume of gas shrunk to exactly  $\frac{1}{2}$ ; hence there is in one volume of nitrogen monoxyd  $\frac{1}{2}$  vol. N +  $\frac{1}{2}$  vol. O or made into full units 1 vol. N + 1 vol. O, the symbol of the gas is NO. There is no contraction in the union.

Specific gravity of NO found by weighing 1 vol.  
= 1.0379 (air = 1)

$$\frac{1}{2} \text{ vol. N} = 0.4856$$

$$\frac{1}{2} \text{ vol. O} = 0.5525$$

$$\hline 1.0381$$


Calculated specific gravity equal to the experimental, hence *NO* represents the *molecular volume* of the gas.

*Properties of NO.* The gas cannot occur in nature. Why? Because it changes to  $\text{NO}^2$  on meeting the oxygen of the air. For the same reason we *do not know* whether it has odor or taste. Does not act on litmus paper. Its actions on the breathing organs are the same as those of  $\text{NO}^2$ , for the same reason as above. If the gas be conducted into a solution of copperas  $\text{Fe}(\text{SO}^4) + 7\text{H}^2\text{O}$  or  $\text{Fe}(\text{Cl}^2) + 2\text{H}^2\text{O}$  the solution turns *dark brown*, or even *inky black*. The gas is *completely absorbed*. Distinction from all other gases. This reaction enables us to recognize and identify a nitrate in an unknown mixture, even very minute quantities of the nitrate. Proceed as follows with this test: Bring the unknown solution (1 c.c.) into the bottom of a test-tube *T*, Fig. 50, add 2 c.c. of concentrated  $\text{H}^2(\text{SO}^4)$  and mix. Then take into a glass tube *t*, which has been drawn out to a capillary, a strong solution of copperas (ferrous sulfate)  $\text{Fe}(\text{SO}^4)$ . Lower *t* into *T*, so that the point just touches surface of the liquid mixture, and let run out a few cubic centimeters. The two liquids are then unmixed, in two superimposed strata. Within a short time a dark ring will develop at the contact of the strata. If no dark ring develops then the unknown substance does *not contain any nitrate*.

The student should practice this reaction using as unknown a solution which was made up from one drop of strong  $\text{H}(\text{NO}^3)$  and 50 c.c. of water (approximately 0.2 per cent.  $\text{HNO}^3$ ).


*Nitrites*,  $\text{Na}(\text{NO}^2)$ ,  $\text{K}(\text{NO}^2)$ ,  $\text{Ag}(\text{NO}^2)$ . *Di-nitrogen-trioxyd*,  $\text{N}^2\text{O}^3$ . Both  $\text{Na}(\text{NO}^3)$  and  $\text{K}(\text{NO}^3)$  loose oxygen when heated at low red heat and more rapidly at bright red heat. Since neither the metal

FIG. 50.




nor nitrogen is given off, the ratio between the three elements must become changed; a new body forms. This we can demonstrate readily in two ways: (1) by acting upon the residue with concentrated  $\text{H}^2\text{SO}^4$  when copious brown fumes are given off; (Recall that the nitrates +  $\text{H}^2\text{SO}^4$  do not give


any fumes.) (2) by adding solution of  $\text{Ag}(\text{NO}^3)$  to the water solution of the residue—a white precipitate falls out, which is not very soluble in water. This silver salt does not contain water, and by its decomposition we can find the ratio of Ag, N, O, by using the apparatus and method followed with the lead nitrate. Let the silver salt, which shall be named *silver nitrite* (note the substitution of the letter i for the letter a in nitrate) be  $\text{Ag}(\text{N}^m\text{O}^{p-q})$  then we shall obtain by the decomposition



Let the fumes be decomposed by copper gauze at red heat and you get the nitrogen.




Ag : N : O = 1 : 1 : 2 hence the formula of the nitrite is



The hydrogen salt cannot be made, it is *too unstable*.

*Preparing K(NO<sup>2</sup>).* Heat niter in an iron pot or crucible to melting heat, then add 2 parts of metallic lead for each part of niter. The affinity of lead for oxygen helps the decomposition of the niter :




Molecular weight  $\text{K}(\text{NO}^3) = 101$ , atomic weight of  $\text{Pb} = 207$ , hence 1 part niter + 2 parts lead, correspond to theory. In practice, however, the reactions are not complete. Some lead is apt to remain



unoxydized, some oxygen escapes, the residue is, usually,  $K(NO^2)$  mixed with  $PbO$ ,  $KNO^3$  and  $K^2O$ . Dissolve the fused mass in little boiling water and let cool. The  $PbO$  will settle, the  $K(NO^3)$  will crystallize. Decant (that is pour off) the liquid and evaporate to dryness, finally heat to melting and pour into pencil moulds, same as used for  $NaOH$  and  $KOH$ .  $K(NO^2)$  is a reagent in use for the separation of cobalt from nickel, as well as for other operations. The salt usually shows an alkaline reaction from  $KOH$ , can be neutralized with dilute acetic acid.



In the nitrites, we have undoubtedly a peculiar oxyd of nitrogen. The radical ( $NO^2$ ) is not that oxyd, its reactions are quite different. This oxyd is  $N^2O^3$ , for  $K^2O.N^2O^3 = 2K(NO^2)^2$ . This oxyd—*dinitrogen trioxyd*—is contained in the brown fumes, as gas. It may be condensed at  $-15^\circ C$ . into a deeply blue liquid, which boils even at the freezing-point of water. A *blue solution* results from the action of  $NO$  gas upon a solution of  $H(NO^3)$  in water (specific gravity = 1.25) a *green solution*, when the specific gravity is 1.35 (because this liquid then contains both  $N^2O^3$  and  $N^2O^4$ , ( $2NO^2$ )). The same colors result when acids of the given specific gravities act upon certain oxidizable bodies. Any one not knowing this fact will often waste much time trying to find copper (blue nitrate) because he sees a blue solution, or chromium because he sees a green solution when acting on *galena*, the lead sulfid, for instance. The student should get this information by experiment to fasten it the more firmly.

If a solution of  $\text{Na}(\text{NO}^3)$  or  $\text{K}(\text{NO}^3)$  be shaken with amalgamated zinc (zinc coated with mercury) the solution will then contain *nitrite*, to be detected by the addition of the liquid to starch paste containing potassium iodid and some free  $\text{H}^2\text{SO}^4$  (dilute). The paste turns *blue*, because



The rain water from a thunder shower will give this reaction, showing that it contains *nitrite*; azote and oxygen have become united by the electric discharges of the storm.

*Dinitrogen monoxyd*,  $\text{N}^2\text{O}$ , *nitrous oxyd*, *laughing-gas*. This gas arises when the monoxyd  $\text{NO}$  is allowed to stand in contact with easily oxydizable substances; finely divided zinc, iron filings, sulfites ( $\text{Na}^2(\text{SO}^3)$ ), and many others. One-half of the oxygen is removed, or 2 molecules  $\text{NO}$  furnish one molecule  $\text{N}^2\text{O}$ .



We proceed to prove this by acting with heated potassium upon the gas in a knee-tube, exactly as described for  $\text{NO}$ . The action is quite as energetic. After cooling the volume is the same as before the action. Hence one volume gas contains one volume nitrogen. Weight of one vol. gas (spec. gr.) = 1.5270  
minus weight of one vol.  $\text{N}$  (spec.gr.) = 0.9713  
0.5557


but 0.5557 is very near  $\frac{1.1056}{2} = 0.5528 = \frac{1}{2}$  vol. oxygen.

Hence the gas is  $\text{NO}^{\frac{1}{2}}$  or  $\text{N}^2\text{O}$ .

*The properties of the gas  $\text{N}^2\text{O}$  are remarkable.* It was discovered by Priestly in 1776, and Sir H. Davy demonstrated its composition. The gas is colorless, possesses a faint sweetish taste and slight aromatic odor. One cubic centimeter at  $0^\circ \text{C}$ . weighs 0.001974 gram. The gas is quite soluble in cold water. At  $0^\circ \text{C}$ . one volume water absorbs 1.3 volumes of the gas, but at  $20^\circ \text{C}$ . only 0.67 volume. A taper burns in the gas more brightly than in air, almost as in oxygen gas. The gas can be taken into the lungs (breathed) without any discomfort. Quite on the contrary; its action when breathed, is that of a stimulant. The nerves become excited, the effect is similar to that of alcohol and ether—intoxication ensues, either numbness or acute mania. The effect is not by any means alike on all persons. Sir H. Davy gave it the name laughing-gas, pleasure gas. Its application in dentistry is well known. Dentists use it with nervous persons, who are unwilling to stand up against pain. Though not quite without danger, serious after-effects are less likely through its use than through other anæsthetics—ether or chloroform. The dentists buy the gas in the compressed state, in copper cylinders. At  $0^\circ \text{C}$ . the gas becomes liquid under a pressure of 30 atmospheres, that means when 30 volumes are compressed into the space of one volume, approximately; 300 litres of the gas yield 400 c.c. of liquid  $\text{N}^2\text{O}$ . When this liquid is allowed to evaporate it produces intense cold, like liquid air, and part of it becomes solid as


snow. The liquid  $\text{N}^2\text{O}$  boils at  $-89^\circ \text{C.}$ , and becomes solid at  $-100^\circ \text{C.}$  For use of the dentists the compression of the gas is only carried so far that about 10 volumes are compressed into 1 volume; which means a pressure of 150 pounds per square inch.

*Preparation of the gas on manufacturing scale.* The salt  $\text{NH}^4(\text{NO}^3)$ , ammonium niter or nitrate, breaks up into  $\text{N}^2\text{O}$  and  $\text{H}^2\text{O}$  when heated :



and thus furnishes an excellent raw material. It

FIG. 51.



is not expensive. Apparatus as shown in Fig. 51 will enable you to make the gas quickly.  $R$  is a small retort, into which has been put the ammonium nitrate at  $A$ . The stand  $S$  holds the retort.  $L$  is a

Liebig condenser with the water inflow at *i* and its outflow at *o*. The condensed water and the gas separate in the flask *C*, the gas passes into the drying tube *D* and issues dry at *G*, whence it may be conducted into any desired holder, or to a compressing pump. Substitute large iron vessels for the small glass vessels and you have the manufacturing plant.

*Recapitulation of the oxyds of nitrogen :*

$\text{N}^2\text{O}^5$  (contained in the nitrates) does not exist in free state.

$\text{N}^2\text{O}^4$  (contained in the very unstable hyponitrates) brown fumes.

$\text{N}^2\text{O}^3$  (contained in the nitrites) blue liquid in free state.

$\text{N}^2\text{O}^2$  (forms no salts) originator of brown fumes.

$\text{N}^2\text{O}$  (forms no salts) laughing-gas.

Nitrogen shows five different valences, from mono- to penta-valent, but all of them are weak. With other elements similar tendencies are observed, the smaller the affinity between two elements, the greater the number of combinations into which they enter.

## CHAPTER XI.

### AMMONIA, A VOLATILE ALKALI. A COMPLEX METALLIC RADICAL.

AMONG other facts concerning the action of KOH and NaOH, we gathered that these hydroxyds and their water solutions can dissolve zinc with the evolution of hydrogen; when the hydrogen is thus generated we say it is in the *nascent state*, meaning by this word (literally "being born") that there is a special force or energy connected with it, and which the gas has lost after it is once set free and allowed to collect or to dissipate. It is a play of the affinities. We observe that a solution of indigo is *not* decolorized by shaking it with hydrogen gas. Yet when zinc, dilute sulfuric acid and indigo are brought together, the liquid becomes colorless.

Blue indigo + Zn + H<sup>2</sup>SO<sup>4</sup> + water = white indigo + Zn(SO<sup>4</sup>) + water. Hydrogen is not evolved until all the blue indigo is changed into white indigo, and we ascribe the change to *nascent* hydrogen. Thus also when niter, NaHO, and zinc act together, the escaping gas possesses not only a peculiar odor, but the gas turns *red* litmus to *blue*. There must be with the hydrogen a new *volatile* body which possesses the properties of the *alkaline hydroxyds*. Let this, as yet suspicious, body be named *ammonia*.




Passing the gas into water, the latter soon acquires *alkaline* reaction and the power to neutralize acids. If these neutralized acids be evaporated, crystals form, characteristic for each acid, the same as if those acids had been neutralized by KOH. Yet if these crystals be heated over an open flame, they will completely disappear, totally differing from any of the metallic salts. The total volatilization is proof that the body, which here takes the place of a metal, cannot contain either zinc or sodium, and since besides these, only the elements N, O, H had been in the generating solution, they alone can constitute the new body. Hydrogen we know to be a metal, because it takes the place of a metal in the salts, producing the hydrogen salts or acids. Oxygen and hydrogen together form water, which can take the place of a metallic oxyd, but not of a metal, hence we must conclude that the new body must contain nitrogen.

*Investigation.* The first step will be the preparation of a sufficient supply of the *ammonia gas*. A word about the name. We know that the name of the Egyptian sun god was *Rha Ammon*; also that the name of a powerful tribe of Bedouins in the desert to the southeast of the Dead Sea, in Palestine, was *Beni Ammon*, in Hebrew, the sons of Ammon, as we would say the sons of the sun. It is my belief that the name *ammonia* is connected in some way with Rha Ammon, though I cannot say how. The Egyptian priests must have known this body, and for its revivifying, penetrating odor likened it

to the effect of the sun's rays. There is no historical record in existence to substantiate my view. In the Latin translation of Geber's works (ninth century A. D.) the name appears *sal armeniacum*, which would mean the salt of Armenia, but that stands also for rock salt. I believe it is a misspelling.


*Preparation of ammonia.* Put into a 500 c.c. flask (Fig. 52) 25 grams of NaOH, or about that much,

FIG. 52.




75 c.c. of water, 20 grams of zinc (in turnings), a piece of bright sheet-iron, and 5 grams of niter. Stopper the flask and connect by rubber tubing with a Will's condenser. The flask should stand on a tripod upon wire gauze. Heat gently and keep up a slow evolution of gas. The condenser is charged with 20 c.c. of water and 5 c.c. of HCl concentrated. It is best to place a tube, *T*, between flask and condenser to receive any liquid spatterings. Will's condenser is especially adapted, be-

cause either of the bulbs, *a*, *b*, can hold more than the volume of liquid indicated, hence no danger of the running back of the liquid into *T* should a vacuum occur in *F*, nor a running out at *C* should the escape of gas become tumultuous at any time. The addition of some litmus to the liquid in *W* is advisable; so long as the color remains red we know that the liquid is still able to take up more ammonia. Any hydrogen mixed with the ammonia will escape at the point, *C*, of the condenser. The action must not be overhurried. In hurrying much more hydrogen escapes, doing no work. As soon as all the air is displaced from *F*, *T*, and *W*, you will note a tendency of the liquid in *W* to advance against the gas, to pass up into the bulb, *a*, an action which indicates the strong affinity between ammonia and HCl. It will take several hours to accomplish the complete change of the niter into ammonia. Towards the end it will be well to set *T* into a glass containing boiling water to drive out the ammonia from the liquid which may have been condensed in *T*. Then empty liquid from *W* into an evaporating dish, and evaporate over a water-bath. A white, granular residue will be obtained composed of cubic crystals like those of common salt, NaCl or KCl, which is another indication that ammonia must be a body similar to Na and K. We will name this white salt ammonium chlorid = Am.Cl. In the drug trade it is named *sal ammoniac*. Bringing this salt together with Na(OH) or K(OH) or Ca(HO)<sup>2</sup> we observe at once a pungent odor of ammonia.





It follows from this action that there must be contained in the salt *one hydrogen besides the ammonia*. Important observation to be remembered. Of the three hydroxyds the one best servicable for making ammonia is  $\text{Ca(HO)}^2$ . Why? Because it is a powder. Even better is the oxyd  $\text{CaO}$ , because if an excess of it be added, then this excess will retain the water in the form of  $\text{Ca(HO)}^2$ . Even by rubbing together  $\text{Am.Cl}$  and  $\text{CaO}$ , ammonia is set free. The process follows along the equation




Pure dry ammonia gas results.

#### COMPOSITION OF AMMONIA.

Let the generating apparatus be set up as shown

FIG. 53.




in Fig. 53. In the small flask (not more than 100 c.c.) put the mixture  $M$  ( $4\text{CaO} + 1\text{AmCl}$ ) so that the

bulb is nearly full. To insure perfect dryness of the gas, let it pass through two U-tubes *U* and *U* both filled with pieces of burnt lime, pea size.

*Proof that ammonia contains hydrogen.* Rig a hard glass tube *T*,  $\frac{1}{4}$ " diameter, 12" long, as shown in Fig. 54. Between two asbestos plugs *b* and *b'* fill in coarse copper oxyd and ignite both asbestos and oxyd thoroughly, before filling the tube. Let the plugs be 3 to 4 inches apart. Arrange shield *S* so that only the filled part becomes heated, the empty

FIG. 54.



part projecting beyond the shield. Fill the tube with ammonia gas by attaching *t* to *P* in Fig. 53. Then heat *a* to redness, and let the gas pass slowly (by heating generating bulb very gently). Soon we see moisture appear in the cool tube at *W*. Meanwhile connect the outlet of *T* by means of a bent tube with the test-tube *B*, which stands inverted in basin *C* both filled with dilute  $\text{H}^2(\text{SO}^4)$ . Whatever gas collects in *B* must be nitrogen (prove by its negative actions), because any ammonia gas passing out undecomposed will become absorbed by the dilute acid. That the condensed moisture at *W* is water we prove by bringing together with it a small piece of potassium. But water could form only if the ammonia contained *hydrogen*.

*Proof that ammonia contains no oxygen.* We replace the copper oxyd at *a*, Fig. 54, by a bright copper wire, or a strip of bright copper foil and repeat the experiment at red heat. The copper does *not cover itself with a film of oxyd*. This is proof of the absence of oxygen.

Hence ammonia must be



*Demonstration that the ratio,  $\frac{m}{p} = \frac{1}{3}$ .* Let a small volume of the pure ammonia gas pass into the eudiometer *E*, Fig. 55, over perfectly dry mercury.

FIG. 55.





FIG. 56.



Dry the eudiometer, inside and outside, most thoroughly, before filling in the dry mercury. A eudiometer is a glass tube open at one end, made of strong glass, either graduated or not graduated, but having two thin platinum wires fused into the glass near the closed, or upper, end — +, so that an electric current of high potential may be sent across in form of a spark. Let *m* designate the portion of



the mercury meniscus after the ammonia gas has entered. Then turn on the current which has been transformed to high potential by means of a Rhumkorf coil. We notice at once an increase in the gas volume; rapid increase at first, but slowing down by geometric progression until the maximum enlargement of the volume has been reached at  $2m$  as shown in Fig. 56. The volume has doubled; the hydrogen and nitrogen are now merely mixed together, the energy of the shocks from the sparks having broken the chemical bond. Now let us assume that the entire volume of the gas be hydrogen, *i. e.*, 2 vols., let one vol. of pure oxygen be added, making altogether 3 vols. of gas. Let the spark pass between the wires (under proper precautions, that is covering the mercury trough with a towel and holding down eudiometer with the left hand). Had our assumption been correct there would now only be contained in tube aqueous vapor and liquid water. We must remove this water, because we started with dry gas, by bringing into the gas a ball of  $K(OH)$  fused to a thin, soft wire.  $KOH$  absorbs water eagerly. A ball of fused  $CaCl^2$  will answer also. We watch the gas from time to time and remove the drying ball when the volume remains constant for a full hour. Since we know that nitrogen will certainly be in the residue, we must have a certain and unknown excess of oxygen after the explosion, for we put oxygen equal to  $\frac{1}{2}$  of the total gas  $H + N$ . Let this excess of oxygen =  $dO$ , then the residue will be =  $N + dO$ . Suppose we

started with one vol. ammonia gas = 10 c.c. By decomposition this became = 20 c.c. By addition of one vol. of oxygen = 10 c.c. this became = 30 c.c. Now we explode the mixture and remove aqueous vapor, and find that 30 c.c. have shrunk to 7.5 c.c. Then  $30 - 7.5$  c.c. = 22.5 c.c. here disappeared in form of  $H^2O$ ;  $\frac{2}{3}$  of this contraction was hydrogen,  $\frac{1}{3}$  oxygen.

$$\frac{2 \times 22.5}{3} = 15 \text{ c.c. of H}$$

$$\frac{22.5}{3} = 7.5 \text{ c.c. of O}$$

But we had used 10 c.c. of O, hence d O =  $10 - 7.5 = 2.5$ ; hence  $N + d O = N + 2.5 = 7.5$  (gas volume after explosion);  $N = 7.5 - 2.5 = 5$  c.c.; hence ratio  $\frac{m}{p} = \frac{5}{15} = \frac{1}{3}$ . The formula of ammonia is  $NH^3$ . Into 10 c.c. of  $H^3N$  are compressed 15 c.c.  $H + 5$  c.c. N. Into 1 c.c. of  $H^3N$  are compressed 1.5 H + 0.5 N, or *two* volumes are condensed into one.

$1\frac{1}{2}$ vols. hydrogen weigh gram . . .	0.1038
$\frac{1}{2}$ vol. nitrogen weighs gram . . .	0.4856
	<hr/>
	0.5894

which is the calculated specific gravity or *volume weight* of ammonia, and agrees fairly well with the experimental specific gravity of 0.596. The *molecular weight* ( $H = 1$ ) is  $14 + 3 = 17$ .

*The chemical nature of ammonia—ammonium.* Ammonia gas is easily absorbed by water. 1 c.c. of water at  $0^\circ C$ . will absorb 1050 c.c. of the gas;


much heat is produced by the absorption. The resulting liquid, be it concentrated or dilute, has the same pungent odor as the gas. The liquid possesses the *biting taste* of potassium and sodium hydrates. Raises a blister on the tongue or lips. The liquid is lighter than water; at  $+14^{\circ}\text{C}$ . its specific gravity is 0.8844, and contains 35.0 per cent. of  $\text{NH}^3$ . This liquid goes in the drug trade by the name *aqua ammonia* or *ammonia water*. We chemists call it *ammonium hydrate*. Because from the actions of this liquid towards the acids, we conclude that as soon as the gas,  $\text{NH}^3$ , comes together with water it combines with the latter, thus



One hydrogen leaves  $\text{H}^2\text{O}$ , and attaching itself to  $\text{NH}^3$  brings forth the *metallic radical* ( $\text{NH}^4$ ). Heretofore we have only had non-metallic complex radicals, such as ( $\text{SO}^4$ ), ( $\text{NO}^3$ ). This radical, ( $\text{NH}^4$ ), we name *ammonium*. Note carefully the difference in the words ammonia and ammonium, and try not to confound them in your speech. Ammonia is the real, actual compound,  $\text{NH}^3$ ; ammonium the hypothetical metal,  $\text{NH}^4$ . I say hypothetical because we have no means at present to isolate it; in attempting to tear ( $\text{NH}^4$ ) from ( $\text{HO}$ ) we always get  $\text{NH}^3 + \text{H}^2\text{O} = \text{ammonia} + \text{water}$ . And yet there is no unproven assumption in chemistry more certain than this, that ammonium would show all the physical characters of a metal if we could separate and condense it, even as hydrogen itself would do.

*Indirect or circumstantial proof of the ammonium theory.* Let us introduce (Fig. 57) into the tube,  $T$ , over mercury, one volume of dry  $\text{HCl}$  and one volume of dry  $\text{NH}_3$ . The one gas is strongly acid, the other strongly alkaline. In order to get the volumes equal let  $t$  be a small tube holding 10 c.c. up to the mark  $m$ . If  $t$  be filled with mercury and then held so that the upper rim of the sticker falls together with the level of the mercury in the trough,  $V$ ,

FIG. 57.



$V$ , and if now the gas-delivery tube be brought under the rim and the gas allowed to enter until all the mercury is displaced, that is, until the bubbles come up on the outside, and if we do the same with the other gas, then we have measured the gases exactly at the same pressure and temperature. For the transmission of the gas from  $t$  into  $T$  we incline  $T$  as much as possible, and then with the left hand  $t$  is inclined in the opposite direction, the open end of  $t$  shoved under the open end of  $T$ ; the closed end of  $t$  is then pressed down, and the entire volume of gas

passes above the mercury into *T*. When the two gases meet a white cloud forms; the mercury rises and occupies practically the whole of *T*. Both gases have disappeared, no hydrogen has been liberated, we have



But the resulting body is a white salt crystallized in cubes and octahedrons, or a combination of the two, exactly the same as KCl and NaCl. By action of  $\text{H}^2\text{SO}^4$  upon this salt we obtain HCl, as we did from NaCl and KCl. Hence we deduce that a body must be in the salt, in every respect equal to either sodium or potassium. That is ammonium,  $\text{NH}^4$ .





Some chemists always write Am. instead of  $(\text{NH}^4)$  in order to lay stress upon the metallic nature of the group  $(\text{NH}^4)$ . Because one volume HCl combines with one volume  $\text{NH}^3$  we say ammonium is a monad, a monovalent radical; hence there is



one nitrate  $(\text{NH}^4).(\text{NO}^3)$

two sulfates  $\begin{cases} (\text{NH}^4).\text{H}.\text{(SO}^4) \text{ acid sulfate, bisulfate} \\ (\text{NH}^4)^2.\text{(SO}^4) \text{ neutral sulfate} \end{cases}$


one chlorid  $(\text{NH}^4).\text{Cl}.$

Knowing now the composition and nature of ammonia and ammonium, we can write the equation of its formation from  $\text{Na}(\text{OH})$ , Zn,  $\text{Na}(\text{NO}^3)$ , or  $\text{K}(\text{NO}^3)$  thus, in two stages,






The incentive for the action is the tendency of the zinc and sodium to form the salt  $\text{Na}^2\text{ZnO}^2$  (sodium zincate), then the hydrogen in the *nascent state* attacks the niter, substitutes itself for the oxygen and changes the latter into  $\text{H}^2\text{O}$ . A very similar action takes place when zinc is in presence of dilute  $\text{H}^2\text{SO}^4$  and  $\text{H}(\text{NO}^3)$ . Here the incentive lies in the tendency of zinc to displace the hydrogen in  $\text{H}^2\text{SO}^4$  and to form  $\text{Zn}(\text{SO}^4)$ . Then the nascent hydrogen will attack  $\text{HNO}^3$  as in the alkaline solution. Thus:



It also follows that a similar action will take place when zinc acts upon a very dilute water-solution of  $\text{H}(\text{NO}^3)$  itself, but in this case only part of the nitrogen becomes changed into  $\text{NH}^3$ .




Of 9 molecules  $\text{H}(\text{NO}^3)$  only *one* becomes changed into  $\text{NH}^3$ . The student must practice on these equations specially, because on them depends a method, and a very good one it is, to determine in an unknown substance the percentage of nitrate and nitrite. For it is evident that the action must be similar upon nitrites; one molecule of nitrite requires 2H less for its conversion into  $\text{NH}^3$ . Besides



these, which we may designate as inorganic or mineral processes, there are many other processes by which ammonia is generated, namely the processes of *fermentation* and *putrefaction*; the latter process is only a variety of fermentation and often referred to as putrid, or stinking, fermentation. 1. Ammonia forms in the curing of tobacco. The air-dry leaves being made into piles, soon begin to feel warm and give out a strong odor of ammonia. The nicotin of the leaf which contains much nitrogen is broken up by the ferment (a fungus) and the tobacco loses its rankness. The chemical reactions are complex and could not be understood by you at your present state of knowledge. 2. Ammonia forms in the fermentation of urine, because this secretion contains much urea—a highly nitrogenous substance like nicotine. 3. Ammonia forms in the distillation of soft coal; because the coal contains from one to two per cent. of nitrogen; and although one can only get—say 20 lbs. at most—of ammonium sulfate, by bringing the gases from the distillation in contact with  $\text{H}^2\text{SO}^4$ , yet so many millions of tons of coal are distilled every year, that the total of ammonium sulfate is enormous, and constitutes in fact the only *commercially important* source of supply of ammonia. The great bulk of this goes back to the field as fertilizer, to supply the growing crops with nitrogen. Much is however consumed chemically in the arts and manufactures. Though not immediately a subject for the miner or metallurgist, yet every engineer ought to be acquainted with these short statements.

Ammonia combines with many bodies, among others with silver chlorid and calcium chlorid. Note this, so that you do not attempt to dry the gas by passing through  $\text{CaCl}^2$ . On the other hand, we may utilize this fact to procure *liquid*  $\text{NH}^3$  (not ammonia water).  $\text{AgCl}$  is better suited than  $\text{CaCl}^2$ . Bring the dried, pulverized  $\text{AgCl}$  into a tube and pass over it, slowly, dry  $\text{NH}^3$  so long as the latter is absorbed. Then bend a strong glass tube  $F$  into a knee Fig. 58, and pour the  $\text{AgCl.NH}^3$  into this tube. Clean the tube at  $d$  and draw it out over the lamp, making a good strong job. The tube will be then as shown at  $a$  in Fig. 58. The  $\text{AgCl.NH}^3$  lies at

FIG. 58.




$S$ . Immerse the pointed end into snow or ice and heat  $S$  slowly. Soon a colorless, very mobile liquid will condense at  $L$ ; this is liquid  $\text{NH}^3$ . The liquefaction is, in this case, produced by the self-pressure of the gas as it becomes expelled at  $S$ . If allowed to stand the liquid will disappear because the  $\text{AgCl}$  has once again taken up  $\text{NH}^3$ . The experiment may be repeated over and over. At  $-40^\circ$  C. the gas becomes liquid without any extra pressure. At  $-90^\circ$  C. the liquid turns into a snow-

white solid, which melts at  $-75^{\circ}\text{C}$ . and has no odor, owing to the absence of tension at this low temperature. The tension of liquid  $\text{NH}^3$  at  $0^{\circ}\text{C}$ . is 7 atmospheres, and its specific gravity = 0.63 (water = 1).

In the chemical factories aqua ammonia or ammonium hydrate is prepared from  $(\text{NH}^4)^2\text{SO}^4$  by means of milk of lime in large iron cylinders *C*, Fig. 59.

$(\text{NH}^4)^2\text{SO}^4 + \text{Ca}(\text{HO})^2 + \text{water} + \text{heat} = 2\text{NH}^3 + \text{Ca}(\text{SO}^4) + 2\text{H}^2\text{O} + \text{water}$ . It is necessary to keep

FIG. 59.



the mixture in constant motion by means of a stirring wheel *W*, the shaft of which passes through well tightened stuffing boxes *B B'*. The manhole *M* serves to introduce materials; the live steam enters at *S*; the ammonia gas together with a certain quantity of water vapor passed out at *A*. *O* is the discharge opening, and the pulley *P* transmits the power to the wheel *W*. From *A* the gas passes

through a cooler in which the aqueous vapor also condenses and flows partly as gas, partly as highly concentrated liquid into glass balloons (carboys), or iron cylinders if the liquid is to be shipped into hot climates. The carboys must be most carefully packed in straw or salt hay, for if a carboy should break, the hold of a ship—for instance—could not be entered; the gas is deadly, causing immediate suffocation.

The concentrated aqua ammonia is much used in the Carré ice machine, the construction of which is explained in physics. Its principle is that a concentrated solution of ammonia gives out  $\frac{1}{3}$  its  $\text{NH}_3$  as gas, without any water, when gently heated. If the gas be taken by a suitable pump and forced into a strong tube, which is cooled by running water, so as to remove the heat of compression, and if the compressed gas be allowed to flow into another tube which is surrounded by water—stagnant—then this water will fall below its freezing-point, it will become ice. Why? Because the compressed gas in expanding into the tube (from which the air has been pumped) will absorb the heat necessary for its expansion from the surrounding objects. Theoretically no ammonia is lost, because the expanded gas is again taken hold of by the pump and compressed as before. Practically a certain portion is lost through unavoidable leakage of the apparatus, and must be replenished.

The aqua ammonia also serves as raw material for the preparation or manufacture of the other

salts, the *nitrate* (for the manufacture of laughing gas), the *chlorid* (sal ammoniac), the *carbonate* and the *bicarbonate*. The latter salts are also known as *hartshorn salt*, because they were formerly obtained by distilling hartshorn or other kinds of horn. It has been mentioned that horn and skin as well as hair are so-called *albumenoid* bodies (like albumen—white of egg) containing from 15 per cent. to 17 per cent. of nitrogen. Now if the horn be heated in a closed vessel, it will char and give off fumes as well as gases. Amongst these is the *ammonium carbonate*, and it may be condensed.

Both carbonate and bicarbonate evaporate in the air, giving a strong odor of ammonia, hence known as *volatile salt*, *smelling salts*, to be applied to fainting persons. Owing to their easy and complete volatilization, these salts are used instead of baking soda in the making of fine cake. Being incorporated into the dough (flour, milk, sugar, eggs), the salt becomes gas in the oven, raises the dough, and leaves neither taste nor smell.

$(\text{NH}^4)^2\text{CO}^3 = \text{normal or neutral carbonate.}$

$\text{H}^2\text{O} \cdot (\text{NH}^4)^2\text{CO}^3 \cdot \text{CO}^2 = \text{bicarbonate.}$

The *neutral*  $(\text{NH}^4)^2\text{CO}^3$  is a highly unstable salt. As soon as heat is applied it breaks up. The transparent or translucent crusts which we buy as neutral carbonate have normally the composition :

$2((\text{NH}^4)^2\text{CO}^3) \cdot \text{H}^2\text{CO}^3 = \text{sesquicarbonate, or } 1\frac{1}{2} \text{ carbonate.}$

The salt is made by heating together in an iron retort ;




The bicarbonate is usually a granular substance, but may easily be obtained in long prismatic, orthorhombic crystals, white or colorless. Composition as above,  $(\text{NH}^4)^2.\text{CO}^3.\text{H}^2\text{CO}^3$ .

It is prepared by passing lime-gas into a saturated solution of the sesquicarbonate. Being less soluble than the latter salt, the bicarbonate falls out in grains.

*Sal ammoniac, ammonium chlorid*,  $(\text{NH}^4).\text{Cl}$ . A white or colorless salt. Sometimes a loose, fluffy

FIG. 60.



powder composed of small cubic crystals. Mostly in lumpy crusts, dense, as if they had been melted, and exceedingly fibrous, tough; will not grind into powder. To study the formation of this body, place two watch glasses alongside of each other, pour into the one concentrated  $\text{HCl}$ , into the other concentrated aqua ammonia, and cover the two with an




inverted beaker glass or bell jar. Both liquids giving out gas, there will be at once a white cloud. The glass and table cover themselves with snowy sal ammoniac, feathered and fern-like aggregates of small crystals. At the point of sublimation or volatilization the salt breaks up into  $\text{HCl} + \text{NH}^3$ . One smells both at the same time, but so soon as the temperature drops reunion ensues, hence the salt can be sublimated in earthenware vessels, glazed, to avoid yellow stains. In Fig. 60, *a* is earthenware, *b* is earthenware, *c* is a cast-iron vessel into which the vessel, *a*, fits, *M* is the crude salt, and *S* represents the sublimated crusts of the salt.

*Use of sal ammoniac in soldering.* Soldering means the joining together of two metal surfaces by means of a film of liquid metal (solder) whose melting-point is much lower than the melting-point of the metals which are to be joined. Solder is either *hard* or *soft*. Hard solder is an alloy of 2 pts. lead and 1 pt. tin; soft solder contains one lead, one tin. The soldering-hammer or iron is a pointed piece of copper, the point of which is coated with tin. Tin will not stick to heated copper, because the latter covers itself with a film of oxyd, as you well know. But if the hot copper be rubbed together with the tin and sal ammoniac, then the two metals will cling together; the tin fairly jumps at the copper. Why? Because we have seen that the  $(\text{NH}^4).\text{Cl}$  when heated breaks up into  $\text{HCl} + \text{NH}^3$ . But  $\text{HCl}$  at once converts the oxyd film into fusible chlorids,  $\text{Cu}^2\text{Cl}^2$ ; the metals become bright and join. After

the hammer is well tinned it will hold liquid solder. And if now  $(\text{NH}^4)\text{Cl}$  in powder be strewn over the surfaces which are to be joined (brass for instance) the  $(\text{NH}^4)\text{Cl}$  will cleanse them, the solder will adhere. Liquid  $\text{HCl}$  or hydrochloric acid may be used, but is inconvenient and not so effective, because it evaporates too rapidly. Sal ammoniac is used in medicine, and also for freezing mixtures, because its solution in water absorbs much heat.

#### SOLVAY PROCESS.

The Solvay process, also known as the ammonia-soda process, has become of such great importance that we must know something about it. Nearly one-half of the world's supply of soda-ash is now manufactured by this method. Like many other processes, the reactions underlying it were known many years before the mechanical difficulties of applying them could be overcome. The reactions involved here are



The last two equations show the uses made of by-products, which help to make the process a commercial success.

The equations represent the following operations :

1. The formation of  $\text{NH}^4\text{HCO}^3$ , which immediately

reacts with  $\text{NaCl}$ , producing the most insoluble combination possible, when  $\text{CO}^2$  is passed into purified ammoniacal brine. 2. The calcination of the  $\text{NaHCO}^3$  to form the normal carbonate,  $\text{Na}^2\text{CO}^3$ . 3. The decomposition of limestone in a lime kiln to produce  $\text{CaO}$  for by-product recovery, and  $\text{CO}^2$  for operation 1. 4. The recovery of ammonia to be used over again.

The initial supply of  $\text{NH}^3$  is obtained from compounds that are the by-products of the coal-gas works. Ammonium sulphate is the principle compound used. From the equations one might think that only the initial supply of  $\text{NH}^3$  would be needed for continuous operation; but leakages have to be made up from an outside source.

The  $\text{CO}^2$  is obtained from the lime kilns and also from the calcination of the bicarbonate,  $\text{NaHCO}^3$ . The lime-kiln gases are produced in continuous kilns, and are cooled and purified before being passed to the carbonating towers. These gases contain about 35 per cent. of  $\text{CO}^2$ .

The purified brine is first saturated with  $\text{NH}^3$ , the heat produced being taken up by cooling coils in the tank. The ammoniacal brine is then pumped into towers, supplied with perforated diaphragms to make good contact between the liquid and the  $\text{CO}^2$  which is forced in under pressure.

The milky liquid drawn out from the bottom of the towers contains the  $\text{NaHCO}^3$  in suspension. This is separated and washed by means of centrifugal machines.

The bicarbonate is then calcined in covered cast-iron ovens to produce the  $\text{Na}^2\text{CO}^3$ , soda-ash. The gaseous products are cooled and sent back to be used again. The soda-ash produced by this process is more fluffy than the Leblanc soda-ash, hence it is often recalcined to increase its density. Ordinarily the Solvay soda-ash is exceptionally pure.

## CHAPTER XII.

### ORIGIN AND OCCURRENCE OF NITER.

ALL nitrates being easily soluble in water, we can expect to find any of them as mineral only under exceptional conditions; that is, in places protected against water. Such conditions exist in the rainless parts of North America and South America, Asia, and Africa. Europe has no rainless districts. But the Continent of Australia might yet prove to be a storehouse of niter; it is hardly explored as yet. Nitrates or any nitrogen compounds have never yet been found among the rock-forming minerals. Thus we are forced to the conclusion that the *air* contains the entire supply of nitrogen from which plants draw this element; store it in their structure first as *protoplasm*, then by differentiation into more complex tissues and combinations as the *gluten* of the cereal seeds, as *nicotine* in tobacco, as *morphine* in the poppies and multitudes of others. The animals feed upon the plants, building up still more complex tissues, such as muscles, nerves, skin, hair. These in their turn become metamorphosed, changed into matters rejected or excreted. Where animals congregate in numbers, as in cattle-yards, stables, etc., one notices the soil reeking with ammonia and ammonium carbonate


from the fermentation of the excreta. Before long a white film appears upon the surface (when dry weather sets in): the white film is made up of small prismatic crystals of *niter*. Hence we conclude that ammonia and ammonium carbonate become oxidized in the presence of a sufficiency of an alkaline body—become nitrates. If the niter film be scraped off from time to time, the raw material for the manufacture of niter is given in these scrapings. It will only be necessary to extract this niter earth with water, strain the liquid through cotton or linen cloth, and evaporate to the point of crystallization and let cool, when the nitrate will form in large crystals, mostly colored yellow and known as *crude niter*. By resolution and crystallization, at the last, *pure niter* results. Through this method fully one-third of the niter of commerce is still at this time made in India, notably in the valley of the Ganges in the Kingdom of Bengal. By piling up soil rich in humus into heaps or walls, 3 to 4 feet deep and 6 feet high, and covering these heaps with a shed, a so-called *niter plantation* can be built. A system of gutters or pipes distributes the liquid animal excretion—urine—over the tops of the heaps, wood ashes are strewn over from time to time (to supply the potassium), and before long the white niter film will appear on the *wind face* of the heaps. It is scraped off regularly; the extracted earth being always returned to the heap, and thus a continuous process of niter-production becomes instituted. Likewise the earth floor of stables was dug up from time to time



in all the villages of France and the rest of the continent of Europe during the Napoleonic wars, to supply the niter for the enormous consumption of gunpowder. Sweden is the only country where this custom still prevails.

However, the discovery of an extended territory in Peru, under the surface of which lies a natural deposit of *soda niter*, has gradually produced a thorough revolution in the niter industry. The former Peruvian Province, Tarapacca, now belonging to Chili, is formed by a plateau with an average alti-

FIG. 61.



tude of 3,000 feet. It forms the first steps, so to speak of the giant stairway of the Andes, the highest step of which is 26,000 feet. The surface rocks are all of volcanic origin, being both basalt, porphyry and trachyte. Between ridges of these rocks extend flat basins whose surfaces are destitute of vegetation except a few plants which have become acclimated. The region is quite *rainless*, a perfect desert. Fig.

61 is a section through a niter basin. *s, s* gives a vertical section of the different layers; *a*, top layer of ashy gray sand and pebbles, *b*, conglomerate in which the same material contained in the top layer is cemented together with clay and salt, *c*, massive niter, sometimes 8 to 9 ft. thick and perfectly white, a mixture of sodium nitrate, sodium chlorid, potassium sulfate, sodium sulfate, *sodium iodid*, sodium iodate with more or less fine sand and clay, *d*, pure sodium chlorid (common salt), *e*, clay and loam, *f*, bed rock-granite, porphyry, basalt. The natives give to the niter the name *caliche*. The mining operations consist in sinking bore holes to the surface of bed *e*, the clay. The diameter of the hole admits a workman of small stature, who widens out a small chamber and charges it with 500 to 600 lbs. of powder or a corresponding quantity of dynamite. The hole is then filled first with dry, then on top with wet, sand. The explosion usually breaks up the niter bed in a circle of about 90 to 100 feet in diameter. The niter blocks are then sorted out and transported to the leaching works some distance away. In 1873 the niter bed was estimated to underlie a territory of 550 square miles, and each square mile to contain about four million tons of niter, which would give a total of 2,200 millions of tons or an annual yield of two million tons for a thousand years. But since the present consumption is about ten million tons per year, the enormous deposit will not last over 200 years. How came this deposit to be? The nitrogen must have been

brought by the instrumentality of either plants or animals or both ; our present knowledge admits of no other source. As we find the Chincha Islands just off the coast of Peru, and upon these islands millions and millions of tons of *guano*, Indian word for the excrements of birds, it has been suggested by some that this depression or gap in the Andean Mountain chain was used in past ages as the transition point for the east-west migration of birds and they made a halting place on the shores of existing lakes (now dry desert). This theory, while accounting for the nitrogen, leaves out the phosphate contained in the guano. There are no phosphates in the niter beds ; phosphates are much less soluble than the nitrate of sodium ; they would surely be found with the niter if guano had been the source of the nitrogen. Equally fallacious is the theory of sea-weeds as original nitrogenous material. " We do not know ! " remains as answer.

#### CONVERSION OF SODA NITER INTO POTASH NITER.


Although  $\text{Na}(\text{NO}^3)$  contains more  $(\text{NO}^3)$  per unit weight than  $\text{K}(\text{NO}^3)$ ,  $\text{Na}(\text{NO}^3)$  being 85,  $\text{KNO}^3$  being 101, yet experiments proved the unsuitability of  $\text{Na}(\text{NO}^3)$  for gun powder. Thus the conversion of  $\text{Na}(\text{NO}^3)$  into  $\text{K}(\text{NO}^3)$  became necessary. It is readily done by the reaction



$\text{Na}(\text{NO}^3)$  is produced in Chili,  $\text{KCl}$  is produced (see under potassium) in the salt mines at Stassfurt, Germany. Conversion comprises the following

stages or operations: (1) Dissolve the soda niter in 1.5 parts of boiling water in a large iron vessel or open boiler, *B* (Fig. 62). (2) Dissolve KCl in three parts of water. (3) Hang into *B* a perforated sheet-iron vessel, *P*, by means of chain, *C C*, and a crane. (4) Add the KCl to the boiling  $\text{Na}(\text{NO}_3)$  solution in quantity corresponding to the ratio niter = 85, KCl = 74.5. (5) Common salt, NaCl, will begin to

FIG. 62.



fall out at once as a fine, granular body, because it is less soluble in the given water than either of the two others. (6) NaCl crystals will collect in the perforated or basket vessel, *P*, more and more as the liquid is boiled away to one-half. (7) The vessel, *P*, is lifted out and washed with boiling water, which displaces the adhering niter liquor. The washings are boiled down and added to liquid in *B*. This liquid is run into shallow iron pans, where the

impure  $K(NO^3)$  now comes out as a massive crystal crust as the liquid cools down. (8) The mother liquor, being chiefly  $NaCl$  with some  $K(NO^3)$ , is boiled down, the salt falling into basket as at first. (9) The massive, impure  $K(NO^3)$  is redissolved and boiled down until no  $NaCl$  falls, and is then run into vats, where it cools under constant agitation. The agitation causes the niter to fall out in small, loose crystals like granulated sugar. The niter meal is raked and dipped out, thrown upon drying floors, and then packed into bags and shipped. After this refining process the niter still holds about one-half per cent. of  $NaCl$ , which does not, however, interfere with its further use in powder making. The  $NaCl$  produced by this process is chiefly used for pickling meat, as the small quantity of adhering niter is desirable as a better preserving agent even than the salt.

## CHAPTER XIII.

### THE MANUFACTURE OF HYDROGEN SULFATE OR SULFURIC ACID ON A COMMERCIAL SCALE.

1. *Direct proof that  $H^2SO^4$  results by the union of  $SO^3$  with  $H^2O$ .* Sulfur disappears when heated with concentrated  $HNO^3$ . It disappears easier if  $HCl$  is also present. If the liquid be then evaporated on a water-bath a point will be reached when no further evaporation takes place. Water must be added several times and the liquid evaporated to get rid of  $HCl$  and excess of  $HNO^3$ . The residue has all the characters of hydrogen sulfate. Let the action be made with a known weight of the flowers of sulfur, say 0.5 gram. Then let us add water, after the evaporation and also 3 grams of pure lead oxyd ( $PbO$ ), and let everything be transferred to a clean porcelain crucible of known weight, and be evaporated to dryness. We know that lead vitriol will stand a low red heat without decomposition, and also that lead nitrate decomposes below red heat into  $PbO + 2NO^2 + O^2$ . If then we heat the crucible to redness until the weight be constant, we shall have the total 3 grams of lead oxyd, the total 0.5 gram of sulfur and the oxygen which was taken up by the sulfur to form the hydrogen sulfate, but



there will be no water, because  $\text{PbO}$  has displaced the water in  $\text{H}^2\text{O}.\text{SO}^3$  forming  $\text{PbO}.\text{SO}^3$ .


We find, after ignition, that the contents of the crucible weigh 4.25 grams; 0.75 gram of oxygen has been taken up by 0.5 sulfur, for  $4.25 - 3.0 - 0.5 = 0.75$ ; hence  $\frac{0.5}{3.2} : \frac{0.75}{16} = 0.01563 : 0.0463 = 1 : 3$ , hence  $\text{SO}^3$ . The reaction between sulfur and hydrogen nitrate alone is:  $\text{S} + 2\text{H}(\text{NO}^3) + \text{heat} = \text{H}^2\text{O} + \text{SO}^3 + 2\text{NO} = \text{H}^2(\text{SO}^4) + 2\text{NO}$ . (The  $\text{HCl}$  when added acts as a catalyzer possibly through the momentary formation of a chlorid of  $\text{S}$ .) Thus 32 lbs. of  $\text{S}$  will yield with 126 lbs. of  $\text{H}(\text{NO}^3)$ , 98 lbs. of  $\text{H}^2(\text{SO}^4)$ . Value of 1 lb.  $\text{S} = 0.5$  cent; of 1 lb. nitric acid, specific gravity 1.48 = 7.5 cents. But acid of 1.48 specific gravity contains only 88 per

cent. of  $\text{H}(\text{NO}^3)$ , hence 1 lb. of  $\text{HNO}^3$  costs  $\frac{7.5 \times 100}{88}$


= 8.52 cents. Thus 1 lb. of  $\text{H}^2(\text{SO}^4)$ , if made by this reaction would cost in material alone

$$\left. \begin{array}{l} \text{for S } \frac{32 \times 0.5}{98} = 0.16 \text{ cent} \\ \text{for H}(\text{NO}^3) \frac{126 \times 8.5}{98} = 10.9 \text{ cents} \end{array} \right\} = 11.06 \text{ cents.}$$



Now, the 66° Bé. sulfuric acid which contains 93 per cent.  $\text{H}^2\text{SO}^4$  is sold for 2.5 cents per lb. or nearly  $\frac{1}{4}$  of the 11.06 cents. The oxygen of the niter is too expensive for our purpose. Supposing we change  $\text{S}$  into  $\text{SO}^2$ , by burning the sulfur in air, the oxygen of which costs nothing, and then conducting the  $\text{SO}^2$  into the  $\text{H}(\text{NO}^3)$ , then we obtain



and by adding water




By this reaction the cost of materials will be cut down to  $\frac{11}{3}$  or 3.66 cents. Even this is too much. But we have seen that the gas NO becomes NO<sup>2</sup> in presence of air, and that NO<sup>2</sup> reacts upon SO<sup>2</sup> thus



Deduction: The gas NO can be made the *carrier* of the atmospheric oxygen or even more pertinently the *transfer agent*. This reaction, then, points out the road for the *cheap* manufacture of H<sup>2</sup>(SO<sup>4</sup>), for, theoretically we need only to buy the sulfur. The theoretical conditions would become realized in practice, if we could burn the sulfur in pure oxygen. But the latter requires KClO<sup>3</sup>, and turns out more expensive even than the niter-oxygen. The chemical engineer is forced to make a compromise between what is best and what is cheapest. For, since 1 volume SO<sup>2</sup> contains 1 volume of oxygen we must admit for every volume of sulfur vapor, 5 volumes of air, hence follows the mixture of 5 volumes, one of which is SO<sup>2</sup> and four are N. In addition to this, 2.5 more volumes of air must be admitted to furnish the one-half volume required for the transfer by NO so that SO<sup>3</sup> may result. Thus 6 volumes of indifferent, or useless, nitrogen must be steadily removed from the vessel in which the transferring action takes place. This

cannot be done without at the same time removing the NO (the transfer agent), while the  $\text{H}^2\text{SO}^4$  condenses to a liquid and thus removes itself. The French chemist Gay-Lussac's ingenuity intervened, however, and saved a very large percentage of NO; and thus made the low price of the sulfuric acid possible, while yet leaving a margin of profit in the manufacture. Another saving in the cost came in with the substitution of *pyrite* for sulfur. The mineral pyrite is  $\text{FeS}^2$  and contains  $2 \times 32 = 64$  sulfur for every 56 of iron, or 1 pyrite = 0.533 of sulfur, a little more than  $\frac{1}{2}$  its weight. As some sulfur always remains with the iron, we may say that pure pyrite yields 50 per cent. of sulfur in form of  $\text{SO}^2$  if properly handled. The action occurs thus:



240 pyrite + 176 oxygen = 160 iron oxyd + 256 sulfur dioxyd. If burnt in air we have an addition of 4 vols. of N = 616 nitrogen. These numbers, of course mean weight. 1 c.c. of  $\text{SO}^2$  weighs 0.00285 gram hence

$$256 \text{ grams } \text{SO}^2 = \frac{256}{0.00285} = 89824 \text{ c.c.} = 89.8 \text{ liters.}$$

1 c.c. of N weighs = 0.001256 gram, hence 616 grams

$$\text{N} = \frac{616}{0.001256} = 490.4 \text{ liters.}$$

For 1 liter of  $\text{SO}^2$  we have about 5 liters of nitro-

gen. But we must provide with this also the oxygen necessary to make  $\text{SO}^2$  into  $\text{SO}^3$ .  $\text{SO}^2$  contains  $\frac{1}{2}$  volume S and one volume O, hence 1 volume  $\text{SO}^2$  requires  $\frac{1}{2}$  volume of O or  $2\frac{1}{2}$  volumes of air, thus making a total of 1 volume  $\text{SO}^2 + 5$  volumes of N +  $2\frac{1}{2}$  volumes of air = 8.5 volumes, leaving the pyrite burners. In this mixture of gases the volume percentage of  $\text{SO}^2$  is 11.76. If this percentage falls below 4, the conversion becomes incomplete, as demonstrated by experience. As was to be expected, experience showed the necessity of an excess of oxygen over that which the calculation demands. Hence the average composition of the gas mixture, in leaving the pyrite burner, is like this: N = 81;  $\text{SO}^2 = 8.8$ ; O = 9.6 in 100 volumes. The plant (equal to what in laboratory speech we call apparatus) for the manufacture of sulfuric acid comprises the following principal parts: 1, *the burners*; 2, *the Glover tower*; 3, *the lead chamber*; 4, *the Gay-Lussac tower*; 5, *the concentrating outfit*. The diagram, Fig. 63, shows the arrangement of the plant in ground plan. *BBB* are three burners or kilns for pyrite; *D* is the dust chamber, of brickwork for the purification from solid particles of the gas; *G* is the Glover tower; *MC* is the mixing lead-chamber and *C'* is the main lead-chamber; *GL* is the Gay-Lussac tower with the chimney attachment *Ch*, from which the waste gases escape into the air. In Fig. 64 the plant appears in sectional elevation. The pyrite burner shows as a rectangular shaft or stack about 12 feet high and built of fire-brick. Several of them




are generally built close together in a row (see ground plan). Two opposite doors 1, 1 allow the iron oxyd to be drawn out, while a bell and hopper arrangement, 2, serves to introduce the pyrite, without losing any gas. At 3 is a strong cast-iron grate in the form of a cone; through the canal 4 air is admitted to this grate. Small inlets 5, 5, 5 admit air to the upper part, when necessary. The burner is started with a wood fire, until the lower furnace walls are at red heat. Then small charges of pyrite are introduced until the stack is full to within 6 feet of the top. The burning of the sulfur into  $\text{SO}^2$  furnishes all of the heat needed from this period on. A 1-foot iron pipe takes the gases from each burner into the dust chamber *D* which is built of common brick; two vertical partitions 7, 8 divide the chamber into three sections and force the gases to a broken up and down course shown by the arrows. Doors 9, 9 permit the removal of the ore-dust from time to time. A two foot iron pipe, 10, takes the gases to the *Glover tower* *G* so named after its inventor. It forms a brick stack of square section; the inside is lined with hard glazed bricks which resist the action of the acids. A perforated arch of such bricks, 11, serves to support a structure of loose, hard, stoneware bricks and cylinders 12. The idea is to present a very large surface over which the acid which comes from tank *T* above the tower, can spread and come in contact with the hot gases.

*Purpose and function of the Glover tower.* (1) To cool down the gases to the temperature needed in



the chambers. (2) To utilize the *nitrose* from the Gay-Lussac tower. The term "nitrose" was introduced to designate the combination  $\text{H}^2\text{SO}^4.\text{NO}$ . The nitrose is pumped from the tank, *T*, under the Gay-Lussac tower, by means of air pressure through lead pipes into the one-half of the tank, *T*, on top of the Glover, and the *dilute* acid formed in the chamber, *MC*, is pumped through pipe line, *13*, into the other half of *T*. From these tanks properly regulated streams pass through air-tight openings of the top of the tower upon the cylinder which forms the apex of the stoneware pyramid. Here the concentrated nitrose in mixing with the dilute chamber acid sets free the *NO*, while the now semi-concentrated acid in running over the large brick surface comes to boiling in contact with the hot burner gases, splits into concentrated acid, which collects below the arch, and is drawn thence by automatic syphon in the tank, *T*, under the tower, whence it is pumped to the top of the Gay-Lussac tower to begin a new circuit of absorbing *NO*, etc., *ad infinitum*. Aqueous vapor from the boiling acid in the meantime has mingled with the liberated *NO*, and with the other gases passes through pipe, *14*, into the mixing chamber, *MC*. Here the action sets in :



The  $\text{H}^2\text{SO}^4$  + water vapor forms a dense white fog, which becomes liquid when it strikes a cool surface ; thus the principal precipitation occurs along the sides of the chamber, whence the acid runs down

into the leaden pan, 15. Owing to the misty, foggy condition of the acid such an immense size of the chambers is required. From an average of fifteen English works, the chamber space is 21 cubic feet for *one pound* of sulfur burned in 24 hours. A plant, therefore, which is to produce five tons of 66° Bé. acid in 24 hours will require a chamber space

$$V = 10000 \times 0.94 \times 21 \times \frac{32}{98} = 64457 \text{ cubic feet.}$$

Making the chamber's cross section 20' wide by 16' high we get an area of 320 square feet, and hence  $\frac{64457}{320} = 201.4$  feet. Such a length would be best broken into 3 chambers; to wit: A mixing chamber 50 feet long, a main chamber 101.4 feet long, an end chamber 50 feet long. The cost of the lead for these three chambers will be arrived at thus:

$$2 \text{ sides } (201.4 \times 16 \text{ each}) = 6445 \text{ square feet.}$$

$$1 \text{ top } 201.4 \times 20 = 4028 \text{ square feet.}$$

$$1 \text{ bottom } 201.4 \times 22 = 4430 \text{ square feet.}$$

$$6 \text{ ends } (20 \times 16 \text{ each}) = 1920 \text{ square feet.}$$

For Gay-Lussac tower and

$$\text{connection pipes} = 2000 \text{ square feet.}$$

---

Total	18823 square feet.
-------	--------------------

Sheet-lead is rolled of many thicknesses, which are counted as so many pounds per square foot, hence 1 lb., 2 lbs., 3 lbs., . . . 12 lbs. sheet lead. Six lbs. per square foot is thick enough for the chambers, hence the total weight of lead will be  $18823 \times 6 = 112938$  pounds, and at 6 cents per lb., this represents a cost of \$6,776.28.

*The Gay-Lussac tower and its functions.* The purpose is to expose a maximum of surface covered with a moving film of concentrated  $\text{H}^2\text{SO}^4$  to the ascending gas mixture from the last chamber. When thus exposed the  $\text{H}^2\text{SO}^4$  will absorb  $\text{NO}$ . The absorption is proportional to the concentration

FIG. 65.





FIG. 66.




of the acid. Much aqueous vapor is in the gas mixture, which will be absorbed, and hence dilute the acid, lowering its capacity for  $\text{NO}$ . Thus comes the more recent practice of setting up two towers, one for drying the gases, the second for the absorption proper. A, Fig. 65, is a vertical section of the tower as commonly in use. Over a perforated arch

1 there lies a column of coke-pieces of fist size. From a shallow tray 2, studded with many small dripping tubes falls the concentrated acid over the coke, soaks into the latter and finally comes out as nitrose through the arch 1 into a tank  $T''$  under the tower. The gases enter  $I$  and leave at  $II$  into the chimney  $Ch$ . A pipe 3 passes through the leaden top (tightly) and feeds the acid from the tank  $T$  into the tray 2.

A more perfect arrangement is shown in Fig. 66, known as the Lunge-Rohrmann *plate column*. The tower is of sheet-lead in wooden or iron frame. Supporting ledges 4, 4 are provided, fused on (burned on in technical speech) along the walls or sides.

FIG. 67.



Similarly a central support 5 is provided of glazed stoneware. In the cross section Fig. 67, we see that there are four plates, each one perforated by 16 holes, and each hole enclosed within a small square area acting as a shallow reservoir. It is claimed for this system that it is quite superior in action to the coke filling, inasmuch as the surface is very large and yet regular, especially when the perforations of the alternate plates are not directly above

one another, but so that the liquid drops from the upper hole upon the dividing ridge of the lower panel.

*Concentration of the chamber acid.* Experience demands that enough water be delivered to the chamber, as steam or as spray, so that the condensing acid corresponds about to the tri-hydrate  $\text{H}^2(\text{SO}^4) \cdot 3\text{H}^2\text{O}$ . This liquid's specific gravity is  $50^\circ$  Bé. corresponding to 64 per cent.  $\text{H}^2\text{SO}^4$ . For many purposes such a strength is sufficient, for others it must be made as strong as possible. Between the

FIG. 68.





FIG. 69.



two extremes lies the strength of  $60^\circ$  Bé. = 80 per cent.  $\text{H}^2(\text{SO}^4)$ . This grade is known in the trade as *pan acid*, because it is obtained by heating the chamber acid in leaden *pans*. At a higher concentration than 60 per cent., the hot acid begins to at-

tack the lead of the pan, making white lead sulfate and  $\text{SO}^2$ . The further concentration must be carried on in retorts of either glass or platinum. Large glass retorts break easily; those of platinum are very costly. Hence a combination of the two is sometimes used in which the bottom is of platinum, the top or helmet of glass, or the helmet of lead. Figs. 68 and 69 illustrate the Gridley system of concentrating in glass retorts by continuous process. Fig. 68 shows one retort *R* in elevation, while Fig. 69 is a ground plan showing the combination of 4 retorts into a self-acting system. We see the retort set into an iron basin with a layer of coarse sand between iron and glass. Heat is applied to each retort separately by a Bunsen burner of sufficient size. Each retort has an inlet for the weak acid *1* and a syphon outlet *2* (see ground plan specially). The helmet *H* carries the weak-acid distillate into the pipe *P* of sheet-lead. As each retort of a set stands higher than its onese neighbor and lower than the other neighbor, the acid of steadily increasing strength flows from  $R_1$  to  $R_4$  and from  $R_4$  into the cooling worm *W*, which lies in a stream of running water; from the worm the acid of  $65.5^\circ$  Bé. flows into the carboy *K*. By general agreement this is the cheapest way of concentrating sulfuric acid. But only acid of 92 to 93 per cent. can be made by it.

In Figs. 70 and 71 we see an all-platinum still of the most modern construction, for making acid of 98 to 99 per cent.  $\text{H}^2\text{SO}^4$ , as furnished by the firm of Lemaire and Co. of Paris. The strenuous effort



necessary is divided between two stills *A*, *B*, each upon a separate fire-place *F*, *F'*. Each still is a flat, elliptical vessel, Fig. 71, ground plan. The still is in three detachable pieces, 1 the pan, 2 the lid, 3 the helmet and snout to carry off the vapors. (Snout not shown.) The older forms of still are circular. The elongated form is chosen because it gives a more economical use of the heat. The pan 1 is different in the two stills. In *A* the pan has 4 longi-

FIG. 70.





FIG. 71.



tudinal compartments, it consisting in fact of 2 concentric pans, the inner about  $\frac{3}{4}$  inch deeper than the outer. Thus a larger surface of evaporation is gained. Now supposing the acid of  $56^{\circ}$  Bé. being fed into *A* at the point *I*, it will flow along the outer compartment as the arrows point. The current will pass through an opening in the partition at *a* into the inner compartment and circulate to *b*, whence the syphon tube *S*, *S*, *S* will draw it into the still *B* at *G*. The partitions in *B* are transverse and do not touch the bottom, but leave  $\frac{3}{8}$  inch opening. This is necessary for cleansing the pan as from the

highest concentrated acid a small percentage of iron sulfate precipitates. But here too a meandering of the current is brought about as the arrows indicate. At *d* another syphon tube takes the concentrated acid to the cooler *C* whence it goes either to the carboys or iron tanks, because at ordinary temperatures the strongest  $\text{H}^2\text{SO}^4$  does not act upon iron. It is found that by the best care the platinum will dissolve in the strong acid at about the rate of 1 gram per ton of acid. For five tons daily production the bottom pan of the still will therefore lose  $300 \times 5 = 1500$  grams a year and can at best last 2 years.

#### MANUFACTURE OF OIL OF VITRIOL AND SULPHURIC ACID BY THE CONTACT PRINCIPLE

1. *Oil of vitriol and  $\text{SO}^3$  by Winkler's method.* Fundamental facts represented by the following equations:

(a)  $\text{H}^2\text{SO}^4$  (93 per cent. acid) + contact surface at yellow heat =  $\text{H}^2\text{O} + \text{SO}^2 + \text{O} + \text{aq.}$

(b)  $\text{H}^2\text{O} + \text{SO}^2 + \text{O} + \text{aq.} + \text{cold contact surface} = \text{Water} + (\text{SO}^2 + \text{O}), (\text{moist gases}).$

(c)  $\text{SO}^2 + \text{O} + \text{moisture} + \text{spray of concentrated acid} = \text{SO}^2 + \text{O} + \text{less conc. } \text{H}^2\text{SO}^4 (\text{dry gases}).$

(d)  $\text{SO}^2 + \text{O} (\text{dry}) + \text{platinated asbestos (at dull red heat)} = \text{SO}^3 (\text{as white vapor}).$

(e)  $\text{Spray of } \text{H}^2\text{SO}^4 (93 \text{ per cent. acid.}) + n\text{SO}^3 = \text{H}^2\text{SO}^4 + n\text{SO}^3 = \text{oil of vitriol},$

In words: 93 per cent. acid is broken up into  $\text{H}^2\text{O} + \text{SO}^2 + \text{O}$  by being spread over a highly heated surface of acid-resisting material. The resulting mixture of water vapor, sulfur dioxyd and oxygen is passed through cooling tubes in which most of the water vapor becomes liquid water. The gases are then sent over an extended surface of 93 per cent. acid (a fine spray) by which the water-vapor becomes fully absorbed, and lastly the mixture of dry  $\text{SO}^2 + \text{O}$  is passed through a cylinder heated up to dull red heat, and filled with asbestos over which a film of platinum has been spread. Here then the platinum is the carrier for the oxygen, or the *transfer agent*, as in the ordinary process the *NO* is the transfer agent.

Finally the  $\text{SO}^3$  can either be condensed in glass vessels as solid, silky, trioxyd, or it can be sent against a fine spray of 93 per cent. acid, in which the  $\text{SO}^3$  dissolves. By increasing or decreasing the volume of the spray one will be able to get *oil of vitriol* (fuming sulfuric acid) of any degree of strength. Simple as the process appears, there are considerable technical difficulties, notably the difficulty of keeping the apparatus from rapid deterioration.

2. *Making sulfuric acid by contact directly from the pyrite.* The principle which underlies this plan is the same as in the preceeding. Why can we not make  $\text{H}^2\text{SO}^4$  by passing the mixture of gases coming from the pyrite burner, namely nitrogen, sulfur dioxyd, and oxygen, directly over platinated as-

bestus? The question has been asked. At first sight there seems to be no valid objection and the experiment proves the feasibility. The cost should certainly be less, no lead chambers, no expensive stills required for the concentration. Works have been built on this plan and run for a number of years. It seems however that the conversion of  $\text{SO}^2$  into  $\text{SO}^3$  is incomplete owing to the presence of the large volume of nitrogen. Likewise the condensation of the  $\text{SO}^3$  in this diluted condition is very difficult and gives much weak acid, which has to be concentrated after all. Thus the chamber method is not likely to become superseded for some time, or until the present difficulties of the contact method have been finally overcome by ingenuity and experience.

## CHAPTER XIV.

### OTHER COMPOUNDS OF SULFUR.

We may say that sulfur possesses a strong leaning or affinity towards combination with the metals nearly equal in this respect to oxygen and chlorine. The product of the union we call sulfid.




Copper Sulfid    Copper Oxyd    Copper Chlorid.

Oxyds and sulfids are mostly insoluble in water; chlorids are mostly soluble in water. It follows that the so-called metallic ores are either sulfids or oxyds. Thus the iron ores are oxyds and sulfids:  $\text{Fe}^2\text{O}^3$ ,  $\text{Fe}^3\text{O}^4$ ; and  $\text{FeS}^2$ . The copper ores are  $\text{Cu}^2\text{S}$ ,  $\text{CuS}$ ,  $\text{CuFeS}^2$ ,  $\text{Cu}^2\text{O}$ ,  $\text{CuO}$  (exceptionally native copper,  $\text{Cu}$ ). The lead ore is  $\text{PbS}$ , zinc ores are  $\text{ZnS}$  and  $\text{ZnO}$ . Tin ores are  $\text{CuSnS}^2$  and  $\text{SnO}^2$ . The only exception among the common metals is aluminum; of it only the oxyd  $\text{Al}^2\text{O}^3$  is known but no combination with sulfur, except in the form of a *sulfate*,  $\text{Al}^2(\text{SO}^4)^3$ . This means that aluminum has no affinity for sulfur at ordinary temperature and in water solutions. In a general way we conclude that nature utilizes the affinity between sulfur and the metals to concentrate the latter within the rocks and thus make it possible for man to mine and extract them at a profit. Example: Rub together with

mortar and pestle, 1, a drop of mercury and sulfur ; 2, finely divided metallic copper and sulfur. In either case a new, black substance is formed, the sulfids of mercury and copper. Gentle heat in a closed vessel converts the black sulfid of mercury into a dark-red sulfid of the same composition and when ground fine it is called *vermilion*.


The copper sulfid  $\text{CuS}$  has a blue color ; it corresponds to the natural mineral *covellite* or *indigo copper*. Kept at a red heat in a glass tube, which is closed at one end, it changes into the bright grey sulfid  $\text{Cu}_2\text{S}$  thus :



one-half of the sulfur subliming into the cooler part of the tube.

*Hydrogen sulfid* (old name sulphuretted hydrogen).

FIG. 72.




In acting with dilute  $\text{H}^2\text{SO}^4$  or dilute  $\text{HCl}$  upon iron sulfid or zinc sulfid a very peculiar odor appears, due to a rapidly generating gas. Let A, Fig. 72, be a flask holding about 250 c.c., fitted with stopper funnel tube and evolution tube. Bring into it 20 grams of iron sulfid ( $\text{FeS}$ ) and allow the acid to fall in drops from the funnel. (5 %  $\text{H}^2\text{SO}^4$  must



be used). *B* is the wash tube with water and *C* is a U-tube filled with  $\text{CaCl}_2$  in small pea size. The gas will issue dry and pure at the narrow opening *I*. If a match be applied the gas will burn with a pale blue flame and if a dish *D* with clean undressed face and containing cold water be held into the flame, drops of colorless liquid (water) and yellow sulfur will be precipitated upon the dish *D*, whilst the pungent odor of  $\text{SO}_2$  is given off. Hence the gas must be composed of S and H.

*Proof that the compound is  $\text{H}^2\text{S}$ .* Let the gas enter the knee-tube, (Fig. 73), over mercury, until the latter's level is below the knee; mark the level with

FIG. 73.



sticker, *M*. Introduce a piece of tin, *S*, and shove it to near the end of the knee-tube. Bring *S* to red heat with lamp. Tin unites with the sulfur, forming brown  $\text{SnS}$ . After cooling we find volume of gas unchanged. We reason since 1 volume  $\text{H}^n\text{S}^m$  contains 1 volume H, then

Weight of 1 volume  $\text{H}^n\text{S}^m = 1.521$

Deduct wt. of 1 vol, H            = .089

---

1.432 = weight of S.

But 1.432 is equal to the wt. of  $\frac{1}{2}$  vol. sulfur, hence one vol.  $H^nS^m$  contains 1 vol. of  $H + \frac{1}{2}$  vol.  $S$ , or 2 vols. of  $H + 1$  vol.  $S$ —the symbol is  $H^2S$ . 100 grams of  $H^2S$  contain  $S = 94.2$ ;  $H = 5.8$  grams. Now we can write the equation of formation :



*Generation of  $H^2S$  in a steady current at the minimum of cost.* This is a very important problem, because in all analyses, both qualitative and quantitative,  $H^2S$  must be used. Different forms of apparatus have been described by inventors. Of all those the Koenig's apparatus serves the purpose best. It has already been described on pp. 61 to 63, Fig. 28, for the generation of chlorine gas in a steady, long-continued current. The apparatus is universal, may be used whenever a gas is to be produced from a solid by means of a liquid acid. Read over the description on pages 61 to 63, with the following changes: The generating tube,  $G$ , is to be charged with pieces of iron sulfid,  $FeS$ , of hazel-nut to pea size, the funnel being removed, up to the end of the funnel tube. (For chlorine we only fill the tube one-third.) The funnel bulb contains 5 per cent.  $H^2SO^4$  (never any stronger), 27 c.c. of concentrated  $H^2SO^4$  in 1000 c.c. of water. This dilute acid acts upon  $FeS$  at ordinary temperature, but more energetically at  $60^\circ C.$ , to which temperature the small flame at  $b$  heats the water in the jacket,  $J$ . Regulate the stop-cock at  $F$  so that a drop

of acid will issue per second. The gas will then flow from the goose-neck, *L*, in a strong, even current. The other product, which is a water solution of  $\text{FeSO}^4$  with a small quantity of free acid, will be discharged through the rubber tube, *R*, into the flask, *W*. Thus the acid from above will act upon a material surface always clean, always under equal conditions, and hence produce the same quantity of gas per minute.

*Note.* The pressure of the gas is equal to a column of water of the length of the funnel tube, provided that the length of the tube, *R*, from its lowest point in the bend to the discharge at *P* be equal to the length of the funnel tube or greater. If you should connect the rubber from the goose-neck to a long tube and dip the latter to the bottom of a high beaker-glass filled with water, so that the length of the water column be greater than the length of the funnel tube, then the gas will not go through the liquid, as it should do, but it will escape through the funnel or at *P*. The funnel can be supplied from a large bottle by means of a syphon and a pinch-cock or a glass stop-cock. The funnel holds sufficient acid for any ordinary operation in analysis. One often neglects to look after things at the right time, and so in this case the whole charge of iron sulfid might be used up by sheer neglect if the acid supply kept on running.

*Properties of  $\text{H}^2\text{S}$ .* Strong, unpleasant odor. Colorless. At  $11^\circ \text{C}$ . under a pressure of 15 atmospheres,  $15 \times 15 = 225$  lbs. per square inch, the gas

becomes a mobile, colorless liquid. The latter becomes a white, snow-like mass of crystals at  $-85^{\circ}\text{C}$ . The gas acts as a poison on man and animals, producing faintness, headache; if persistently breathed it causes death. A horse was placed in a large room which had been made thoroughly air-tight. One-half per cent. by volume of  $\text{H}^2\text{S}$  was mixed with the air of the room, then closed. The horse died in 15 minutes. Always generate the gas under a well-drawing hood or in the open air. One volume  $\text{H}^2\text{S}$  weighs 1.180 if the same volume of air weighs 1.000. By calculation the specific gravity is 1.1747 when air = 1, and 18 for H = 1. One c.c. of air weighs 0.001293 gram, therefore 1 c.c.  $\text{H}^2\text{S}$  will weigh  $0.001293 \times 1.180 = 0.001526$  at  $0^{\circ}\text{C}$ . and 760 mm. mercury pressure.

One volume water at  $15^{\circ}\text{C}$ . absorbs nearly three volumes  $\text{H}^2\text{S}$ . The resulting solution contains  $0.001526 \times 3 \times 100 = 0.4578$  p. c. of  $\text{H}^2\text{S}$  by weight, about *0.5 per cent.*, and is named  *$\text{H}^2\text{S}$  water*. The solution does not keep. It decomposes rapidly in the sunlight, thus  $\text{H}^2\text{S} + \text{water} + \text{O} + \text{sunlight} = \text{S} + \text{H}^2\text{O} + \text{water}$ . It will keep longer if the water has been boiled, then cooled quickly, before passing the  $\text{H}^2\text{S}$  into it, and if the bottle is then sealed air-tight before the oxygen of the air can dissolve again in the water. Brown-glass bottles are best, because the active sun rays do not penetrate much through such glass.  $\text{H}^2\text{S}$  water freshly prepared is a very handy reagent both in qualitative and quantitative work. The  $\text{H}^2\text{S}$  water shows *slight*

*acid action on litmus.* The gas itself decomposes more readily than  $\text{H}^2\text{O}$ , thus :

$\text{H}^2\text{S}$  in red-hot glass tube =  $\text{H}^2 + \text{S}$  (sulfur deposits).

$\text{H}^2\text{S} + \text{H}^2\text{SO}^4 = \text{H}^2 + \text{S}$  (sulfur floats on the acid).

$\text{H}^2\text{S} + \text{electric spark} = \text{H}^2 + \text{S}$  (Put 10 c.c. of gas into eudiometer over mercury and let the spark pass over; sulfur deposits as a snowy cloud and 10 c.c. of hydrogen are left. This is another proof of its composition of 1 vol.  $\text{H} + \frac{1}{2}$  vol.  $\text{S}$ .)


Because concentrated  $\text{H}^2\text{SO}^4$  decomposes the gas, we reason that one must not use  $\text{H}^2\text{SO}^4$  as a *dryer*, but  $\text{CaCl}^2$  instead, for this gas.

#### ACTION OF $\text{H}^2\text{S}$ UPON THE ALKALINE HYDROXYDS.

The gas is eagerly absorbed by both weak and strong solutions of  $\text{KOH}$ ,  $\text{NaOH}$ ,  $\text{NH}^4\text{OH}$ .


(a) *Action of  $\text{H}^2\text{S}$  upon  $\text{K}(\text{OH})$  in water at air temperature.*

$\text{H}^2\text{S} + \text{K}(\text{OH}) = \text{K}(\text{SH}) + \text{H}^2\text{O}$  and  $\text{H}^2\text{S} + 2\text{K}(\text{OH}) = \text{K}^2\text{S} + 2\text{H}^2\text{O}$ . Dissolve about 5 grams of  $\text{KOH}$  in 25 c.c. of water. Divide the liquid into 2 equal parts in two test-tubes. Let  $\text{H}^2\text{S}$  pass into one of the tubes until there is no further absorption, that is, until the size of the bubbles does not diminish in the ascent through the liquid, and you smell the  $\text{H}^2\text{S}$  strongly. Then the colorless liquid will contain  $\text{K}(\text{SH})$ , potassium sulfohydroxyd. Now pour the contents of the other tube into the first and you will have




The solution of potassium sulfid is likewise a colorless liquid. The properties of the two bodies are not quite alike. Both, by gentle evaporation over  $\text{H}^2\text{SO}^4$  in a dessicator (a glass vessel with a ground rim and ground cover or plate of glass), give crystals, but they do not belong to the same system.

$\text{K}(\text{SH})$  can stand red heat without decomposition if made in the following way in a glass retort, no water being present:



In this respect it acts like the hydroxyd  $\text{K}(\text{OH})$ . It is a *sulfo base*, which can form with *sulfo acids*, *sulfo salts*.

(b) *Action of  $\text{H}^2\text{S}$  upon  $\text{Na}(\text{HO})$ .*



They act like the potassium compounds.

(c) *Action of  $\text{H}^2\text{S}$  upon  $\text{NH}^4(\text{HO})$ .*



Both colorless solutions.

(d) *Action of  $\text{H}^2\text{S}$  upon  $\text{Ca}(\text{HO})^2$ .*

Make 2 grams of quicklime into milk of lime with 20 c.c. of water and pass  $\text{H}^2\text{S}$  into solution until saturated; the solution becomes clear in measure as the hydroxyd passes into the sulfohydrate, thus







The solution of the sulfohydrate decomposes during evaporation, into  $\text{CaS}$  and  $\text{H}^2\text{S}$ ; the *CaS is not soluble in water.*

$\text{Ca}(\text{SH})^2$  is used by tanners to remove the hair from the hides. Even from the living skin the hair falls out when the solution is repeatedly applied.




*Calcium monosulfid*, *CaS*, is best prepared in the dry way. Mix gypsum with charcoal powder in the proportion of gypsum 3.6 parts, charcoal 1 part, and heat the mixture in a fireclay crucible to full redness in a suitable furnace until the blue flame stops burning at the small opening left in the lid of the crucible.




The porous mass of  $\text{CaS}$  is white, when the materials are pure. It is mostly somewhat reddish-yellow. If exposed to the sunlight during the day it will be luminous during the dark of the night. It is a *phosphorescent body*. It is used for faking ghosts. When mixed with water it decomposes:



The solutions of the alkaline sulfids become yellow after some time of standing in contact with the air. This phenomenon is due to the forming of *poly-sulfid* thus:



As the action advances the color of the liquid deepens until it becomes blood-red; the capacity of the potassium for sulfur is now satisfied, ( $2n = 9$ ) but the oxydation keeps on, the solution deposits sulfur in crystals, the color becomes lighter until, at last, a cloudy solution remains with the sulfur all separated out in crystals; the liquid is  $K(OH)$  water thus:




Oxydation, therefore, means substitution of O in the place of S. *Note.*—Write out these reactions for  $Na^2S$  and  $(NH^4)^2S$ .

#### ACTION OF SULFUR ON THE ALKALINE HYDROXYDS.


Bring flour of sulfur together with a strong solution of  $KOH$  (1 : 3) into a test-tube. At ordinary temperature the action is too slow to be perceptible. On heating, the liquid assumes a yellow color, which grows in intensity at the boiling heat, while the sulfur disappears. No gas is seen to escape, hence the oxygen of the  $KOH$ , as well as the hydrogen, must enter into the new unions. The brown-red liquid acts upon copper or silver like  $K^2S + H^2O$ , that is to say, a black sulfid  $CuS$  or  $Ag^2S$  forms, while  $K^2$  unites with the  $H^2O$  as  $2KHO$ . Hence we may assume that the first action of  $2KHO$  upon  $S$  gives  $K^2S + H^2O^2$ . But the latter  $H^2O^2$  is hydrogen peroxyd, a powerful oxydizing agent, which tends to transfer one O to any oxydizable body in its neighborhood, and this is S. Now we know two oxyds of sulfur,  $SO^2$  and  $SO^3$ , either of

which might form, but in fact quite another combination results, namely,  $S^2O^2$ , and this oxyd combines thus with  $2KHO$  to form  $K^2(S^2O^3)$ . Evidently one can explain this in another way by assuming first the forming of  $SO^2$ , then of  $K^2(SO^3)$ , and then the entering of one  $S$  into this molecule in presence of an excess of  $KHO$  and of  $S$  molecules. The final reaction presents itself in the balanced equation :



$K^2S^9$  is potassium polysulfid (the maximum of saturation),  $K^2(S^2O^3)$  is *potassium thiosulfate* or *potassium hyposulfite*.


If  $NaHO$  be taken instead of  $KHO$  the action will be



$Na^2(S^2O^3) + 10H^2O$  easily crystallizes in large, transparent, colorless crystals. It is the so-called *hypo* of the photographers because its water solution dissolves the insoluble  $AgBr$  of the negative plate, but not the metallic silver which resulted from the action of the developing agent upon the exposed plate, and therefore *fixes* the image.  $Na^2S^2O^3$  also dissolves the insoluble  $Pb(SO^4)$ . It is the chief ingredient of the extracting solution in the *Russell process* for extracting the silver from its ores ; hence a most important substance.

*Technical manufacture of the salt.* Boil together  $NaHO$ ,  $S$  and water until the sulfur is all dissolved.

Then pass  $\text{SO}^2$  gas into the liquid until the yellow color has disappeared and allow the hot liquid to cool. The hyposulfite crystallizes. The proportions are: Solid  $\text{NaHO}$  96 parts, flowers of sulfur 64 parts, water 500 parts. The principle of the action is that  $\text{SO}^2$  takes from the polysulfid one S, turning into  $\text{S}^2\text{O}^2$ , and the latter decomposes  $2\text{NaHO}$  to form  $\text{Na}^2\text{S}^2\text{O}^3 + \text{H}^2\text{O}$ . Thus:




Ultimately there must be a precipitation of sulfur if an excess of  $\text{SO}^2$  is run into the solution.

*Second process.* The raw material is here the waste product from the Leblanc soda process, which is essentially a mixture of  $\text{CaS} + \text{CaO}$ . If this waste be exposed to the action of air for a proper period an oxydation will set in, by which calcium thiosulfate, calcium sulfate and sulfur are produced. If this oxydized material be extracted with water the solution contains chiefly,  $\text{Ca}(\text{S}^2\text{O}^3)$ , calcium thiosulfate. On the addition of  $\text{Na}^2\text{CO}^3$  we obtain a precipitate of  $\text{CaCO}^3$  and a liquid holding  $\text{Na}^2(\text{S}^2\text{O}^3)$ . Evaporation and crystallization furnish commercial *hypo*.


*Note.* The oxyd  $\text{SO}$  or  $\text{S}^2\text{O}^2$  has not yet been obtained in the free state. For as soon as an acid is added to a solution of *hypo* there is an action thus:  $\text{Na}^2(\text{S}^2\text{O}^3) + \text{water} + 2\text{HCl} = 2\text{NaCl} + \text{SO}^2 + \text{S}$ . This precipitation of sulfur, upon acidification, is an excellent means for recognizing the presence of a thiosulfate.

ACTION OF  $H^2S$  UPON THE SOLUTIONS OF METALLIC SALTS.

(1) *Upon solutions of potassium, sodium, ammonium and calcium salts.* We find that no action takes place.





Reasons: If there were action it would have to be thus



But we have seen that:  $K^2S + \text{water} + H^2SO^4 = K^2SO^4 + \text{water} + H^2S$ , and similarly for other sulfids; therefore the end equaling the beginning, there can be no action.

(2) *Upon solutions of iron-zinc-aluminum salts.*



There is no action for the same reason as given above. We generate  $H^2S$  by acting upon  $FeS$  with very dilute  $H^2SO^4$ , hence  $H^2S$  cannot form  $FeS$  in presence of free acid, which must be formed simultaneously with the  $FeS$ :



The same is to be said of zinc. But aluminum does not combine with the sulfur in  $H^2S$  under any circumstance, in water solution.


(3) *Action of water solution of  $K^2S$ ,  $Na^2S$ ,  $(NH^4)^2S$  upon the solutions of iron, zinc and aluminum salts.*

There is always a precipitate formed, but this is not the same thing for all the three metals; thus:




$FeS$  forms as a voluminous *black* precipitate. Why? Because there is no  $H^2SO^4$  formed, but  $K^2SO^4$ , and  $K$  is the most positive of the metals, most difficult to be displaced in its salts.

$Na^2S, (NH^4)^2S$  act the same. Write the equations.



$ZnS$  is a *white* flocculent precipitate. Reason as before.

Write equations for  $ZnSO^4 + Na^2S, (NH^4)^2S$ .



as there is no affinity between  $Al$  and  $S$ , the latter must form  $H^2S$ . Aluminum hydroxyd forms instead of the sulfid.

(4) *Action of  $H^2S$  upon the solutions of lead, copper, tin, mercury, silver and gold salts.* Here we find that precipitation of the sulfids occurs whether the solutions be neutral or whether they contain free acid.

$PbS$ , a flocculent, *brownish-black* precipitate from cold solutions, a *bluish-grey* from a hot solution.



$\text{CuS}$ , a flocculent *brownish-black* precipitate.


$\text{SnS}$ , a flocculent *brown* precipitate from *stannous* salts.

$\text{SnS}^2$ , a flocculent *yellow* precipitate from *stannic* salts.

$\text{Ag}^2\text{S}$ , a flocculent *black* precipitate.

$\text{Hg}^2\text{S}$ , a flocculent *brown-black* precipitate.

$\text{Au}^2\text{S}^3$ , a fine, granular, *brown* precipitate.



This means that the gold sulfid has very slight stability, heat causing the *metal* to precipitate. The marked difference in the action of the metal solutions towards  $\text{H}^2\text{S}$  will suggest at once the utilization for separating a mixture of salts by means of  $\text{H}^2\text{S}$ . Let a solution be made up containing the following chlorids:  $\text{SnCl}^4$ ,  $\text{CuCl}^2$ ,  $\text{HgCl}^2$ ,  $\text{AlCl}^3$ ,  $\text{FeCl}^2$ ,  $\text{ZnCl}^2$ ,  $\text{CaCl}^2$ ,  $\text{NaCl}$  and  $\text{KCl}$ . Let  $\text{HCl}$  be added and then  $\text{H}^2\text{S}$  passed through it, the liquid being warm, but not boiling. A precipitate falls, being a mixture of *SnS*<sup>2</sup>, *CuS*, *HgS*. When the liquid smells strongly of  $\text{H}^2\text{S}$  filter and call the precipitate of mixed sulfids (*A*). The filtrate is then made alkaline with  $\text{NH}^4\text{OH}$ , and another precipitate falls, of what? Of *Al(OH)*<sup>3</sup>, *FeS*, *ZnS*. Why? Because the filtrate contains  $\text{H}^2\text{S}$  in solution which becomes  $(\text{NH}^4)^2\text{S}$  when  $\text{NH}^4\text{HO}$  is added, and the  $(\text{NH}^4)^2$  acts upon the *Al*, *Fe*, *Zn* salts as soon as the free  $\text{HCl}$  is neutralized by  $(\text{NH}^4)\text{HO}$ . Pass  $\text{H}^2\text{S}$  into the liquid in order to make a *complete* precipitation, until liquid smells strongly of ammonium sulfid.

Filter and call this precipitate (*B*). Into the filtrate pass  $\text{CO}^2$  gas. Why? Because we know that calcium carbonate is insoluble in water, and this solution being alkaline with ammonia, the  $\text{CO}^2$  will form  $(\text{NH}^4)^2\text{CO}^3$  and this will give with  $\text{CaCl}^2$ ,  $\text{CaCO}^3$  plus  $(\text{NH}^4)\text{Cl}$ . Filter. Name this precipitate (*C*). In filtrate can now be only  $\text{KCl} + \text{NaCl} + \text{NH}^4\text{Cl} + (\text{NH}^4)^2\text{CO}^3 + (\text{NH}^4)^2\text{S}$ . Evaporate to dryness in porcelain dish, then heat over direct flame until the ammonium salts have smoked away. Dissolve the residue ( $\text{KCl}$ ,  $\text{NaCl}$ ) in 1 or 2 c.c. of water. We have not come across an action by which we can well separate the 2 alkali metals from one another except the difference in the solubility of the sulfates and the nitrates. Convert  $\text{KCl} + \text{NaCl}$  into  $\text{K}^2\text{SO}^4 + \text{Na}^2\text{SO}^4$ . How? By adding a few drops of  $\text{H}^2\text{SO}^4$  to the 1 or 2 c.c. of liquid; by evaporating this liquid to dryness and by then heating to red heat until no odor of  $\text{H}^2\text{SO}^4$  comes off. Because we know by experience the following actions:



Dissolve the residue in little boiling water; bring a drop upon a piece of glass, let evaporate and examine under the microscope. As the  $\text{K}^2\text{SO}^4$  is very much less soluble than the  $\text{Na}^2\text{SO}^4$ , the former will crystallize first from the margin of the drops inward. In the center will be found the long prismatic crystals of  $\text{Na}^2\text{SO}^4$ , outside the stumpy, almost cubic crystals of  $\text{K}^2\text{SO}^4$ .

If the nitrate test is to be made the first steps must be the conversion of  $KCl$ ,  $NaCl$  into  $KNO^3$ ,  $NaNO^3$ . We remember that  $AgCl$  is insoluble in water and dilute acids, also that metallic silver dissolves in  $HNO^3$  of medium concentration forming  $AgNO^3$ . If therefore we add the  $AgNO^3$  solution to the  $KCl$ ,  $NaCl$  solutions so long as the white cloud is forming, while stirring, then a change will occur thus,



Filtering off  $AgCl$ , a drop of the filtrate evaporated upon a glass slide will give the characteristic crystals of the two nitrates if both metals ( $K$ ,  $Na$ ) are present in the unknown substance, or the crystals of either alone, as the case may be. Since  $K$  salts impart to the mantle of the gas flame a *purple* color, and  $Na$  salts an *orange-yellow color*, properly applied flame test may decide the question of the presence or absence of either of the two metals, but it is not here the place to treat of these phenomena in greater detail.


Let us now turn our attention to the *precipitate* (A), and see how far our knowledge will enable us to separate and recognize the metals which it does, or may contain.

(a) *HgS*. All salts of mercury are volatile either as a whole, or at least the metal itself. If the mixture of sulfids includes  $HgS$ , we must get a sublimate of some kind by heating this mixture in a hard-glass tube, closed at one end. Three inches of the tube will suffice. (Precipitate must be dried


before placing it in the tube). Heat closed end to full redness : The sulfids melt, HgS sublimes. When tube has cooled, break off the closed end, crush glass and sulfids, act upon with  $\text{HNO}^3$ , pick out the glass and evaporate to dryness, then add water.

(b) If copper is present, a blue solution results, if tin, a white, pulverulent body of  $\text{SnO}^2$ ; if lead, a white powder of  $\text{PbSO}^4$ . (Why? Because  $3\text{PbS} + 8\text{HNO}^3 = 3\text{PbSO}^4 + 4\text{H}^2\text{O} + 8\text{NO}$ .)

Filter and wash the filter with water. Squirt the white powder into a small beaker-glass and add a few grams of  $\text{Na}^2(\text{S}^2\text{O}^3)$  sodium thiosulfate. (Hunt up this body under the action of S upon KOH, NaOH and study its formation.) Stir well 5 to 10 minutes, filter.  $\text{Na}^2\text{S}^2\text{O}^3$  acts upon  $\text{PbSO}^4$  thus :






Lead-sodium thiosulphate is soluble in water, hence if you filter, the filtrate will contain the lead salt, and if a white residue be left on the filter it must be  $\text{SnO}^2$ . Boil the filtrate and blue-grey PbS falls out :



(c) Mix the filter ashes with the problematic  $\text{SnO}^2$  together with a little  $\text{Na}^2\text{CO}^3$ . Press mixture into a cavity on a piece of charcoal, apply a good reducing flame with a blow-pipe; break out the fused mass with a knife-point, grind up in a small mortar with water. If white flakes with metallic lustre show

themselves after washing away the charcoal, then the white powder was surely tin. We have thus proved the metals in precipitate (A) with no other means but such as our progressively acquired knowledge has furnished us.

*Precipitate (B).* We have present, presumably, FeS, ZnS, Al(OH)<sup>3</sup>. It was found that all three dissolve in HCl:




By boiling solution with 1 c.c. HNO<sup>3</sup>, we shall convert FeCl<sup>2</sup> into FeCl<sup>3</sup>, thus:

$3\text{FeCl}^2 + 3\text{HCl} + \text{HNO}^3 = 3\text{FeCl}^3 + 2\text{H}^2\text{O} + \text{NO}$ , the solution turns yellow. NH<sup>4</sup>(HO) decomposes FeCl<sup>3</sup>, AlCl<sup>3</sup>, thus:




Fe(HO)<sup>3</sup>, a *flocculent, brown* substance, insoluble in water and NH<sup>4</sup>HO. But ZnCl<sup>2</sup> behaves differently towards NH<sup>4</sup>HO; it gives first a *white flocculent* precipitate, but on adding more NH<sup>4</sup>HO the precipitate dissolves.




$\text{Zn}(\text{HO})^2 + 2\text{NH}^4\text{HO} = \text{Zn}(\text{NH}^4\text{O})^2 + 2\text{H}^2\text{O}$ , soluble in water and NH<sup>4</sup>(HO). Hence it follows that we can separate Zn from Fe + Al by means of NH<sup>4</sup>HO. Add NH<sup>4</sup>HO to the hot liquid and stir until it smells strongly of NH<sup>4</sup>HO; then, filter.


Into filtrate pass  $H^2S$  and white  $ZnS$  falls. Why? Make answer yourself from what was given above. The forming of a white precipitate proves the presence of zinc. Wash the precipitate from  $NH^4HO$  from filter into a beaker glass or porcelain dish, add  $Na(HO)$ , a couple of grams, and boil.  $Fe(HO)^3$  is insoluble in  $Na(HO)$ ;  $Al(HO)^3$  is soluble in  $Na(HO)$ , because it combines with the latter to make soluble sodium aluminate.



Filter. To prove  $Al$  in the filtrate, acidify with  $HCl$  and then add  $NH^4HO$ . If a white, flocculent precipitate falls, aluminum is present. Thus:



To prove that the brown substance is iron hydroxyd, dry it on a filter. Put some of the dry mass in a cavity on charcoal and heat in a reducing (yellow) flame with blowpipe; then test the mass with a magnet. If the substance clings to magnet the presence of iron is proved. Why? Because by the reducing flame the trioxyd  $Fe^2O^3$  is changed either to *metallic iron*, or to the *tetroxyd*  $Fe^3O^4$  and *both* are magnetic. Thus:



*Precipitate (C).* The white substance thrown down by  $(NH^4)^2CO^3$  can only be  $CaCO^3$  according to our present state of knowledge. Later on we will learn of several more elements whose carbonates



will fall with the calcium. But to prove the substance absolutely, ignite it with the soda at yellow heat. Then moisten with 1 or 2 drops of water. If calcium oxyd, it will slake, *i. e.* get warm and swell up a little. If it does, add a few drops of very dilute  $\text{H}^2\text{SO}^4$  and heat, it will all dissolve; then allow one drop of liquid to evaporate on a glass slide, and examine the crystals under a microscope. They will be the characteristic needle-shaped or arrow-head shaped twins of  $\text{CaSO}^4$ .

Thus our analysis is finished. It was introduced at this point to show you that analytical chemistry simply means the thinking application of the chemical knowledge which we gradually accumulate by experiment.

#### SOME OTHER SULFUR COMPOUNDS.

*Hydrogen persulfid*,  $\text{H}^2\text{S}^2$ . Whenever you acidify a solution of ammonium, sodium, or potassium sulfid and then boil the solution, there will be at first only the odor of  $\text{H}^2\text{S}$ . But in measure as this odor becomes less pronounced, there will appear another odor, pungent and offensive, somewhat resembling the odor of fresh onions. As the eyes begin to smart if you peel and cut up an onion, so will the eyes become affected if exposed to the fumes arising from the above boiling solution. This odor is due to  $\text{H}^2\text{S}^2$ . This latter body is at ordinary temperature a yellow liquid, little soluble in water, and therefore separating from water in small oily drops. Preparation: Pour yellow  $\text{Na}^2\text{S}$  or  $(\text{NH}^4)^2\text{S}$  into an

excess of dilute  $\text{HCl}$  or  $\text{H}^2\text{SO}^4$  and let the milky liquid stand for some hours. Then pour off the liquid and you will find this yellow, evil-smelling substance at the bottom. There is no technical use for it at present.

*Sulfur and chlorine,  $\text{SCl}$ .* Place flowers of sulfur (2 grams) in a small tubulated retort and conduct dry chlorine gas into the retort. The chlorine is eagerly absorbed, generating heat, which must be removed by cold water on the outside. A red liquid forms. When the sulfur has all disappeared, close the tubulus of the retort and let the neck of the retort project into a receiver, cooled by hydrant water. Then heat the retort. A yellow-red liquid condenses in the receiver and finally sulfur remains in the retort. The original liquid was therefore a solution of sulfur in sulfur chlorid. The latter has specific gravity of 1.68. It fumes at the air because the air-moisture decomposes it:



The one property which makes this sulfur chlorid technically valuable is its solvent power for sulfur. 100  $\text{SCl}$  dissolve 73  $\text{S}$ . If soft india rubber is placed in this solution of sulfur in  $\text{SCl}$ , the rubber will absorb sulfur and become *vulcanized*. There are other combinations of  $\text{S} + \text{Cl}$  and  $\text{S} + \text{Cl} + \text{O}$ , but of no practical interest.

*Sulfur and carbon,  $\text{CS}^2$ .* Vapors of sulfur combine with carbon at a red heat forming a vaporous compound  $\text{CS}^2$ , *carbon disulfid*. On a large scale the substance is made in vertical retorts of cast-iron

lined with fire clay. The retorts are filled with small bits of charcoal and stand walled in so that they may be brought to red heat. The sulfur is introduced through an inclined tube at the bottom of the retort. The vapors of  $\text{CS}^2$  are condensed against cold water.  $\text{CS}^2$  sinks to bottom of condenser and can be drawn off. The first or raw product is far from pure and has a most offensive odor. By redistillation and shaking with metal chips as lead, copper or mercury, a colorless liquid is obtained whose odor is rather agreeable. In 1840 one pound of  $\text{CS}^2$  was worth \$5.00; in 1860 only 5 cents; to such an extent had the process been improved.

*Properties.*  $\text{CS}^2$  is a very mobile liquid. Sp. G. = 1.268. Boils at  $46^\circ \text{C}$ . High index of refraction. Very inflammable, burns with bluish-white flame. When a current of air blows through the liquid  $\text{CS}^2$ , soon snowy crystals form on the glass ( $\text{CS}^2 \cdot \text{H}^2\text{O}$ ); the temperature falls to  $-18^\circ \text{C}$ .; water dropped on the liquid freezes instantly.  $\text{CS}^2$  easily dissolves animal and vegetable fats and oils, and it dissolves sulfur. Upon these facts are based the technical applications of carbon disulfid. (1). To extract the sulfur from the sulfur earth, where the latter is abundant (near active or near extinct volcanoes). (2). To extract the fat from bones (glue factories, stock-yard slaughter-houses). (3). To extract the oil from the seeds (cotton, flax, rape, poppy, olive, nut) more thoroughly than can be done by mere pressing of the crushed seeds. (4). To remove fat from raw wool; from woolen cloth

after dyeing with certain colors. In all of these processes much care must be exercised on account of the inflammability and the poisonous effects of the carbon disulfid. It produces first headache, drowsiness, stupefaction ; if inhaled for a long time, death.

## CHAPTER XV.

### CARBON COMPOUNDS ; ORGANIC BODIES.

WE discovered in the lime gas the oxyd of a peculiar element which we named carbon from its resemblance and action to charcoal. As element we find carbon among minerals in two forms: as diamond and as graphite. Diamond is the hardest, graphite the softest of all minerals. Both resist all agents with energy. A yellow heat is required to enable oxygen to combine with adamantine carbon as well as with graphite carbon. It was a great feat to even suspect that diamond was merely crystallized carbon. Isaac Newton concluded that diamond must be a combustible substance, on account of its high index of refraction. The Academy of Florence in 1694 showed by experiment that diamond disappears in the focus of a large lens. Lavoisier showed, in 1786, that lime gas results from the burning of diamond, and Humphrey Davy, in 1807, proved it to be pure carbon. No attempt at making diamond artificially has been a success, up to this time, and it is not known by what way Nature arrives at the product. That diamond has been found in meteoric iron together with amorphous carbon (Koenig, 1889) indicates a high temperature as one of the conditions. On the other

hand the perfect crystals found in flexible sandstone in Brazil indicate a low temperature as one of the conditions.

Graphite is of very common occurrence in the archæan rocks either in mica schist or in white crystalline limestone. Sometimes large bunches together, but mostly in isolated scales. Graphite forms in pig-iron, also in gas retorts at high heat. It burns more readily than diamond, especially in oxygen gas. Long boiling with fuming  $\text{HNO}^3$  converts it into a yellow substance which has been named graphitic acid.

The *mineral coal* cannot correctly be named carbon, not even the finest anthracite of Pennsylvania; it contains however up to 95 per cent. of carbon. The great storage of carbon in the earth must be looked for in the limestones; and in the air, though the percentage of carbon dioxyd in the atmosphere be but 0.02 per cent. Yet all this immense quantity of carbon is of very little value to mankind. As  $\text{CO}^2$  it is equivalent to *spent energy*, to an uncoiled spring. At this point intervenes *organic life* contained in the animated *cell* filled with that most mysterious substance, the *protoplasm*. Under its influence carbon dioxyd is split up into carbon and oxygen, a part of the oxygen returns to the air, the remainder together with carbon and hydrogen serving in building up the body of the plants.



## THE COMBINATIONS OF CARBON AND OXYGEN.

We saw that the limestone gas is an oxyd of carbon, and furthermore that this oxyd changes into a combustible oxyd by the action of zinc and heat. There are then two oxyds of very differing properties. Being now in possession of the required knowledge we will proceed to the properties and composition of these oxyds.

*Carbon dioxyd*,  $CO^2$ , often erroneously named *carbonic acid*. This is our limestone gas as well as the gas evolved from all carbonates by the addition of HCl or any other so-called acid. This is a substance of fundamental importance, and its properties should be memorized by any engineer, because what is called perfect or complete combustion means nothing but the conversion by burning of carbon into  $CO^2$ , and because by this combustion the highest heat-value is obtained from coal or any other fuel.

*Physical properties.* Carbon dioxyd is a colorless gas without odor; but it gives to water a pleasant, prickling taste. Its specific gravity is 1.524, hence the gas is found near the floor when it issues into the workings of a tunnel, shaft, or other mine-working, or in caves such as the *dog-cave*, where a dog dies rapidly, while a standing man, alongside, breathes freely. It spec. heat is 0.22 (water = 1), 1 liter at  $0^{\circ}$  C. weighs 1.97 grams. The gas can be liquefied at the freezing-point of water under a pressure of 38.5 atmospheres; at  $31^{\circ}$  C. a pressure of 74 atmospheres is required. If two strong metallic

vessels (cylinders) be connected by a flexible metallic tube, and if one vessel has been previously charged with sodium-hydrogen carbonate and sulfuric acid in such a way that the agents only come together when the cylinder is tilted, then pure  $\text{CO}_2$  will be generated, and not being able to escape, it will liquefy in the second cylinder under its own pressure. The temperature rises to  $40^\circ \text{C}$ ., so that upon screwing a metallic receiver upon the second vessel, the liquid  $\text{CO}_2$  will distill into the receiver and the temperature will drop quickly to about  $-80^\circ \text{C}$ . and solidify into white crystal-flakes like snow. (Application as a very intense freezing mixture.) Liquid  $\text{CO}_2$  is colorless and mobile, its specific gravity at ordinary temperature being nearly that of water.  $\text{CO}_2$  is quite soluble in water at ordinary temperature, and much more so under pressure. The sparkling or boiling cold-springs, as well as the artificial soda-water are charged with the gas under pressure and release it when the pressure is taken off. If air be mixed with 4 vol. per cent. of  $\text{CO}_2$  it becomes unfit for the proper oxygenation of the blood in the lungs. In larger proportion it *suffocates* at once. Note ventilation of rooms and mines on this account. The poisonous action is negative.

*Chemical properties.* Moist litmus paper will turn to a reddish-purple in a test-tube which has been filled with the gas. We assume (without proof) that the water-solution contains a weak acid  $\text{H}_2\text{O}.\text{CO}_2 = \text{H}^2(\text{CO}_3)$ . The gas is eagerly absorbed

by  $\text{NaHO}$ ,  $\text{KHO}$ ,  $\text{Ca}(\text{HO})^2$ ,  $\text{NH}^4(\text{HO})$  and even by  $\text{Na}^2\text{CO}^3$  which changes into  $2\text{NaH}(\text{CO}^3)$ . These latter actions we utilize for the identification of the gas, for its abstraction from a mixture of gases in gas analysis. Because if the white precipitate of  $\text{Ca}(\text{CO}^3)$  be filtered off and then be acted upon by dilute  $\text{HCl}$  solution, a colorless and odorless gas will be evolved which can only be  $\text{CO}^2$ , for  $\text{SO}^2$  and  $\text{N}^2\text{O}^3$  are each pungently odoriferous.

*Composition of  $\text{CO}^2$ .* If a piece of purest charcoal be placed in a knee-tube over mercury (see proof of  $\text{SO}^2$ ), the tube be partly filled with pure oxygen, and the coal heated to redness it will burn with a bright light. When the temperature has returned to normal the volume of gas is unchanged. Hence 1 volume  $\text{CO}^2$  contains 1 volume O.

But 1 volume  $\text{CO}^2$  weighs 1.965 grams.


1 volume O weighs 1.429 grams.

---

0.536 grams.

hence 0.536 is the weight of carbon which has entered into combination. We know that the gas is  $\text{CO}^2$  for it is completely absorbed by introducing a drop of  $\text{NaHO}$  solution. Unfortunately carbon is practically non-volatile at feasible temperatures, hence we are ignorant of the weight of one volume of this element. But since  $\text{CO}^2$  and  $\text{SO}^2$  are in many ways similar and because we did prove in the case of  $\text{SO}^2$  that the difference between the weights of equal volumes of  $\text{SO}^2$  and oxygen is equal to  $\frac{1}{2}$  volume of sulfur, therefore we can assume that 0.536


equals the weight of  $\frac{1}{2}$  volume of carbon, and hence our gas contains in one volume



a contraction of 3 to 2.

On the other hand it has been found by burning a known weight of diamond, that 1 gram of the latter (pure carbon) yields 3.666 grams of  $\text{CO}^2$ , therefore in it are combined C 1.00 with O = 2,660. Having previously found that the volume ratio is  $\text{CO}^2$  we arrive now at the atomic weight of carbon, C by the proportion  $\frac{2,666}{2 \times 16} = \frac{1}{\text{C}}$  or  $\text{C} = \frac{1}{0.8331} = 12.003$ . *The atomic weight of carbon is 12.* The weight of 1 volume carbon vapor is 1,072. The molecular weight of  $\text{CO}^2 = 44$ .

*Carbon monoxyd, CO*, sometimes called *carbonous oxyd*. This gas may be generated in numerous ways of which the most ordinary is the passing of dry  $\text{CO}^2$  over well ignited charcoal at a red heat. Use the apparatus given under lime gas, substituting the charcoal for the zinc. Let the resulting gas pass through solution of  $\text{NaHO}$  which will absorb any  $\text{CO}^2$  that may have escaped decomposition. It will be seen that for every bubble of  $\text{CO}^2$  passing into the charcoal, there will be two bubbles of carbon monoxyd coming out.



This gas always generates when any fuel burns imperfectly, that is when there is a lack of oxygen.


If you observe an anthracite stove fire sometime after fresh coal has been added (opening the charging door) you will see *blue* flames bursting out all over the coal. This is very characteristic of CO—that is  $\text{CO} + \text{O} + \text{heat} = \text{CO}^2$  with blue flame. Carbon monoxyd explodes with  $\frac{1}{2}$  volume of oxygen or  $2\frac{1}{2}$  volumes of air. It can therefore be used in gas engines.

The gas is neutral, has a gravity of 0.967, is only liquefiable at  $-139^\circ \text{C}$ . and a pressure of 35.5 atmospheres. It freezes at  $-207^\circ \text{C}$ . It is little soluble in water. It is very poisonous in a positive way inasmuch as it combines with the red corpuscles of the blood, changing the color to purplish. As such it is the worst enemy to the rescuing parties who enter a coal mine after an explosion, since the combustion is usually imperfect and the gas has nearly the same gravity as air. The charcoal poisoning is due to this gas; or, any boiler-room may become deadly from it if the draft should stop on reverse; its presence cannot be told by the odor. CO is quite eagerly absorbed by solutions of *cuprous chlorid*  $\text{Cu}^2\text{Cl}^2$  either in HCl or in  $\text{NH}^4.\text{HO}$ . (Use in gas-analysis.)

*Proof that composition is CO.* Bring into a eudiometer over mercury 1 volume of the gas +  $\frac{1}{2}$  volume of O and explode. There will be left 1 volume of  $\text{CO}^2$ , it follows that 1 volume of CO contains the same volume of carbon as 1 volume of  $\text{CO}^2$ . But we saw above that 1 volume of  $\text{CO}^2$  contains 1 volume of O, hence 1 volume CO contains  $\frac{1}{2}$  volume of C +  $\frac{1}{2}$  volume of O.

## STRUCTURE OF PLANTS.

The structural unit of the plant is the *cell*. The natural shape of a plant or animal cell is the sphere, *A* (Fig. 74). When cells crowd each other in an aggregate, their shape becomes polyhedral, *C* (Fig. 74). When crowded only in one direction the cell is apt to become an elongated bag—cotton or linen fiber, *B* (Fig. 74). Each live cell has three



essential parts: The cell wall, *A2*, the nucleus, *A1*, the sap or protoplasm, *A3*, filling the cell space. From 80 to 90 per cent. of the sap is merely water. The substance of the cell wall is named *cellulose*. Since the body of the plant is made up of root, stem, branches, leaves, the flowers being merely modified leaves, and since all these are made up either



of live or dead cells, it follows that a plant, after the sap is dried out, is practically nothing but cellulose, with more or less secondary substances, such as starch, coloring matter, resin, which had been dissolved in the sap or had exuded from the cells into the intercellular space. Wood is impure cellulose. The purest form of cellulose is cotton, pith, paper pulp. Cellulose enters into so many applications, and forms as wood one of the engineer's sources of fuel, that we must devote some time and exertion to its study.

Let some cotton, or filter paper, be dried at  $105^{\circ}$  C. in an air-bath until its weight shall have become constant. We need not try whether it will burn; we know it will. But let us contrive to burn it completely in such a way that the products of the combustion can be collected and weighed. The products of the burning are flame and ashes. Flame is a mixture of gases. These gases we must collect. We do this first so that we may study the kinds (qualitative) of elements in the gases, and then, in a more guarded experiment, the quantities.

(1). We burn 1 gram in a crucible. A very small quantity of ashes remains. The ashes partly soluble in a little water, show alkaline reaction, and effervesce with  $\text{HCl}$ —( $\text{K}^2\text{CO}^3$ ). This potassium carbonate was not originally in the sap; the potassium was combined with the carbon acids, which we will consider hereafter. Combustion converts these salts into the carbonate.

(2). We burn 1 gram in an open tube and draw

(by means of an aspirator) the gases through water. The latter does not acquire acid reaction, hence sulfur and chlorine are not contained in the cellulose. While the substance burns we note the deposit of water in the cool end of the tube, therefore the cellulose contains *hydrogen*. We draw a part of the burning gases through a tube filled with lime water and note a strong white turbidity appearing showing the presence of *carbon* in the cellulose.


(3). We mix a gram of the substance with soda-lime ( $\text{CaCO}_3 + 2\text{Na}(\text{HO})$ ) and heat in a tube closed at one end—no smell of *ammonia* is noticed. Moist, red litmus held into the open end does not turn blue, hence *nitrogen* is absent, and thus we have proved that cellulose consists merely of  $\text{C}^m$ ,  $\text{H}^n$ ,  $\text{O}^p$ .

And now we proceed to find the numerical values of  $m$ ,  $n$  and  $p$ . By experiment we know that when hydrogen passes over  $\text{CuO}$  at red heat  $\text{H}^2\text{O}$  will be formed and metallic copper. Similarly  $\text{CuO}$  heated with charcoal will give metallic copper +  $\text{CO}$ . And if  $\text{CO}$  be passed at red heat over  $\text{CuO}$  it will give  $\text{Cu} + \text{CO}^2$ . Upon these facts we can now contrive a process and suitable apparatus for the purpose in hand.

In Fig. 75 the hard, infusible glass tube  $T$ , 20 inches long,  $\frac{5}{8}$  inch wide, is furnished with perforated stoppers, and is placed into the sheet-iron charcoal furnace  $F$ , being supported the entire length by a half cylinder of iron, thus preventing sagging and deformation. The tube is charged at  $1$  with 3 inches of coarse copper oxyd, then comes 10 inches

of copper oxyd with which has been intimately mixed the cellulose (0.5 gram cut up with scissors). Then comes at 3, 5 to 6 inches of coiled copper gauze, which has been partly converted into  $\text{CuO}$  by heat and oxygen. Everything must

FIG. 75.



be thoroughly dried before charging, as we desire to catch and weigh the water formed by the combustion. Into the stopper 5 fits the bulb-tube  $C$ , filled with pieces of  $\text{CaCl}^2$ . To this tube is joined by a short rubber connection the bulb-tube  $P$ , known as a Liebig *potash bulb*, and to this is joined a second smaller bulb-tube  $C'$  filled with  $\text{CaCl}^2$ . Through stopper 4 passes the inlet tube with rubber tube and spring clamp 6. The rubber tube leads to a holder for air with tubes for drying and purifying the air. The tubes are filled with  $\text{Na}(\text{HO})$  in pieces to retain  $\text{CO}^2$ , which is always in the air, and  $\text{CaCl}^2$  to take up the moisture. The bulb  $P$  is partly filled with a 1:3 solution of  $\text{NaHO}$  in water. First we weigh the tube  $C$  by itself, and  $P + C'$  together. During the process of weighing, the tubes  $C$ ,  $P$ ,  $C'$  are air-tightly closed by means of pieces of rubber and glass rod. Let the weight of  $C = a$  grams, the weight of  $P + C' =$

b grams. After all is again joined, open the clamp 6, so that bubbles will be seen to rise in the liquid of *P*; the level in the farther bulb will rise until the bubbles can pass from the middle bulb through the joining neck. The bubbles should pass at the rate of one per second. Now we place a sheet-iron diaphragm in the furnace, astride the tube *T*, just where the copper gauze begins, heap in first some ignited pieces of charcoal, then black pieces, and help the fire by gentle fanning. When the tube and gauze are red hot remove the shield or diaphragm 2 inches to the left, and even up the coal. Every 10 minutes move the shield 2 inches further to the left, until we have reached the left end of the furnace, and then keep up a uniform red heat for 10 minutes more. The cellulose is burnt up by this time, and the products of the combustion have been carried forward into the absorption tubes by the slow current of air. It may be that a small quantity of water still lingers this side of *C*. We draw *T* to the right to heat the end of it and cause the water to evaporate and to get into *C*. Now we detach first *P* from *C*, put on the glass plugs, then draw out *C* and put on the plug. Then we weigh. The weight of *C* will now be  $a'$  grams, that of  $P + C' = b'$  grams; the increase in *C* being due to water, the increase in  $P + C'$  being due to carbon dioxyd. Having started with 0.5 gram of cellulose

$a' - a$  will be  $0.2772 \text{ H}^2\text{O}$

$b' - b$  will be  $0.8147 \text{ CO}^2$

$$\frac{H^2}{H^2O} = \frac{1}{9}; \quad H^2 = \frac{H^2O}{9}; \quad \text{hence } 0.2772 \text{ } H^2O = \\ 0.0308 \text{ } H$$

$$\frac{C}{CO^2} = \frac{3}{11}; \quad C = \frac{3}{11} CO^2; \quad \text{hence } 0.8147 \text{ } CO^2 = \\ 0.2222 \text{ } C$$

$$0.5 \text{ gr. cellulose contains } 0.2222 \text{ } C; \quad 0.0308 \text{ } H; \\ 0.2470 \text{ } O.$$

For the difference  $0.5 - 0.2222 - 0.0308 = 0.2470$  must be oxygen, we have proved by the experiments above, that cellulose can contain only C, H, O. Expressed in percentage we get

$$C = 44.44$$

$$H = 6.16$$

$$O = 49.45$$

Dividing these percentages by the atomic weights we obtain the atomic quotients thus,

$$\frac{44.44}{12} = 3.7033$$

$$\frac{6.16}{1} = 6.1600$$

$$\frac{49.45}{16} = 3.0906$$


We reduce these to units of oxygen, because oxygen has the smallest quotient,

$$\frac{3.7033}{3.0906} = 1.198 \text{ or } 1.20$$

$$\frac{6.1600}{3.0906} = 1.993 \text{ or } 2.00$$

$$\frac{3.0906}{3.0906} = 1.000 \text{ or } 1.00$$

and obtain the nearest whole numbers which are 12 — 20 — 10. But these numbers are divisible by 2, therefore in the molecule of cellulose the three elements are contained in the ratio



The weight of the molecule is  $6 \times 12 + 10 \times 1 + 5 \times 16 = 162$

The first point striking our attention is that hydrogen and oxygen are in this molecule exactly in the same ratio as that of water, so that we might write the formula  $\text{C}^6(\text{H}^2\text{O})^5$ , a *hydrate of carbon*. As there are a number of other bodies of this type it is well to distinguish them as a group by the name *carbon hydrates*, although we know well that  $\text{H} + \text{O}$  are not grouped in the molecule as  $\text{H}^2\text{O}$ . For if they were, the cellulose would have to part, on heating, into  $\text{C} + \text{H}^2\text{O}$ , but it does so only in part as we shall see presently in the destructive distillation of wood.

*Some of the Properties of Cellulose or Wood Fiber.*

It is colorless or white. It is insoluble in water, in alcohol, in ether, whether cold or boiling. It is not affected by dilute solutions of the alkalies and acids at ordinary temperatures. Persistent boiling with these dilute agents slowly dissolves it.


*Vegetable parchment* = *papyrine*, results when good filter paper is immersed for a few seconds into a mixture of 1 vol. conc.  $\text{H}^2\text{SO}^4 + \frac{1}{2}$  vol. water. After withdrawing the paper from the acid it must




be washed out with much water and finally with a 2 per cent. solution of ammonium hydrate, to remove every trace of adhering acid. It is then dried. After drying, water has little effect upon it; it does not blot, and resembles dried skin.


*Nitro-cellulose, gun-cotton,  $C^6H^7(NO^2)^3O^5$ , tri-nitro-cellulose.* Let cleansed, dry cotton be immersed for 5 minutes in a liquid consisting of 1 vol. fuming  $HNO^3 + 3$  vols. conc.  $H^2SO^4$ . Let it then be squeezed out and thrown into much cold water and washed until all acid reaction has disappeared. On drying we find that externally the cotton has retained shape, color and cell structure; its weight has increased 50 per cent., its tensile strength has considerably decreased. But it has acquired the faculty to explode with great force, when struck a sharp blow with a hammer upon an anvil. Ignited by a match it burns instantly with a flash but without detonation. If the material be put into a closed space, such as a drill hole, and then ignited it will rend the enclosing material same as gunpowder.

*Explanation.* Analysis shows that the composition of the altered cotton leads to the formula  $C^6H^7N^3O^{11}$ , the substance having become a *nitro body*. Three atoms of hydrogen have been removed from the cellulose and in their places have been substituted  $3N + 6O = 3NO^2$ . We can represent this graphically thus:





The change can be represented by the equation :



There is no evolution of gas. The water is taken up by the concentrated  $\text{H}^2\text{SO}^4$  and the  $\text{HNO}^3$  suffers no loss of energy through dilution. If ordinary conc.  $\text{HNO}^3$  be taken, a different product results. It appears the same to the eye, but dissolves in a mixture of alcohol and ether. The product is essentially  $\text{C}^6\text{H}^8(\text{NO}^2)^2\text{O}^5 =$  dinitrocellulose. It bears the name *pyroxyline* and its ether-alcohol solution is called *collodion*. Collodion produces a transparent film when allowed to run over a clean glass plate. This is the film used in the *wet plate* process of photography. It is also used in surgery to keep a fresh wound from contamination with the poison microbes of the air.

*Gun-cotton (trinitrocellulose) as an explosive.* The products of the detonation of gun-cotton are  $\text{CO} + \text{N} + \text{CO}^2 + \text{H}^2\text{O}$ . Two molecules will give  $9\text{CO} + 3\text{CO}^3 + 6\text{N} + 7\text{H}^2\text{O}$ .

1 molecule  $\text{C}^6\text{H}^7(\text{NO}^2)^3\text{O}^5$  weighs  $72 + 7 + 138 + 80 = 297$ .

$2 \times 297 = 594$  grams of  $\text{C}^6\text{H}^7(\text{NO}^2)^3\text{O}^5$  give  $250\text{CO} + 132\text{CO}^2 + 84\text{N} + 126\text{H}^2\text{O}$ . 1 gram of  $\text{C}^6\text{H}^7(\text{NO}^2)^3\text{O}^5$  gives  $0.4242\text{CO} + 0.2222\text{CO}^2 + 0.1414\text{N} + 0.2121\text{H}^2\text{O} = 0.9999$  substance.

Since 1 c.c. CO weighs at  $0^{\circ}$  C. (760 mm)

$$= 0.00125 \text{ gr.}$$

$$1 \text{ c.c. CO}^2 = 0.001977 \text{ gr.}$$

$$1 \text{ c.c. N} = 0.001256 \text{ gr.}$$

$$1 \text{ c.c. H}^2\text{O (steam) at } 100^{\circ} \text{ C.} = 0.000589 \text{ gr.}$$

Therefore 0.4242 gram CO occupy the space of

$$\frac{0.4242}{0.00125} = 339 \text{ c.c.}$$

0.2222 gram  $\text{CO}^2$  occupy the space of

$$\frac{0.2222}{0.00199} = 112 \text{ c.c.}$$

0.1414 gram N occupy the space of

$$\frac{0.1414}{0.001257} = 113 \text{ c.c.}$$

0.2120 gram  $\text{H}^2\text{O}$  occupy the space of

$$\frac{0.2120}{0.0005896} = 360 \text{ c.c.}$$

But the volume V, equal to  $(339 + 112 + 113) = 564$  c.c. of  $\text{CO} + \text{CO}^2 + \text{N}$  at  $0^{\circ}$  C., will be at  $100^{\circ}$  C. equal to  $564 + (564 \times 0.37) = 772$  c.c., and the total gas at  $100^{\circ}$  C.  $= 772 + 360 = 1132$ , or over three times the volume of the gases which are evolved from 1 gram of black powder. Assuming that 1 gram of gun-cotton fills the space of 1 c.c., and that the sensible temperature of the gases after explosion be  $1000^{\circ}$  C., after proper deductions for loss, the pressure exerted by the gases will then be (for the permanent gases alone)  $772 + 285 = 1057$  atmospheres per 6 sq. centimeters. For the aqueous vapor we get, at the least, a pressure of 360 atmos-

pheres per 6 square centimeters, a total of 1417 atmospheres pressing upon 6 square centimeters. 1 square inch = 6.452 square centimeters. Hence we get  $1417 \times 15 = 21255$  lbs. pressure on one square inch nearly. When used in underground workings the air becomes poisoned, partly by the carbon monoxyd, but chiefly from the brown niter gases,  $\text{NO}^2$ . This shows that the explosion does not break up the gun-cotton always alike. NO forms instead of  $\text{CO}^2$ , and then after explosion, when the gases mix with the air, NO becomes  $\text{NO}^2$ .


THE ACTION OF HEAT UPON CELLULOSE. DESTRUCTIVE  
DISTILLATION OF WOOD. CHARCOAL MAKING.

Let a splinter of dry wood be heated over an open flame, in a narrow glass tube which has been closed at one end. We notice the wood burning first yellow, then brown, then black, while a dense yellow vapor is evolved. The vapor condenses partly into a brown-red liquid. The vapor escaping from the tube burns, is inflammable. In order to study each of these three main products, one must repeat the experiment on a larger scale, and so that the products may be completely collected, especially the gas. The following arrangement of apparatus Fig. 76 will answer and yet be simple.

A strong glass tube *T* closed at one end lies in a furnace *F* as in the previous experiment. The tube holds a stick of dry wood 20 grs. in weight. The tube *T* reaches into a glass receiver *R*, home-made from a large test-tube. A rubber band *2* makes this

connection tight. *B* is an open bell jar with stop-cock *6* connected by angle tube *5* and rubber tube *4* with the receiver. The bell stands in a large beaker glass *H*, two-thirds full of water. With open stop-cock *6* and sucking at the rubber tube *4* the bell is filled with water; the stop-cock *6* is then closed. Live charcoal is piled into the furnace to the left of the shield *S*. The distillation begins; the gases drive the air out of *T* and *R*. As soon as the yellow cloud fills *R*, we connect the rubber tube *4* with *R*, and by so doing will have fair assurance that

FIG. 76.



whatever gases are now collecting in *B* will have been produced from the breaking up of the cellulose molecule. Care is taken to give *T* a slight inclination forward, to wit, by raising the closed end. The condensed liquid will then run into *R*, and not backward, where it might cause a breaking of red-hot glass. When the flow of gas becomes slower we move *S* into the position 7, and fill in fresh live coal. Be watchful of the glass tube *T*. If its temperature rises above cherry redness it is apt to bulge out, become weak and perforated, an event which prema-

turely ends the experiment. We maintain the heat until the volume of gas in *B* becomes stationary. All this time we have been lifting the bell *B* so that the liquid *inside* remained higher than the outside level, which means a steady suction or partial vacuum. The latter is an additional precaution against the distention or bulging of the softened, red-hot glass tube. If now a mark is made upon the bell to indicate the height of the level when water stands evenly inside and outside, (the coal being removed and the tube having acquired ordinary temperature) we will have the volume of gas *V* which can be obtained from the distillation of 20 grams of dry wood. We then close the stop-cock *C*, and set aside the bell and holder. We detach the receiver and weigh it with contents, then without contents, and thus find the weight of the latter. Lastly we take out the charcoal from the tube and find its weight.

(1) *The charcoal.* Varies in color from red-brown to deep black, according to the temperature. The higher the heat, the deeper the black. The brown charcoal is imperfectly done, as shown by the following analysis: C, 70.4; H, 4.6; O, 24.2; ashes, 0.85.

But even the hardest and blackest charcoal is not all carbon; temp.  $1300^{\circ}$  C.: C, 90.8; H, 1.6; O, 6.5; ashes, 1.15.


(2) *The brown liquid.* Even in the receiver, *R*, one notices that the liquid consists of two parts; one watery, *mobile*, brown liquid, we designate *pyro-*



*ligneous acid*; the other portion, dark black-brown, viscous, not very liquid, shall be named *wood-tar*. Both portions possess a penetrating, peculiar, not unpleasant odor—the odor of smouldering wood, *empyreumatic odor* from Greek, fire odor. The taste of both portions is bitter, unpleasant. Some of the tar collects over the pyroligneous acid, some at the bottom, and some remains suspended. This proves tar to be a mixture of several bodies. The pyroligneous acid is so named because of its acid reaction on litmus, and from *pyro*, fire; *lignum*, wood, *firewood acid*. Let the two portions be separated by filtering through charcoal powder which lies in a cotton-plugged funnel. The tar particles adhere to the charcoal, pyroligneous acid passing through. We add solution of NaOH until the acid reaction on litmus ceases. We bring the liquid into a flask with perforated stopper, a glass tube, bent over so as to connect with a cooling tube, passing through the hole. On boiling, a colorless liquid passes over which shows the strong odor still very markedly. But the liquid is inflammable; its specific gravity is less than water. This substance is known as *wood spirits*. A closer study reveals that this liquid may be separated by fractionating into water plus concentrated spirits, and the latter again fractionated gives the pure wood spirits, also known as *wood alcohol*, *methyl alcohol*. It is a very mobile liquid, which boils at  $60.5^{\circ}\text{C}$ . Its specific gravity at  $4^{\circ}\text{C}$ . = 0.81 (water = 1). Mixes with water in all proportions. Burns with bright bluish hot

flame. It is very useful in the chemistry of dye-stuffs as a solvent and modifying agent.

The analysis gives for it the symbol  $\text{CH}^4\text{O}$ . We evaporate the remaining liquid in a boiling flask to dryness. A black mass results which we calcine in a porcelain dish, that is heated over the open flame until the black color has become grey-oily; gases and vapors pass off. We return it then to the distilling flask with  $\text{H}^2\text{SO}^4$  and water. A colorless liquid condenses, very mobile, very sour. Because, we reason, if the original acid in the pyroligneous liquid be  $\text{HA}$  then by adding  $\text{NaHO}$  we get  $\text{NaA} + \text{H}^2\text{O}$ . Then acting with  $\text{H}^2\text{SO}^4$  must give us




The analysis gives for  $\text{HA}$  the formula  $\text{H.C}^2\text{H}^3\text{O}^2$ . We name it *acetic acid* from *acetum* = Latin for vinegar, for the acid resulting from the fermentation of fruit juices has the same composition and properties as this wood acid. Acetic acid  $\text{H.C}^2\text{O}^3\text{O}^2$ —short symbol,  $\text{HA}$ —is a liquid, colorless mobile; with very sharp odor. At  $0^\circ \text{C}$ . this liquid acid becomes solid, crystallizes; and strangely these crystals only melt at  $16^\circ \text{C}$ ., having sp. gr. 1.0553. This most concentrated acid is known as *glacial acetic acid*. When water is mixed with this glacial acid, the specific gravity *increases* although the water is less heavy. This can only be explained by assuming that the molecules of the strong acid are in their free state farther apart than in the water-mixed state, that the addition of water causes the drawing together of these molecules. The spe-

cific gravity increases until 22 parts of water have been added to 78 pts. of acid, when the specific gravity = 1.0748. Beyond, with more water the specific gravity decreases. A mixture of acid 43, water 57 has again the same specific gravity as the pure acid. HA forms salts with all metals except gold; and the normal salts are all *easily soluble in water*. The *acetates* are much used in dyeing, especially  $\text{Pb(A)}^2$  and  $\text{Al(A)}^3$ . In the manufacture of charcoal, the pyroligneous acid is neutralized with  $\text{Ca(HO)}^2$  slaked lime, in place of  $\text{Na(HO)}$  because cheaper. Hence the crude  $\text{Ca(A)}^2$  is the product of commerce, shipped from the woods to the large cities, where the acid itself is then distilled.

(3) *The tar or wood tar.* It has been stated above that by this name is meant the oily dark-colored material which either rises as a scum to the top of the wood acid, or sinks to the bottom of the latter. We separate it from the watery liquid and proceed to its examination. The expression Tar-jacket for a sailor came in use through someone's observing that the steeping of wood or linen in tar, made those substances impervious to water, and prevented both dry-rot and wet-rot (the decaying of wood). Hence the practice rose to soak the wooden planks of boats and ships with the tar, also the entire rigging, especially halyards and shrouds and cables, also the overalls of the sailors, with this strong-smelling material. Later on fence posts, telegraph poles, etc., were steeped, and lastly, railroad ties.

Evidently mine timber would be benefited by means of repeated coats of tar, wherever mines are half wet, and the timbers succumb rapidly to the rot, *i. e.*, fungus and microbes. The curing of hams, bacon, fish, by first salting and then drying them out over a smoking, smouldering wood fire, is a very ancient practice. The tar oils condense on the surface of the meat and keep off the microbes of putrefaction until the meat is dried out and no longer

FIG. 77.





subject to the action of these organisms. Tar burns freely and is used as a so-called liquid fuel. Let the tar be heated in a retort, which is furnished with a thermometer, Fig. 77, *t*; the neck of the retort reaching by a reducer *B*, the tube of a Liebig cooler *L*, the joint *R-N* made tight by a rubber band. Cold water enters from a hydrant through

connecting rubber 1 into the cooler, the warm water leaving through rubber tube 2. The flame from burner *F* is so regulated that the liquid keeps gently boiling. We notice that the mercury in thermometer will be steadily rising, while a clear liquid collects in the graduate *G*. The rising thermometer can mean two things: (a) The tar is a mixture of liquids having their boiling points at different temperatures. (b) The tar is a uniform chemical substance, which, however, breaks up by the heat into bodies of different boiling points. Evidently it makes no difference for practical purposes whether the truth lies in condition (a) or whether it lies in condition (b). If the graduate *G* be emptied when 10 c.c. have been condensed, and so every other 10 c.c. be collected separately (distillation by fractions), it will be found that the specific gravity of each fraction is higher than that of the preceding fraction, varying from 0.82 to 0.87. Before the Pennsylvania coal oil had come into general use, these tar oils took the place as illuminating agents, the lighter part going by the name of *photogen*, while the heavier part was known as *solar oil*. To the latter clung the strong penetrating odor. If this portion be shaken with NaOH or KOH in water-solution for some time, and the two liquids be allowed to stand quietly, they will separate, the oil above, the KOH water-solution below. In a separatory funnel the aqueous liquid can be drawn off clean. We make acid with dilute  $\text{H}^2\text{SO}^4$ , and find again an oil separating at the top. This oil possesses a penetrat-



ing odor, and a bitter astringent taste. It used to be called *creosote*, is now called *carbolic acid* or *phenyl hydroxyd*. The combustion analysis (refer to cellulose) gives the ratio of the elements as  $C^5H^6O$ , but because this body as we have seen combines with  $KOH$ , and because it also dissolves in strong  $H^2SO^4$ , we write its formula,  $C^6H^5(HO)$ .  $C^6H^5$  as radical we call *phenol* or *phenyl*. Towards strong bases it acts as an acid, towards strong acids as a base. Thus:




Carbolic acid or *creosote* is at ordinary temperature a crystallized solid without color, but possessing a strong smell. It melts quickly into a thick oil. One part dissolves in 10 parts of water. It is very poisonous to animal life. It is used as a germ-destroyer, but its chief use is to act as base for the manufacture of *picric acid*. Coal-tar contains more carbolic acid than wood-tar.

*Trinitrophenol*,  $C^6H^2(NO^2)^3OH$ , *picric acid*. Take about 1 c.c. of liquid carbolic acid. Pour this into 10 c.c. of fuming  $HNO^3$ . The action is apt to be violent, copious brown fumes being formed. Boil until brown fumes stop. Then pour liquid into much cold water. The water takes a deep yellow color and a yellow sediment falls out. The yellow substance is picric acid. Pour off the yellow



liquid and add to the residue again about 50 c.c. of water, and boil. The picric acid being more soluble in boiling water, a brown resin-like substance remains. As the boiling yellow liquid cools down it throws out yellow, scaly crystals, the picric acid; *picros* = bitter, on account of the strong bitter taste. Solution reddens litmus; can be neutralized with KOH or  $\text{NH}^4\text{HO}$  when yellow needles fall from the solution, representing the potassium, or ammonium picrate. Both these salts are very explosive, more so than the acid itself. The ammonium picrate is used in the smokeless powders.



The solution of picric acid is used in dyeing silk and wool a beautiful golden-yellow color.

*Paraffin*,  $\text{C}^{17}\text{H}^{36}$  to  $\text{C}^{24}\text{H}^{50}$ . On continuing the distillation of the tar, after the solar oil sp. gr. 0.86, we find that the vapors condense in the tube to a semi-liquid, buttery product, and finally they condense to a rigid solid. By pressing the semi-liquid between paper, we find scaly crystals in the paper and a liquid pressed through the latter. The liquid is known as *paraffin oil* for lubrication, and the crystals go under the name of *paraffin*. So also does the material which solidifies at once into a rigid solid. After refining, the paraffin is used for *imitation wax* candles. As given above, the solid is not a homogeneous body, but a succession of hydrocarbons of the general formula  $\text{C}^n\text{H}^{2n+2}$ . The melting-point is


the arithmetical mean of those of the different members.

Paraffin is useful in the laboratory to soak corks in, to paint over labels with, and in fact to cover all metallic apparatus which is not subjected to heat, for it is a body very indifferent to chemical action.

Only hot concentrated  $\text{HNO}^3$  or hot  $\text{H}^2\text{SO}^4$  will decompose it.

*Examination of the wood gases.* In this examination we make use of the solubilities of the gases in

FIG. 78.




liquids or their absorption by solids. The sum of the processes we designate as *gas analysis*. The most convenient apparatus for the purpose has been designed by Professors Winkler and Hempel; though a number of others of no less ingenuity have con-

tributed their share. In Fig. 78 *B* represents the *burette* or measuring vessel, a cylindrical glass tube graduated into 100 c.c. and  $\frac{1}{5}$  c.c. The tube stands in the cast-iron foot *F* so that it will not be upset easily. *F* is bored out at 1 to admit a small tubulature to which a strong rubber tube is wired. The stop-cock *S* has a capillary perforation which leads to the capillary tube 3. With the latter is united by the wired rubber 5 the *capillary* transfer tube 4 in form of a double L. The hollow volumes or canals of both tubes do not exceed 0.01 c.c. *L*, Fig. 78, is the *level tube* in foot *F* to which the rubber tube from foot of *B* is joined and wired at 6. The rubber tube should be as long as the cylinder is high, so that the foot of one cylinder may be raised to the top of the other cylinder. This arrangement permits the reading of the gas volume under constant pressure, *i. e.*, the pressure of the atmosphere, and in so much as the temperature of the room does not materially change during one set of absorptions, we may set down the temperature as constant and thus avoid the reductions of volumes to 0° C. and 760 mm.

Fig. 79 shows the pipette or absorption vessel for absorbing CO<sup>2</sup>. The glass bulb, *C*, is seen filled with small coils of iron wire gauze, which are introduced by means of the wide mouth and stopper at *S*. This bulb communicates through tube, 1, with the reservoir-bulb, *D*, which is open at the top. A capillary glass tube leads in single loop to the rubber tube, 3, the latter being closed by the pinch-cock, 4. The whole combination is fastened to the

iron frame, *F*, which has taken the place of the former wooden frame, the spilling of the strong agents over the wood making it soon unsightly and insecure to stand up. The absorbing liquid is a solution of 15 grams of KOH or NaOH in 50 c.c. of water. This solution is poured into *D* (pinch-cock being open); the air is forced into *D*, the pressure driving the liquid gradually through the capillary

FIG. 79.




tube leading to 3, until the liquid appears at the top of 3. The coils of iron wire gauze being moistened with the alkali, expose a very large surface to the gas and cause an almost instantaneous absorption of the  $\text{CO}^2$ .

Fig. 80 represents a *double pipette*, to be used for liquids which are easily deteriorated by the oxygen of the air, like potassium pyrogallate for oxygen, and acid or ammoniacal solution of  $\text{Cu}^2\text{Cl}^2$  for CO. The capillary is provided with a pinch-cock, as in previous form. Being filled as the

figure indicates, the bulb, *A*, being the absorption bulb, is quite full, whilst the liquid rises in *B* only to the lower outlet. *C* contains the same liquid, but only partially filled. Hence the air has no access to the liquid in *A* and *B*, and can spoil only the liquid in *C* and *D*. When the gas enters *A* through the capillary, it pushes the absorption

FIG. 80.




fluid into *B*, which is large enough to hold the entire volume of *A*. As the fluid rises in *B*, the nitrogen forces down the liquid in *C* until the bubbles can pass into *D*, and when, after the absorption, the gas leaves *A* returning to the burette, air will bubble from *D* into *C*. But it loses its oxygen to the liquid, and thus there is only nitrogen between the levels in *B* and *C*. In others words, *C* and *D* are simply *traps*.

The pipette, Fig. 81, serves when the absorption liquid attacks metal and yet a large wet surface is to be exposed to the gas; using, for instance, oil of vitriol

or bromine water. Here the bulb *B* is filled with glass beads. These were put in by the glass blower before he narrowed the neck which unites *B* to *A*. Through the glass blower's dexterity we are thus in possession of an elegant set of apparatus. But suppose one of the pipettes meets with disaster just when

FIG. 81.




you need the set most. What then? A week would pass before you get a duplicate from the dealer. Fig. 82 shows how an equivalent may be made quickly by means of 2 Erlenmeyer flasks, *E*, *E'*. The flasks should hold 200 c.c. each. The flasks have doubly-perforated stoppers: Through one of the holes passes the siphon tube 1,  $\frac{1}{8}$ " inside diameter, reaching to near the bottom of both flasks. Into the second hole of stopper *E* is stuck a piece of thermometer tube—there are always broken thermometers about a laboratory. The thermometer has a capillary canal, just what we need. The piece



is about  $2\frac{1}{2}$  inches long; the sharp edges are rounded over a flame; the tube is stuck in so that the lower edge is flush with the stopper, avoiding any spaces for the lodging of gas. A piece of rubber tubing (black) 2" long is wired over the upper end and a pinch-cock clamped just above the end of the capillary tube. Through the second hole of stopper *E'* passes a short glass tube 3,  $\frac{1}{8}$ " inside, to which a piece of rubber tubing 4 is attached (for convenience in blowing air). *E* may be filled with coils of iron wire gauze, or with glass beads to give absorb-


FIG. 82.



ing surface. In filling with absorption-liquid remove stoppers, pour into each flask so that the levels are even and nearly up to  $\frac{1}{2}$  vol. of flask. Then insert stoppers very tightly, open pinch-cock and blow into rubber tube 4 until the liquid appears above the pinch-cock; then close the latter and *E* will remain full, ready to receive the gas. The two flasks form one unit. To prevent deterioration of stoppers, they should be coated with

molten paraffin and after pressing them into the necks, copper wire should be stretched across the tops to prevent their slipping out, as the paraffin is a splendid lubricator. Such an apparatus gives perfect results. If the measuring cylinder, the burette, should break, you can construct a very satisfactory substitute as follows: Make a square block of wood 4" side length and 3" thick. Bore an inch hole clear through the center and a slot *S*, Fig. 84, clear through the side,  $\frac{1}{4}$ " wide. Then nail

FIGS. 83, 84, 85.



a square of heavy sheet lead ( $\frac{1}{4}$ " thick) across the bottom as at *L*, Fig. 83. The top edges of the block should be bevelled. This will give you an excellent stand, or foot for the burette. Select a piece of glass tube as nearly cylindrical as possible of from  $\frac{5}{8}$ " to  $\frac{1}{4}$ " inside dia. about three feet long. Scrub the inside of the tube with warm NaOH solution to

remove any fat; then rinse and dry. Round off the sharp edges of one end, fit a cork or rubber stopper with one hole, into which you have put a  $\frac{1}{4}$ " glass tube, bent as shown at *t*, Fig. 83. Close *t* with a bit of wax plug; then set it into the foot, after first sticking some cotton into the hole. Fill with water to about 1 inch above the wood and scratch with a file a line *m* at the upper rim of the meniscus. Weigh into a beaker glass 100 grams of water not nearer than 10 mgr. which is equal to 0.01 c.c. Pour this water into the tube. But remember that the beaker as well as the tube should be moistened before either weighing or pouring, for some water will adhere to the glass wall, thus causing an error.

Now make another scratch at the upper rim of the meniscus *m'*. Now empty the tube and cut off the latter 1" above *m'* at *m''*, round off the sharp edges over the flame and fit a stopper into which you have stuck 2" of thermometer tube, the edge of the latter being flush with the under side of the stopper. Fasten a strip of white writing paper (several lengths may be stuck together) and lay off upon it the distance *m-m'* accurately. Divide this space into 100 equal parts, and each part into fifths. Number every 10 c.c. and draw out in india ink, cut it into a strip  $\frac{1}{2}$ " wide, paste upon the tube with a mixture of starch paste and liquid glue. Remember that the wet paper expands and that the two end marks will fall beyond the scratches on the glass. In drying the contraction will bring the


paper back to the mark. When dry cover the back of the paper with molten paraffin, but not the glass. Press down the upper stopper so that its under face coincides with the zero mark  $m'$  as shown in figure; the remainder of tube being already cut off. Now fasten the tube into the foot by pouring liquid wax into the narrow ring between glass and wood, the slot having been plugged with wood. Finally put 2" of rubber tube upon the capillary and a pinch-cock. Thus the top of the burette will look as shown in Fig. 85. It is in every respect as good as a fully glass blown instrument, except that the scale is not apt to be quite as accurate, yet sufficient for all technical purposes. Never omit the wiring of the rubber, for if the rubber slips off the examination is lost. The analysis of our wood gas can now be proceeded with.

*Note.* All gases are more or less soluble in water. To avoid the error from this source completely the gas must be caught over mercury and measured over mercury—level glass and burette are filled with mercury. The error may be overcome partially, by saturating the water in the burette with the gas, *i. e.*, shaking the water with the gas, and the same with the absorbing liquids. The most soluble in water is  $\text{CO}^2$ , hence it is well to let water in burette saturate itself with  $\text{CO}^2$ , before the analysis.

To fill the burette with gas from the holder will be the first operation, taking advantageously just 100 c.c. to avoid figuring. In Fig 86 the arrangement of apparatus is shown in the moment of trans-

fer, when the gas has passed from *B* into the absorption bulb *A*, and the liquid from the latter into

FIG. 86.



the tank bulb *C*. Stop-cock *1* and pinch-cock *2* are now closed; the level cylinder dropped upon the

table. The operator seizes the frame of the pipette and the scaffold upon which the pipette stands, with both hands and shakes so that the liquid in *A* splashes all about the bulb and remains in contact with the gas; at the least for 5 minutes. Then open the cocks and the gas will flow back into *B* where the vol. is recorded.

The order in which the absorbents are applied is: (1) KOH for  $\text{CO}^2$ ; (2) pyrogallate for O; (3) fuming sulfuric hydrate (oil of vitriol) for the heavy hydrocarbons; (4) cuprous chlorid in ammoniacal or in HCl solution for CO; (5) spongy metallic palladium for H.

In applying these to our gas, the gas from cellulose or wood, we find a shrinkage of volume after each application, except for oxygen. This element is only found when air has become mixed with the gas. The largest shrinkage is that due to hydrogen. After the absorption of H by palladium there is still a large volume of gas left. It must be, therefore, a gas which we have not met with before. We find it to be highly inflammable, and it burns with a pale, nearly colorless flame, if ignited at a platinum tip (for if the flame comes in contact with glass, the former is always of orange-yellow color from the sodium in the composition of the glass).


MARSH GAS, METHANE, METHYLHYDRID,  $\text{CH}^4$ .

When the gas in question burns we can easily demonstrate the forming of  $\text{CO}^2$  and water, hence the gas must contain C and H, but might also contain O.



To get at the relative proportions of  $C + H$  we burn a measured volume within a closed vessel, so that none of the products can escape. Heretofore we used a simple eudiometer, the gas being above mercury. Hempel has constructed a pipette for this purpose. Fig. 87 shows this piece of apparatus. The bulb *A* is shaped into a neck or dome and thence is fused to the capillary. Into the neck two

FIG. 87.



platinum wires are inserted by fusion, the same as in the eudiometer; these wires lead to the pole binders of an induction coil, thence to the battery. *A* being completely filled with water, we transfer from the burette, say 10 c.c. of gas, into *A*. We cannot use pure oxygen, because the explosion might easily shatter the relatively thin and weak bulb. We take air instead, thus diluting the effect of the

explosion by means of the inert nitrogen. 100 volumes of air contain 20.9 volumes O. As regards the volume of air to be admitted we can reason thus: If the total 10 c.c. were carbon gas then we would need 20 c.c. of O, for 1 vol. of  $\text{CO}^2$  contains  $\frac{1}{2}$  vol. C + 1 vol. O. If the whole 10 c.c. were hydrogen, then we should need 5 c.c. of oxygen, a total of 25 c.c. of oxygen or  $\frac{25 \times 100}{21} = 119$  c.c. of

air. As both C and H are present, this volume of air will surely give us an excess of oxygen. Let this volume of air be transferred from the burette in 2 installments, then let the cock *S* be closed. The expansion of the gases, consequent to the explosion, will then throw itself upon the bit of rubber tube, between the pinch-cock and the capillary, hence the rubber tube must be very strongly wired upon the glass, or the rubber will be surely flung off as a bullet from a gun barrel; the pinch-cock also must be wired so that it cannot open under the pressure. A screw clamp is, therefore, preferable to the spring clamp. The rubber tube should be frequently renewed, as frequent expansion is apt to destroy the elasticity of the rubber. Let the pipette be forcibly shaken to effect a mixing of air and gas. Make connection at commutator; the spark is seen passing between the platinum wires, but no explosion occurs; it should take place and mostly does take place. When it does not occur the cause of failure lies in too much dilution, the 10 c.c. of gas taken having already much nitrogen admixed. Under these cir-

cumstances we generate in a voltameter *fulminating gas* ( $O + 2H$ ), and let, say, 10 c.c. into *A*. Explosion is now certain; the fulminating gas will act the same as the cap attached to the fuse, when a charge of dynamite is to be exploded. In our meaning the term explosion = instantaneous oxydation or combustion, or burning of *C* and *H*. A slower burning may be called a fire. The volume of fulminating gas need not to be measured accurately—because it simply disappears in the explosion as water, whose volume is trifling in comparison to the volume of the gas. After the explosion we open cock *S* and note a strong current setting from *B* into *A*, and this means disappearance of a certain portion of the gas. The disappeared portion we name the *contraction*, = *C'*, in this case 30.1 c.c. The volume of the gas mixture *V*, before the explosion was

$$V = G + A \text{ (G = gas 10 c.c. ; A = air 119 c.c.)}$$

$$V = 10 + 119 = 129 \text{ c.c.}$$

After the explosion

$$V' = V - C' \text{ (contraction)} = 129 - 30.1 = 98.9$$

*C'* itself is made up of *H* + *O* which disappeared as  $H^2O$ .

$$2C'/3 = H ; C'/3 = O.$$

We pass the mixture into the KHO pipette and find a shrinkage of 9.8 c.c., which must represent the  $CO^2$  formed from 10 c.c. of the gas. But 1 volume  $CO^2$  contains 1 volume *O*, hence 9.8 c.c. of the 24.9 c.c. of *O* added went into the forming of  $CO^2$ ; which

means that 10 c.c. of the gas contained 5 c.c. or  $\frac{1}{2}$  vol. carbon vapor. We transfer the gas into the pyrogallate pipette and shake 10 minutes to absorb the remaining oxygen.

We find shrinkage = 5.1 c.c. oxygen ; hence, deduct from total oxygen = 24.9 c.c.

Consumed for  $\text{CO}^2$  = 9.8 c.c.

Remaining oxygen = 5.1 c.c.

---

14.9

Therefore in contraction  $C'$  enter  $24.9 - 14.9 = 10$  c.c. of  $O$ . And as hydrogen =  $\frac{2}{3}C'$  ; oxygen  $\frac{1}{3}C'$  ; the 10 c.c. of  $O$  correspond to 20 c.c. of hydrogen, or if the original 10 c.c. of gas be made = 1 vol. it follows that 1 vol. of our unknown gas contains  $\frac{1}{2}$  vol.  $C + 2$  vols.  $H$  ; or 1 vol.  $C + 4$  vols.  $H$  and its formula or symbolic expression is



Here we have a body representing a compression of 2.5 : 1 ; 5 volumes of  $C + H$  compressed into 2 vols.  $\text{CH}^4$ . We have not had, therefore, a similar compound. Nevertheless the fact is proved by the specific gravity which is by calculation

$\frac{1}{2}$  volume of carbon vapor = 0.4146

2 volumes of hydrogen = 0.1382

---

0.5528

Several experimenters have found by independent methods, that is, both by weighing the gas, and by its velocity of diffusion, the number 0.5589 which is very close to the calculated figure. The name

marsh gas (*sumpf gas* in German) refers to the evolution of this same gas from swampy meadows where it sometimes gives rise to the peculiar phenomenon of the *will-of-the-wisp* (*irr-licht* = erring or wandering light), when by accident the gas becomes inflamed at one spot and the pale flame then setting fire to the adjacent bubbles thus produces the impression of a wandering or jumping flame, once thought to be *spirits*. In the summer months the gas always comes abundantly when the mud in swamp rivers is disturbed or poked into.


Marsh gas has neither odor nor taste. Mixed with air it causes no ill effects when breathed, is therefore not a poisonous gas such as CO, or a suffocating gas such as CO<sup>2</sup>. 100 volumes of water absorb at 20° C., *i. e.*, the ordinary temperature of the air, 3.5 volumes of marsh gas.

When mixed with 8 to 10 volumes of air, or 2 volumes of oxygen, the gas explodes with great violence; but the temperature at which it ignites is higher than that at which hydrogen or hydrogen sulfid explode. These gases require only low red heat, while marsh gas requires yellow or white heat, showing that the elements C + H cling very strongly together.

1 volume CH<sup>4</sup> + 3 or 4 vols. air does not explode; with 1 vol. CH<sup>4</sup> + 5½ to 6 vols. air the explosion is weak. And so again with 14 volumes of air, the explosion is weak; with still more air the CH<sup>4</sup> merely burns over the flame of a candle or lamp—no explosion. Now all this is knowledge of extreme importance to

the engineer who engages in coal-mining. For  $\text{CH}^4$  is the gas which issues from the fissures of the coal beds; and causes the fire danger in those mines, from its life-destroying explosions. The simplest instrument for the detection of the danger, *i. e.*, the presence of the marsh gas in the mine, breasts, stopes, levels, was devised by Sir Humphrey Davy, eighty-eight years ago. He it was who first inquired into

FIG. 88.




the nature of a flame and thus discovered a means against the ignition of the inflammable mixture of air and marsh gas. Let *A*, Fig. 88, be the basin of an ordinary oil lamp with a wick holder and wick *4*; the oil level at *3*, the flame *1*. *B* is fine mesh wire cloth made into a truncated cone and the cone is held by the brass ring *8*. Thus no air can get to



the flame unless it pass through the gauze, and the product of the combustion must also pass through the gauze, since the top of the cone is closed by gauze. If the air contains  $\text{CH}^4$  admixed, the flame begins to tremble, becomes longer and a bluish mantle forms around it. The whole interior may be filled with such a blue flame. This is due to the fact that with the deficiency of air the carbon only burns to  $\text{CO}$  and not to  $\text{CO}^2$ . But why does not the flame ignite the outside gas mixture? Simply because the wire gauze absorbs the heat so quickly that the temperature outside of it is too low to ignite the gas. The wick requires trimming from time to time and snuffing. These operations are performed with the wire 2, which passes through a stuffing box in the bottom of the lamp. The gauze must never be removed inside of the workings. Unfortunately the light given by such an affair is very weak and thus the men are tempted to remove the gauze and then comes the disaster.

*Theoretical importance of marsh gas.*  $\text{CH}^4$  may be written



Four hydrogen chemical units evenly balancing the 4 affinity bonds of the carbon unit; in other words we say the marsh gas being a fully satisfied chemical complex, all other combinations of carbon

and hydrogen when saturated must be of the same type. Numerically we can express it  $C^nH^{2n+2}$ . When we found that paraffin showed the composition  $C^{24}H^{50}$ , we mean that we have here a compound built upon the *type* of marsh gas, yet while  $CH^4$  equals a percentage composition of 75C + 25H, the percentage in paraffin will be 85.3 C + 14.7 H. All the members of the *marsh gas* or *paraffin series* have been found either as existing in natural bodies or they have been prepared by artificial splitting.

$CH^4$  = methane—marsh gas—a permanent gas.

$C^2H^6$  = ethane, a condensible gas.

$C^3H^8$  = propane, a light liquid.

$C^4H^{10}$  = butane (in butter).

$C^5H^{12}$  = pentane (five).

$C^6H^{14}$  = hexane (six).

$C^7H^{16}$  = heptane (seven).

$C^8H^{18}$  = octane (eight).

$C^9H^{20}$  = nonane (nine).


$C^{10}H^{22}$  = decane (ten).

} Liquids contained in  
gasoline, benzine,  
and kerosene,  
also in tar.



$\begin{array}{c} | \quad | \quad | \\ C^{24}H^{50} \end{array}$  = paraffin ; a white, hard solid.

With the increase of C in the molecule, rise the specific gravity and the boiling-point. There are reasons for assuming in marsh gas the existence of a group ( $CH^3$ ), in which there is still closer contact between the atoms than in  $CH^4$ . This latter becomes then the hydrogen compound of the form  $(CH^3)H$ . The group  $(CH^3)$  = methyl, is furthermore to be considered a *radical* or complex element, such as  $(NO^3)$ ,  $(SO^4)$ ; only with the difference that

this radical is of a positive or metallic character. It replaces hydrogen. Every elementary molecule is composed of at least two atoms. That is, hydrogen in the free state is not H but  $H^2$ ; chlorine not Cl but  $Cl^2$ . Then in the hydrogen molecule H-H we can replace 1H by  $(CH^3)$  as its equivalent.




Or, again,  $H^2(SO^4) + 2(CH^3)H = 2(CH^3).(SO^4) + 4H$ ; the latter compound is then methyl sulfate, same as sodium sulfate. In ethane  $C^2H^6$ , the older conception saw the radical ethyl  $C^2H^5$  united to 1 hydrogen. But it is simpler to think of it as a coalescence of 2 methyl groups.



Two methyl groups joined by  $CH^2$  which latter is the lowest member of the series  $C^nH^{2n}$ .

*Butane,  $C^4H^{10} = (C^4H^9).H = \text{butyl hydrid}$ .* But on the methyl hypothesis there will be two combinations, which have been actually proved to exist, two bodies having exactly the same percentage composition, but distinct properties, to wit:

$CH^3.CH^2.CH^2.CH^3$ , normal butane; and



Such bodies as these, of equal atomic or percentage composition, are called *isomerids*. Isobutane is the *isomerdi* of butane.


For pentane  $C^5H^{12}$  there are three isomerids :

$CH^3.CH^2.CH^2.CH^2.CH^3$ , normal pentane.


$CH \begin{array}{l} \swarrow CH^3 \\ \searrow (CH^2.CH^3) \end{array}$ , di-methyl-ethyl-methane,

and  $\begin{array}{l} CH^3 \\ CH^3 \end{array} > C < \begin{array}{l} CH^3 \\ CH^3 \end{array}$ , tetra-methyl-methane.

In the last we imagine a marsh gas molecule




in which every hydrogen atom has become replaced or substituted by the methyl group. The second isomerid can be represented, to show the similarity, thus




I beg to remind you that this so-called structural representation is not to mean a real picture—for in reality we have to deal with three dimensions, not with two as on this sheet of paper. This is an attempt to express symbolically the difference in properties of the *isomeric* bodies.

In the light of this exposition on marsh gas, we will reconsider the wood alcohol and the acetic acid. For the wood alcohol we had the atomic proportion  $CH^4O$ , it being then suggested that this formula may be written  $CH^3(HO)$ , and now we see that the methyl alcohol is a marsh gas in which 1 H is replaced by the radical (HO) hydroxyl.



For acetic acid  $\text{H.C}^2\text{H}^3\text{O}^2$  we can write




The group  $\text{CO.OH}$ , carbonyl-hydroxyl, is monovalent; by the entering of this group into a hydrocarbon, the latter takes on the properties of an acid. The lowest form is *formic acid* (*ant acid*)  $\text{CHO}^2.\text{H}$  which is found with the acetic acid. Here we can say that the group  $\text{CO.OH}$  is simply coupled with one H.

#### WOOD OR CELLULOSE UNDER PRESSURE AND HEAT.

If compressed cotton, or a stick of dry wood, or dry sawdust be enclosed in a strong glass tube as shown at Fig. 89, *A* in *figure 1*, and if the tube

FIG. 89.



be then fused together as at *C*, *figure 2*, over a blast lamp and the part *D* be pulled off, the wood *A* will

be enclosed air-tight as in *figure 3*. Care must be had that the wall of the tube remains throughout of even thickness. If this tube be then placed within an air-bath, and the temperature be gradually raised to  $320^{\circ}$  C. as indicated by the air thermometer, Fig. 90, then it is evident that the distillation of the

FIG. 90.





FIG. 91.



wood proceeds under increasing pressure : First, by the expansion of the air ; second, by the expansion of the gases and vapors which arise from the wood at this temperature. Let the condition of things remain thus for twenty-four hours. Let cool slowly. On examining the tube we find the wood converted into a jet-black shining mass ; the cellulose structure is effaced altogether and the resemblance to *soft coal* is unmistakable. We open the tube very cautiously (after taking its weight) as follows : Since the pressure is still high (presumably) a sudden breaking



of the tube might have a very shattering effect (like a boiler explosion). Therefore, we approach the pointed end of the tube to a strong flame, as shown in Fig. 91. In measure as the glass nears red heat, it will become soft, and expand under the interior pressure until at the very tip it will be blown out, giving vent to the compressed gases. On weighing the tube, now, it is found that about 2 per cent. of the weight of the wood has disappeared, that much having been converted into permanent gas. Now we wash carefully the mass in the tube by means of alcohol (to dissolve any tar-stuff), and after drying in a current of dry air, find again a diminution in the weight of about 3 to 4 per cent. Altogether about 20 per cent. in weight of the cellulose has vanished. The importance of this experiment will appear in the next section.

## CHAPTER XVI.

### MINERAL COAL AND ITS CHEMISTRY.

WITHOUT mineral coal the industrial development of modern times would have been impossible. Some 250 millions of tons are now mined annually in the United States, much more than all the other minerals together. The conditions under which coal is found, and its association with other rocks, form a subject of *stratigraphic geology*. The points to which your attention is here called refer to the properties of coal and the uses to which coal can be and is applied owing to such properties.

We distinguish (1). *Hard coal or anthracite*. (Greek *anthrās* = coal.) Color intensely black, luster more or less bright. Fracture smooth and spheroidal. Hardness considerable; the pick does not produce much effect; drilling and blasting is necessary.

(2). *Soft coal, bituminous coal, semi-bituminous, blacksmith coal*. Much softer than anthracite, color black, but color of fine powder is brown. Cleaves or breaks into prismatic pieces. When heated in glass tube gives off *dense yellow vapors* which separate into a liquid and into a combustible gas, whereas anthracite gives off *no vapors*, and only very little gas. If the coal melts into a black, thick liquid, it is called *bituminous* (*bitumen* being the Greek word for the

natural pitch also known as *asphaltum*). If the coal merely softens in the heat, does not become quite liquid, then we call it *semi-bituminous*, (half pitchy).

(3). *Cannel coal*, has a black color but differs by absence of luster from the other varieties. It breaks with an irregular surface. When heated it does not melt nor become soft, but yields both condensable vapors and bright burning gas in abundance, (*cannel* is the Scotch of candle). This coal is not as abundant as the other varieties and is of higher value, because it furnishes a larger volume of illuminating gas. It is only used for gas-making.

(4). *Brown coal, lignite*. Dark-brown color, dull in appearance, very soft, can be shoveled from the pit, shows the cell structure of wood and hence the name lignite (*lignum* = wood). When heated it does not soften, but gives gas and tar, although less of these than cannel coal; and more water. It is usually mixed with sand and clay to a much larger extent than the other coals. The beds lie near the surface, that is, they belong to more recent geological times. Very abundant all over Western United States, Central Europe, Asia and Africa. The cities of Northern Germany use this material exclusively for fuel, on account of its cheapness.

(5). *Peat, bog, turf*. Brown or brown-black in color, very loose in structure and crumbly. Acts like brown coal, when heated after having been dried. This material is found always in flat regions, where the water cannot drain off and where no grasses can grow on account of the wetness. But

on the other hand, the different varieties of mosses and algae develop and grow with astonishing rapidity, one generation on top of the other, each generation being very short-lived. Being so near the air the dead mosses undergo a partial rotting with the formation of marsh gas and carbon dioxide. There are bogs known to be 50 feet thick and more. Ordinarily they do not exceed 3 feet in thickness. Yet they furnish to many localities their only fuel—Ireland, Holland, North Germany near the sea. The higher land of Western New York, where the Hudson, the Alleghany, and the Delaware rivers have their beginning, contains many peat bogs, which are more or less utilized. On the flat tops of very high mountains such bogs have been found. By a washing process the peat substance can be somewhat freed from the sand, and it can then be converted into cakes by pressing. Such cakes, when quite air-dry will give a hotter fire than wood, weight for weight.

*Composition of coal.* We find the ultimate or elementary composition by the same procedure which we followed with the cellulose. We burn a known weight, *W*, with copper oxide and oxygen gas and collect the products of the combustion, conveniently for measuring or weighing. We find invariably the same elements to wit: Carbon, hydrogen, oxygen, nitrogen, sulfur. The two latter are *not* in cellulose, but they are found in other parts of plant structure. Nitrogen is *always* contained in the protoplasm, and sulfur sometimes; without the protoplasm no plant can develop—it is the blood of plant

life. The seeds of plants always contain much nitrogen (15 to 17 per cent.). Coal contains from 1 to 3 per cent. of nitrogen. The sulfur varies between wider limits. The sulfur can be in *union* with the carbon and the hydrogen, and is then *invisible*; or combined with iron as yellow pyrite, and is then readily visible.

Coal always leaves a residue after the carbon and hydrogen have been volatilized by oxidation into  $\text{CO}^2$  and  $\text{H}^2\text{O}$ . The residue is called *ashes*, because wood leaves ashes. The two kinds of ashes are very unlike. From coal ashes water does not extract potash, nor any other body; coal ashes are quite insoluble; no alkaline reaction whatever. Neither  $\text{HCl}$  nor  $\text{HNO}^3$  nor  $\text{H}^2\text{SO}^4$  dissolve it, only  $\text{HF}$ . The ashes in fact contain chiefly the oxyds of silicon and aluminum  $\text{SiO}^2$ ,  $\text{Al}^2\text{O}^3$ , and  $\text{Fe}^2\text{O}^3$  when the ashes have a brown color. The clinkering or semi-fusion of the coal ashes is due to  $\text{Fe}^2\text{O}^3$  which acts as a flux upon the other oxyds; white coal ashes never show clinkering. The following analysis, made by me lately, gives the ultimate composition of a soft coal from Kansas. Carbon = 75.35, hydrogen = 5.50, oxygen and nitrogen = 10.10, sulfur = 1.54,  $\text{SiO}^2$  = 4.35,  $\text{Al}^2\text{O}^3$  = 2.70 (being together ashes equal 7.05, snow white) moisture = 0.45.

Notice the total absence of iron oxyd, whence it follows that the sulfur must be combined with the carbon and hydrogen. This coal belongs to the semi-bituminous variety verging upon cannel; for

the residue merely adheres slightly after having been exposed to a high yellow heat, and like cannel it has a dull, black color ; powder brown.


The *proximate composition* of the different varieties is but imperfectly known, or rather guessed at. I mean by proximate composition the molecular structure—the manner of combination of the elements. You can do no better than to imagine the coal to be an intimate mixture of solid hydrocarbons, oxy-hydrocarbons, sulfo-hydrocarbons, nitro-hydrocarbons, amorphous carbon, and mineral particles constituting the ash. The different varieties of hard and soft coal arise from the preponderance of one or more of the above groups of molecules. Reasons for this hypothesis are : (1) The crystalline structure of the coal, revealed in thin translucent plates or sections under the microscope ; (2) The actions of solvents upon the coal, such as ether, benzole, carbon disulfid, potassium or sodium hydroxyds.

*Origin of coals.* That cellulose is the original material there is no reason to doubt ; all hypotheses or theories start with this base. Generally, however, geologists assume that the material for the coal beds consists in the successive growth of tropical forests one on top of the other. My own theory differs from this. The chief reasons are that in many places we find trees standing upright in the coal beds, reaching even into the sandstone or slate strata lying over the coal bed as shown in Fig. 92. Here a section is reproduced from an English coal



mine. *a* is limestone, *b* is fire-clay (under clay), *cc* the lower and upper benches of a coal seam, *e* black coal slate, *f* sandstone, *g* brown slate. *ttt* are fossil trees, whose bark is also coal, but whose interior is sandstone, because these trees belonged to the class of giant reeds, *calamites lepidodendron*, *sigillaria*, etc., and therefore contained a pithy interior and a very strong fibrous rind, very resisting to chemical change. *d* is a band of slate separating the coal-

FIG. 92.



bed into the two benches. The strata *a* and *b* were in horizontal position during the coal-forming times and the conditions of level equal to that of very shallow basins very near the sea level; that is, the general conditions were those of a tropical swamp as we find them in our time along the Amazon River in South America. At first these conditions were favorable to the growth of the great ferns and the gigantic reeds. Later on, the land sinking very slowly, the swamp became too wet for this vegetation and in its stead algae—the lowest type of plant life—which you find always in the shallow pools of

our present swamp woods, the green and brown threads began to flourish abundantly, luxuriantly. An alga is a plant consisting of one cell or of an aggregate of cells, of which however each cell remains a life unit. Each cell has a thin wall of cellulose enclosing a liquid interior in which there is a floating patch of protoplasm, the cell-nucleus or cell-kernel. While these generations of very rapidly growing cells died and accumulated on the bottom of the pool, or rather upon their dead predecessors, and being under water could not *rot*, the rivers or creeks emptying from higher ground into the swamps brought the fine sand and clay which settles very slowly, as you well know from the rivers remaining turbid long after a freshet. In time this material—the *silt*—settles and mixes with the vegetable ooze, and there we have an explanation of the intimate admixture of the ash particles with the coal, and also an explanation of the strong variation of the ash percentage in the different parts of a coal bed. Whenever a slate band occurs in the coal, according to this theory, we presume that an unusual freshet carried the silt faster into the basin than the algae could accumulate, in fact the muddiness interfering with rapid growth. The silt also settled into the hollow trunks of the trees, thus preserving them against collapse by external pressure.

My theory of the algae accounts also for the high percentage of nitrogen, which we find in the coal, because the relative percentage of protoplasm to cellulose or of nitrogenous substance, is larger in

the algae than in the complex cell-structures of trees. During the accumulation of the ooze a decomposition of the dead algae cells began to set in, alike to that which we now-a-days observe in the peat bogs, by which the percentage of carbon in the residue steadily increases, while hydrogen and oxygen, notably the latter, decrease:  $\text{CO}^2$  forming and  $\text{CH}^4$  and  $\text{H}^2\text{O}$ . At last, the ground sinking more rapidly, the influx of silt increases and the vegetation stops, the sea finally encroaches upon the swamp and the materials for sandstone or limestone are brought in. They cause a steadily increasing pressure, under the influence of which internal heat arises, which cannot dissipate as the rocks are very bad conductors of heat. Thus the plant material comes gradually under the conditions of the experiment upon cellulose, described a few pages back. Wherever the pressure was greatest the change towards *carbon* was greatest. Thus we find in Eastern Pennsylvania only anthracite with 90–94 per cent. of carbon, because the side pressure upon the strata was so great, that the latter became greatly bent and even doubled upon themselves as shown in Fig. 93, whilst in the Western States the strata remained in their original horizontal position, as seen in Fig. 94, except in Colorado, and hence we find anthracite in the latter state. The dotted lines in Fig. 93 denote that part of the coal bed which has been removed and lost by surface destruction and the forming of the present topographical outlines. All the details of the structure of the coal

measures belong to geology ; only sufficient had to

FIG. 93.





FIG. 94.



be introduced here to make the chemical theory intelligible.

#### DISTILLATION OF COAL—COKING PROCESS.

By distillation of cellulose—or wood—we obtained charcoal, tar, pyroligneous acid,  $\text{CO}$ ,  $\text{CO}^2$ ,  $\text{CH}^4$ ,  $\text{H}$ ; and since coal is derived from cellulose

we may and should expect similar, if not identical, products. In order to test this proposition we rig up the set of apparatus, Fig. 76, page 275. The tube *T* we charge with the coarsely-powdered coal, so that when the tube lies in horizontal position and has been tapped upon, the coal only fills one-half of the tube. Why? Because at red heat the bituminous and semi-bituminous coals swell up, thus clogging the tube to the escaping gas, which latter, with increasing pressure, will invariably break the tube. We start heating at the front by using the diaphragm or shield *S*. First we note a heavy yellow vapor, from this condenses a brown liquid in the receiver *3*, and the bell *B* fills itself with gas. The first portion of gas we allow to escape, because it is mixed with the air in *T* and *3*. Anthracite gives no vapor, no condensing tar, only a relatively small volume of gas; because anthracite has already undergone the distillation under the influence of great pressure. When, at bright red heat, the evolution of gas becomes very slow, or stops altogether, we disconnect the receiver *3* from *T* and *B* from *3*. We pull the tube *T* from the furnace, let it become cool, and then break the tube. We find the residue more or less bright, porous, dark or light grey in color. With large pores the stuff is more or less friable, with small pores it becomes hard and tough. For bituminous and semi-bituminous coals the weight of the residue is from 50 to 65 per cent. of the original weight of the coal. The technical name of this residue is *coke*, which



word is derived from to cook, and may have been merely a provincial substitute for cake. When brought up to a red heat in a current of air the coke burns *without* making a visible flame, although it still contains a remnant of hydrogen and oxygen, for the complete change of coal to carbon depends upon the temperature and time. At a white heat (in fire-clay crucible), the last remnant of hydrogen can be eliminated. The ordinary coke is, therefore, a mixture of the ashes with carbon (amorphous and also graphitic), and more or less hydrocarbon.

Coke is required as fuel in blast furnace work, especially in the high-stack-furnaces for the reduction of iron ores. Why? Because if coal were used, the latter would convert itself into coke in the furnace, would cause a *loss* of heat energy (elimination and decomposition of the hydrocarbons and oxy-hydrocarbons requires heat addition from outside sources), and the coke thus forming would cement the pieces of ore and flux into a solid cake, through which the large quantities of nitrogen and carbon monoxyd are blown in at the bottom of the furnace could not pass—the furnace would clog or choke and eventually extinguish itself.

For this need of the blast furnaces, immense quantities of coal are converted into coke. The apparatus for making coke is known as coke oven, not kiln or furnace, but *oven*. Why? Because the first coke was made in Dutch bake-ovens. The so-called *bee-hive* coke oven of the present time is merely a slightly modified bake-oven. In this oven all the gas and




the tar are wasted. In the scientifically constructed ovens, the gas and tar are utilized. Different ovens have been constructed in Europe in great profusion. Those mostly used now are the Solvay oven, and the Hoffman oven; they have also been introduced into the United States. The detail of coke-making belongs to *metallurgy*. We turn now to the contents of the receiver (3). As in the distillation of wood we find two liquids, one oily the so-called coal tar, one watery of light-brown color. This watery liquid smells of *ammonia*, and turns red litmus paper blue. Thus it is the reverse of pyroligneous acid from wood distillation. Addition of acid to the ammonia water produces effervescence:  $\text{CO}^2$ ,  $\text{H}^2\text{S}$  are given off, hence the water contains ammonium carbonate and ammonium sulfid. The occurrence of the ammonia proves the presence of nitrogen in the coal. The absence of the acetic acid may be explained from the smaller percentage of oxygen in the coal and from the higher temperature needed to break up the coal, a temperature at which  $\text{C}^2\text{H}^3\text{O}^2.\text{H}$  breaks up into  $\text{CH}^4 + \text{CO}^2$ . The principal market for ammonia being that as *fertilizer*, the cheapest way of extracting the former is to convert it into *sulfate*. The water is neutralized with  $\text{H}^2\text{SO}^4$  and the solution evaporated by means of the waste heat from the ovens. The result is dark-brown, crude sulfate. This is redissolved in the requisite quantity of boiling water, filtered through a bed of charcoal to remove the tar which separates during evaporation, and the liquid is run

into flat basins, where the sulfate crystallizes on cooling. This second product is still yellowish, but good enough for the market.


*The coal tar*, the oily portion of the condensed vapors, is composed essentially like the wood-tar. But certain valuable hydrocarbons are contained in the coal tar in larger percentage. These hydrocarbons are: benzol ( $C^6H^6$ ), toluol ( $C^7H^8$ ), phenol ( $C^6H^5(OH)$ ), naphthaline ( $C^{10}H^8$ ), anthracene ( $C^{14}H^{10}$ ). When the tar is subjected to distillation, light oils pass over first up to a temperature of  $180^\circ$  C. Benzol, toluol, phenol are contained in this portion; at a higher temperature oil passes over which solidifies on cooling. The cooled mass can be recrystallized from an alcohol solution, or it may be purified by sublimation. In this purified state it forms large thin crystals in form of plates with the luster of mother of pearl and a strong peculiar odor. This is the *naphthaline* of trade and is largely used to protect woolen and fur goods against insects: *moth-balls*. Only the oil which passes over between  $80^\circ$  and  $100^\circ$  C. is used for the manufacture of benzol. Between  $100^\circ$  and  $130^\circ$  C. toluol and phenol with some benzol pass over. Pure benzol is a colorless very mobile liquid; boils at  $82^\circ$  C. and crystallizes at  $0^\circ$  C. Specific gravity = 0.85.

*Nitrobenzol*,  $C^6H^5.NO^2$ . Is obtained by treating the benzol with a mixture of conc.  $H^2SO^4$  and very conc. or fuming  $HNO^3$  in cast-iron cylinders, which can be cooled by running water, because the reaction is very energetic; 1 hydrogen of the  $C^6H^6$  is removed as water and nitrobenzol results.




Nitrobenzol is a yellowish heavy oil which boils at  $205^{\circ}\text{C}.$ ; and has a pleasant odor resembling that of the oil made from bitter almonds. If an excess of  $\text{HNO}^3$  is used the dinitrobenzol  $\text{C}^6\text{H}^4(\text{NO}^2)^2$  results.

*Anilin, C<sup>6</sup>H<sup>7</sup>N.* A small quantity of this highly interesting substance is already contained in the tar. But from nitrobenzol it can be obtained in any desired quantity. Anilin forms a colorless liquid, diffracts light strongly, has a peculiar odor and burning taste. The oil boils at  $182^{\circ}\text{C}.$  and solidification sets in at  $8^{\circ}\text{C}.$  Specific gravity = 1.02. Is very slightly soluble in cold water, more so in boiling water, but quite soluble in alcohol, ether, carbon disulfid, and coal oil. It burns with a very smoky flame. Is very poisonous. Chemically anilin may be considered as ammonia in which 1 H has been replaced by the hydrocarbon radical *phenyl* thus




Like ammonia it unites with HCl and other acids and forms salts:  $\text{C}^6\text{H}^5.\text{H}.\text{HNHCl} = \text{anilin chlorid}.$  These salts are mostly easily soluble in alcohol and crystallize readily. Anilin is a fine example of a *complex base*.

*Methylanilin* is another base, arising from a second hydrogen being replaced by methyl, thus:



This body, known as "Mauve de Paris," colors silk and wool a fine violet color.

*Rosanilin, fuchsin.* This was the first splendid dye-stuff prepared from coal-tar through the way of anilin, by means of oxydation; usually  $\text{As}^2\text{O}^5$  is used as the oxydizing agent. 100 parts of anilin oil are poured slowly into 150 parts of a water-solution holding 75 per cent.  $\text{As}^2\text{O}^5$  in an iron vessel with stirring apparatus. The temperature is raised and kept for five hours at  $182^\circ \text{C}$ . Water and unchanged anilin dissolve during this period. The semi-fluid residue has a bronze color, and from it the dye-stuff is extracted by boiling water, filtered, under pressure, through felt. Solution contains the rosanilin as arseniate and the  $\text{As}^2\text{O}^3$  which is formed during the process of oxydation. By saturating the solution with  $\text{NaCl}$  (equal in weight to the residue), the hydrochlorid of rosanilin forms and  $\text{Na}^2\text{HAsO}^4$ . *Red crystals* fall out, in measure, as the liquid cools. By recrystallizing this first product a higher grade is obtained. The red crystals, being the chlorid of rosanilin, are known in the dye-works as *fuchsin*. By acting on the water solution with  $\text{NaOH}$ , the base *rosanilin* is obtained as a white precipitate, which becomes intensely red in contact with any acid. The composition of *rosanilin* is  $\text{C}^{20}\text{H}^{19}\text{N}^3$ . It forms thus:



The  $3\text{O}$  are furnished by  $\text{As}^2\text{O}^5$  or any other oxy-

dizing agent. But we see that another body is here contained, the *toluidin*, besides the anilin. Remembering, however, that the raw oil contains benzol and toluol, the phenomenon is explained.

The crystals of fuchsin are green-golden in appearance, like a brilliant metal. They dissolve in water with intense red color. If well cleansed silk or wool be hung in such a solution, the liquid becomes colorless by degrees, all the coloring fuchsin will have transferred itself to the fibre, producing thereon the beautiful red tints according to the quantity transferred. The color fixes itself, does not require a *mordant* or fixing agent.

*Mauvanilin*,  $(C^6H^5)(C^6H^5)(C^7H^7)N^3.HCl$  gives an orange-yellow dye. A great many other beautiful dye-stuffs have been produced by further substitution of hydrogen in the base by other radicals, as anilin, toluidin, or simply methyl, ethyl, and others. My main purpose in devoting so much space to this subject was to impress upon you the possibilities lying hidden in such a material as the ugly, bad-smelling coal-tar, and the great fortunes which have been made by utilizing it in the right way.



## CHAPTER XVII.

### THE GASES FROM THE DISTILLATION OF COAL— MANUFACTURE OF ILLUMINATING GAS AND GAS COKE.


The contents of the bell jar are at first cloudy from exceedingly small particles of semiliquid bodies, which will condense after a time, forming a thin layer of tar upon the retaining water surface. If now the gases be subjected to the analysis by absorption which has been given in detail under cellulose or wood, it will be found that the composition by quality does not differ much; the relative quantities differ considerably, and even very much when the gas from the early distillation is compared with that which is given off at the end of the operation. We find  $H$ ,  $CH^4$ ,  $CO$ ,  $CO^2$ , non luminous:  $C^2H^4$ ,  $C^3H^6$ ,  $C^4H^8$  and  $C^2H^2$ , (acetylene) as luminous gases.  $NH^3$ ,  $SH^2$ ,  $CS^2$ ,  $CN$  as impurities.

Of the luminous or light-producing hydrocarbons *ethylene* ( $C^2H^4$ ) olefiant gas (oil-making gas) is the most important; of propylene ( $C^3H^6$ ) there is least; of butylene  $C^4H^8$  there is usually from  $\frac{1}{2}$  to  $\frac{1}{3}$  as much as of ethylene. These hydrocarbons combine with chlorine, bromine or iodine, thus:

$C^2H^4 + 2Cl = C^2H^4Cl^2$ , *ethylene chlorid* an oil-like liquid (hence the name olefiant or oil-making.)  
Spec. Gr. = 1.174.



At red heat ethylene breaks up, thus :





It is this action which we call *dissociation*, that accounts for the smoking of a gas flame. If oxygen is present both C and CH<sup>4</sup> are oxydized to 2CO<sup>2</sup> + 2H<sup>2</sup>O. Bromine acts like chlorine upon ethylene : C<sup>2</sup>H<sup>4</sup> + 2Br = C<sup>2</sup>H<sup>4</sup>Br<sup>2</sup> an oily liquid as the preceding one. Propylene and butylene are acted upon similarly : C<sup>3</sup>H<sup>6</sup> + 2Cl = C<sup>3</sup>H<sup>6</sup>Cl<sup>2</sup>; C<sup>4</sup>H<sup>8</sup> + 2Cl = C<sup>4</sup>H<sup>8</sup>Cl<sup>2</sup>, propylene chlorid, butylene chlorid. Hence it follows that we can remove these three gases from a mixture of gases by shaking the mixture with bromine water in a suitable gas pipette. The higher we find their percentage in a given gas the more we are sure of a bright light, provided that the burner be suitably constructed.

#### PLAN FOR GAS WORKS.

Figs. 95 and 95a give the essential pieces of apparatus for the production of illuminating gas by distillation of coal, in ground plan and length elevation. *R, R', R<sup>2</sup>* represent a fire-brick retort and ovens for heating them to a yellow heat. *H* is a sheet-iron pipe 18" to 24" in diameter and known as the "hydraulic main." The elevation shows how the retort connects by a goose-neck pipe with this main and also that the pipe dips under the water level of the main, thus producing a light pressure upon the gas in the retort. Most of the condensible constituents of the gas become liquid in contact with the liquid in the

FIG. 95.





pipes. At *N* there is provided an overflow which drains into a vertical tank or cistern ; thus the level remains always the same in the main. Steam, tar, ammonium salts are condensed largely. A pipe *C* leads from top of main to the purifying towers *S, S', S<sup>2</sup>* which are technically known as *scrubbers*. *S, S'* are *wet* scrubbers, because the gas must pass the extensive surface of the horizontal trays over which flows a film of cold water. All the ammonium compounds and all the tar are removed from the gas in these wet scrubbers, but the gas still contains a notable quantity of hydrogen sulfid. The latter is taken care of in *S<sup>2</sup>* which is a *dry* scrubber, for here the trays are covered each with a layer of *Laming's mass*, to wit: a mixture of  $\text{Ca}(\text{HO})^2$  with  $\text{FeSO}^4 + 7\text{H}^2\text{O}$  (copperas).  $\text{Ca}(\text{HO})^2 + \text{FeSO}^4 = \text{CaSO}^4 + \text{Fe}(\text{HO})^2$  ; it is the latter, ferrous hydroxyd, which is the active agent. At first  $\text{Ca}(\text{HO})^2$  (dry slaked lime) was used alone.  $\text{Ca}(\text{HO})^2 + \text{H}^2\text{S} = \text{CaS} + 2\text{H}^2\text{O}$ . But the action is slow. When the book-backs in the public libraries of London and other cities began to crumble away, the cause was traced to the gas flames and more specially to the sulfuric oxyd produced by them, *i. e.*, by the burning of  $\text{H}^2\text{S}$  to  $\text{SO}^3$ . Then Laming invented the mixture and thus checked the evil, without knocking it out altogether. The carbon disulfid  $\text{CS}^2$  has, so far, resisted all attempts at absorption in the scrubbers. The dry scrubbers must be provided in duplicate or triplicate, because they need frequent renewing of the Laming mixture, which becomes

*foul* as the men say. The gas is now ready to flow through the pipe  $C'''$  into the holder  $G$ . The holder is a sheet-iron tank of circular or cylindrical shape. It is closed at the upper end and dips with the open lower end into water of the walled and cemented cistern  $C$ . The tank  $G$  is held in place by 6 or more guide rolls, and is counterbalanced by the 6 weights  $W, W$ , etc. These consist of cast iron disks, superposed, so that the bell may be made to press upon the gas at any pressure, by adding or removing a disk, from each or to each, of the 6 weights. By adding disks the bell may take the function of a suction pump thus causing the gas to overcome the friction of scrubbers, during the progress of the distillation. The elevation explains the entrance of the gas at *in* and the outflow at *out* if the main valve  $V$  is open. This valve is always accessible through the pit  $P$ . The dimensions of the plant follow from the rate of consumption of the gas. As one 15 candle-power burner consumes 5 cubic feet of gas per hour, and as the burners are needed on the darkest day for 8 hours, and as each family uses on the average 4 burners, we get 160 cubic feet per family per day. 1000 families require 160,000 cubic feet + 10 per cent. for street lighting, total 176,000 cubic feet, hence 18 tons of coal will be required per day giving roughly 9 tons of coke. A bell 30 feet in diameter and 20 feet high will hold 16,000 cubic feet. Ten such would be required to store the 160,000 cubic feet. However the bell serves more as a regulator than as a storage. The

retorts are kept going all through the hours of largest consumption. One holder is sufficient for each 1000 families. Holders of 100 feet diameter and 25 feet high have been built in large cities. The retorts are made of fire-brick material and have a  $\cap$  section. They are 5 to 6 feet long, 18 to 24 in. wide (inside), 12 to 15 inches high. Thickness of bottom 3 to 4 inches, of sides and top  $2\frac{1}{2}$  to 3 inches. The average daily capacity per retort is 5000 cu. feet; hence 32 retorts will be required for 160,000 cu. ft. daily production. The retorts are best arranged in batteries of 7 each with one common fire-place.

#### WATER-GAS.

(a) *Water-Gas.*  $\text{H}^2\text{O} + \text{C} = \text{H}^2 + \text{CO}$  at yellow or white heat. The mixture of hydrogen and carbon monoxyd burns with colorless flame, hence the gas must be made luminous by the addition of gasoline or hydrocarbons of the olefine series. Prof. Lowe of Norristown, Pa., introduced the application of this reaction into the gas industry about 1873. It displaces the distillation process, but has not been introduced in many gas works until more recently in a modified form.

The action  $\text{H}^2\text{O} + \text{C} = \text{CO} + \text{H}^2$  is *endothermic*, heat-consuming.


The action  $\text{C} + 2\text{O} = \text{CO}^2$  is *exothermic*, heat-producing.

Hence the process of preparing water-gas is necessarily a double one. Let 1, 2, Fig. 96, be two



fire-brick cylinders held each in a sheet-steel mantle. Let these cylinders stand upon iron pillars, 10, 10, 10, 10, which will enable a dropping of the hinged bottoms, 3, 3, and thus an emptying of the cylinders of ashes and klinkers. Let the cylinders be filled with pieces of coke or charcoal. At 4, 4 we have charging hoppers. At 5, 5 air-pipes enter the cylinder, and at 6, 6, steam-pipes. At 7, 7 small

FIG. 96.



pipes can introduce coal-tar. The gas passes at 8, 8 into the common pipe 9. Before charging the coke, small wood and shavings are put into the cylinders to start the fire with in one cylinder first. Then compressed air is allowed to enter the through pipe 5, coke is charged until it reaches just below the gas outlet 8, and the valve in the hopper 4, is left open. Under these conditions, a gas mixture passes out from the hopper, which is composed of nitrogen, chiefly


(60 per cent. to 70 per cent.), carbon dioxide and carbon monoxyd ( $N + CO^2 + CO$ ). The sum of  $N + CO^2$  being at the least 75 per cent., it follows that this gas is only a low-grade heat-producer, and is therefore allowed to escape through the hopper. When the top layer of coke has come to bright red heat, the lower layers will be at white heat. The hopper valve is now closed, and the valve at 8 opened. The air valve in 5 is closed, and the steam valve in 6 opened. As the steam impinges upon the white-hot coke, it breaks up into  $H^2 + CO$ . If the valve in the tar pipe *T* be now opened properly, the tar will fall upon the bright red coke and be *dissociated*, *i. e.*, the higher molecules of  $C^nH^n$ ,  $C^nH^{2n}$  and  $C^nH^{2n+2}$  will be broken up into the lower members of the series, into carbon and into marsh-gas. Read again what is said about this under tar and the distillation of coal. All these hydrocarbons will mix with the main mass of hydrogen plus carbon monoxyd, will pass through pipe 9 into the gas holder, or directly to the burners, and yield a fine light, or if more air be allowed to mix with the gas in the burner, a very intense heat, and a non-luminous flame. All this while the white heat of the coke drops down steadily to a red heat and  $H^2O$  can be no longer decomposed. In common language we say that the steam quenches the fire, *i. e.*, extinguishes it. But the second cylinder has been brought up to white heat during the time. We shut off the first cylinder from pipe 9, open the valve 8 and the steam valve, repeating

all the operations as before, while the first cylinder is again blown to white heat—exchanging the steam for air. Thus by means of the two cylinders a perfectly steady stream of water-gas is produced. A battery of three cylinders is more advantageous still, for then we can throw out one after another of the cylinders for the purpose of removing the ashes and klinkers without danger of explosion.

#### FUEL-GAS.

Under this name goes a mixture of gases of inferior grade to the water-gas; cheaper correspondingly, and much in use, at this time, in metallurgical works and also for the feeding of *gas-engines*.


FIG. 97.



*Chemistry.* Steam and air are blown into the coke simultaneously; the air and the steam periods of the water-gas process are thrown into *one* period. One cylinder only is required for the same volume of gas per minute. Let Fig. 97 represent the sec-

tion plan of the cylinder in the level of the steam pipe 6, Fig. 96; let  $S, S_1, S_2$  be the three steam pipes and  $i, i, i$  the parabolic injectors (system of Koerting). Then there will be sucked up by the steam jets a volume of air depending upon the velocity of the steam (pressure) and upon the capacity of the injector. The pressure of the steam must be adjusted to the capacity of the injector so that the exothermic product  $\text{CO}^2 + \text{N}^4$  is larger by about  $\frac{1}{5}$  than the endothermic product  $\text{C} + \text{H}^2\text{O}(\text{CO} + \text{H}^2)$ . The  $\frac{1}{5}$  over energy is consumed by loss of heat through conduction, radiation and fusion of the ashes. The latter, the fusion of the ashes, can not be obtained unless a proper amount of  $\text{CaO}$  be charged with the coal or coke and the percentage of

FIG. 98.



each as well as the composition be known. A lip or outflow must be provided for the slag as shown in Fig. 98, where 1 is the slanting bottom, 2 the dam of the forehearth, 3 a hollow iron beam, through which water circulates to keep open the outflow and prevent the eating away of the firebrick; 4 is the inner level of the liquid slag; 5 is its outer level and when more slag comes it will run over the

dam ; 6 is the twyer or opening for the injector ; 7 is the foundation. If raw soft coal is to be used in this fuel-gas proposition, the cylinder must be provided with a rotating breaker, to prevent the forming of cakes and lumps, which always make poor gas.

#### PETROLEUM, ROCK-OIL, MINERAL OIL.

The term *coal-oil* is most used in the United States. It is not a good name because implying a relationship between coal and this oil. Such a relationship has not been proven in any instance. All the Pennsylvania, Ohio, West Virginia, Indiana oil comes from rocks which lie under the coal measures, very much older in order of formation. *Rock-oil* is a perfectly appropriate term. In regard to the origin of the oil in these rocks, the most plausible opinion is that of Engler who ascribes the oil to enormous numbers of dead fish within the sand, which now forms the oil-carrying strata. Engler obtained oils very similar to coal-oil by subjecting fish to great pressure at temperatures not much above the boiling-point of water.

*Finding of the oil.* Along Oil Creek, Pa., the oil was found but 50 feet under the surface, but for the most part the oil strata are buried deeply. The gushing or flowing of the oil from a fresh hole is due to the accumulated pressure of marsh gas. As this pressure becomes released the automatic flowing ceases, and pumping becomes necessary.

*Physical properties of the oil.* Most oils are colored ; they appear red or brown-red in transmitted



light and opalescent green in reflected light. This double-color or *dichroism* is owing to the presence in the oil of one constituent which has received the proper name: *fluorescin*. The odor is very strong and characteristic. Sometimes the crude oil is quite mobile and sometimes it is thickish, viscous. So also varies the specific gravity from 0.75–0.95. None has yet been found heavier than water. The crude oil is usually quite inflammable.

*Chemical properties.* In general it will hold true to say: Petroleum is a *complex* liquid. The constituting members of the complex are hydrocarbons of the marsh gas or paraffin series ( $C^nH^{2n+2}$ ). The individual members of the series can be separated by fractional distillation many times repeated (see under wood-tar and coal-tar). The members from  $C^4H^{10}$  to  $C^{16}H^{34}$  have actually been prepared by several chemists. The density of the liquid and its boiling-point are the mean of those constants proportionately to the percentages of the constituting hydrocarbons. The heat-value of coal-oil is very high, much higher than that of vegetable and animal fats, because oxygen is absent. Some crude oils, however, contain sulfur compounds. Coal-oil cannot be saponified by  $NaHO$  or  $KHO$ , and by this negative property we distinguish the mineral oil from vegetable or animal oil. (See further on.)

*Refining of crude oil.* The crude oil is stirred together with concentrated sulfuric acid. By this treatment the objectionable sulfur compounds are converted into a solid, resinous body which settles



with the acid, and the cleansed oil is drawn off. The latter is then subjected to the action of steam heat in large iron tank-boilers, and later on to direct fire. Thus are obtained the conventional fractions: 1. *Petroleum ether*, a colorless, very mobile oil of pleasant odor and intoxicating effect, passes over up to  $70^{\circ}$  C. This oil is much used as a solvent for fats and other bodies. 2. *Gasoline* distills between  $70^{\circ}$  C. and  $90^{\circ}$  C. 3. *Benzine* distills over between  $90^{\circ}$  C. and  $150^{\circ}$  C. After this is collected 4. *Kerosene* between  $150^{\circ}$  C. and  $300^{\circ}$  C. This portion is most valued for burning in lamps, because its *flash-point* is high and it is therefore *safe*. 5. *Lubricating oil* distills between  $300^{\circ}$  C. and  $400^{\circ}$  C. 6. *Vaseline* distills next and condenses as a semi-solid in the receiver. 7. Beyond this comes off considerable *paraffin*, a perfect solid (see under tar); a black, spongy residuum remains in the still.

*Determining the flash-point* of a given coal-oil. Pour the oil into a small tin-cup, which stands upon a water bath. With one hand hold a thermometer into the oil, with the other pass a lighted match across the surface of the oil. Note the temperature of the oil at which the vapor above the oil catches fire. This temperature is known as the flash-point and should not be below  $60^{\circ}$  C. for a safe burning oil.

#### NATURAL GAS.

Any bore-hole which is driven into sedimentary rock-formation can be expected to produce gas, provided the strata have not been disturbed much from


their original horizontal condition. *Natural gas* has been found by many analyses to be composed chiefly of marsh-gas ( $\text{CH}_4$ ). If it burns with a sooty flame then there are some of the higher members of the paraffine series present, each as  $\text{C}^2\text{H}^6$ ,  $\text{C}^3\text{H}^8$  . . . Sometimes there is free hydrogen present and in rare instances CO has been found with the marsh-gas.

## CHAPTER XVIII.

### THE HOMOLOGUES OF CELLULOSE—STARCH, DEXTRIN, SUGARS.

So cunning is the work of nature that extraordinary changes are constantly brought about in properties with apparently the same material. The study of a *wheat grain* will illustrate this. Fig. 99 repre-

FIG. 99.



sents the section through the axis of the grain in magnified dimensions. Beginning from the outside there are 3 layers of small, attached cells 1, 2, 3. 1, 2 epicarpium or skin, colorless cells. 3 the endocarpium; these cells contain a yellow coloring matter. 4 the embryo membrane, large cells, containing the nitrogenous body *gluten*, vegetable albumen. 5 a layer of amorphous substance, grey. 6 the meal-kernel made up of large, loose cells, and these cells

are filled with a multitude of white globules which globules constitute what is known as *starch*. 7 is a dark-colored cell, known as the *embryo* or *germ*.

When such a wheat grain is soaked in water or kept wrapt in a wet cloth the grain swells and two sprouts appear at 8, that is, the vital power lying dormant in the protoplasm of the germ cell awakes in presence of water. Cell upon cell is shot outwards until two blades of grass appear above the earth, which gradually develop into a full plant. The material for these cells was drawn from the store of starch and albumen, so cunningly provided within the reach of the embryonic cell within the grain—a veritable fodder-sack.

*The starch, amyllum.* A white, very finely granular substance. Under the microscope we see spheroidal granules built up from concentric layers. Feels soft to the touch. When pure has neither taste nor smell. Is insoluble in cold water. *Hot water* swells the granules until they flow together into a shapeless, transparent paste, starch-paste. The paste dries into a yellowish, hard, horn-like body. The paste is slightly soluble in water, giving a clear solution upon standing. Starch develops in the roots of certain plants, namely, such plants which grow from the root just as well as from a seed—*potato, arrow root, sago*. The seed grain of Indian corn is richer in starch than other seed grains. Hence starch is made from either corn or potatoes. The corn is ground, while the potatoes are cut into a pulp by rotating knives, after a thorough

washing, to remove adhering soil particles. The pulp becomes milky; the milky liquid is strained off through a very fine hair sieve. The sieve retains the cellulose or skin portions, and also the gluten. The milky liquid is allowed to stand, when the white starch will settle to the bottom and is collected after drawing off the water. The starch only needs drying to be at once a commercial product.

*Chemical properties of starch.* (1) The analysis (same as for cellulose) leads to the ratio  $C^6H^{10}O^5$ , that is to say, identical with cellulose. Note nature's trick to give two very different bodies the same composition. Such bodies are named *isomeric* bodies. (2) A solution of iodine in alcohol or in water, solution of KI colors the starch, first purplish-red, and then blue. Heating causes the color to fade, but on cooling the color reappears. Very characteristic reaction for the identification of starch. (3) Starch dissolves in cold, concentrated  $HNO^3$ . The addition of water throws out a white precipitate, a nitro-body similar to gun-cotton, but non-explosive. Boiling very dilute acid (2-3 per cent.) changes the starch into dextrin and finally into glucose. The word starch (stark = strong in German, because starch paste stiffens tissues).

*Dextrin* (from *dexter* = right-handed). A gum-like, syrup-like substance, or dry granular. Much used in place of gum arabic (postage stamps, envelopes), because cheaper and more reliable. It forms from starch, either by heating the dry starch to  $210^\circ C$ . or by moistening the starch with 2 per cent.

$\text{HNO}^3$  solution, and then heating to  $110^\circ \text{C}$ . (best method), or by boiling starch with 2 per cent.  $\text{H}^2\text{SO}^4$  solution until the resultant liquid does not turn blue with iodine solution. Dextrin is  $\text{C}^6\text{H}^{10}\text{O}^5$ , the same as starch and cellulose. Another isomeric form. It is not soluble in alcohol, but easily soluble in water, rotates plane of polarization to right.

*The sugars.* These are bodies formed in the sap of the cells of certain plants; in the seeds (grape, orange, etc.); in the stem (cane, sorghum, sweet corn); in the roots (beets, carrots). All soluble in water; some crystallize, others are syrups; all possess a more or less sweet taste.


*Cane sugar*,  $\text{C}^{12}\text{H}^{22}\text{O}^{11}$ , or  $2\text{C}^6\text{H}^{10}\text{O}^5 + \text{H}^2\text{O}$ . Obtained by neutralizing the somewhat acid sap of the cane, the beet, the sorghum, the maple, by means of chalk ( $\text{CaCO}^3$ ), and by evaporating the clarified liquid rapidly—best in a closed boiling pan from which vapors and air are steadily withdrawn by an air-pump (vacuum pan). From the thick syrup fall small crystals; can be obtained in large, very perfect monoclinic crystals by recrystallization, which are drained from mother liquor in centrifugal filters, and are then known as crude sugar (90–95 per cent.). The refineries remove the 5 or 10 per cent. of impurities. Cane sugar is easily soluble in cold water, 2 parts of sugar in one of water, much more soluble in boiling water, slightly soluble in alcohol, not in ether. Melts at  $160^\circ \text{C}$ ., and on cooling forms a vitreous, glassy body—candy. Heated at  $200^\circ \text{C}$ . it becomes black



sugar or *caramel* (black, horny). By destructive distillation it forms bodies similar to those from starch or cellulose. Cane sugar forms genuine combinations with  $K^2O$ ,  $Na^2O$ ,  $CaO$ ,  $BaO$ ,  $SrO$ , all soluble in water. The combination  $C^{12}H^{22}O^{11}.BaO$  is now much used in the refining process. Concentrated  $H^2SO^4$  acts energetically on sugar,  $SO^2$ ;  $H.CHO^2$  (formic acid) escape, a coaly residue remains. Boiling  $HNO^3 + H^2O$  converts sugar into oxalic acid ( $H^2.C^2O^4$ ), water,  $CO^2$  and  $NO$ . A sugar solution rotates the plane of polarization to the right, like dextrin. If a solution of cane sugar be kept digesting on the water bath with  $HCl$ , or  $H^2SO^4$  (a few per cent.), its right rotation diminishes and turns finally into left rotation; the sugar is *inverted*.

*Glucose, grape-sugar*,  $C^6H^{12}O^6$  or  $C^6H^{10}O^5 + H^2O$ . Is largely contained in honey and in all fruits and berries. Crystallizes in small scaly forms. Less soluble in water than cane sugar (1 glucose— $1\frac{1}{2}$  cold water). More soluble in alcohol than cane sugar. The crystals contain one molecule of  $H^2O$ . Melts at  $70^\circ C$ . Does not taste as strongly as cane sugar. Concentrated  $H^2SO^4$  does *not* change glucose into a coal-like mass; the glucose combines with the acid to form *gluco-sulfuric acid*. Glucose is found in the urine of persons who suffer from *diabetes*. Copper sulfate + glucose + potassium hydrate + water gives at ordinary temperature red or yellow-red cuprous oxyd. Cane sugar gives such a result only by continued boiling. This reaction is the best to distinguish the two sugars from one another.

Large quantities of glucose are manufactured from corn-starch (in U. S.) and from potato starch, (Europe).



Mix 600 grs. of starch with 700 c.c. of luke-warm water into a milk. Mix 100 c.c. of water with 5 c.c. of conc.  $\text{H}^2\text{SO}^4$ . Heat the latter to boiling. Let the milk flow into the boiling liquid, but not so fast as to interrupt the boiling motion. Keep boiling for 3 hours, replenishing the water lost by evaporation. Perfect solution of glucose. Neutralize acid with chalk while boiling. Filter through bone charcoal; evaporate to thick syrup and let cool. In the course of 10 to 12 hours the syrup will change to a stiff mass of small crystals of grape-sugar. Glucose is largely used as a substitute for malt in the brewing of beer, also as an addition to grape juice before fermentation into wine. By heating cellulose with dilute  $\text{H}^2\text{SO}^4$  under pressure in closed boilers, it changes slowly into glucose (saw-dust made into sugar).

*Fruit-sugar, syrup, levulose,  $\text{C}^6\text{H}^{12}\text{O}^6$ .* Gives the slimy consistence to honey. Contained in all berries and fruits with the glucose. Does not crystallize. Rotates plane of polarization to the left. Invert sugar (see above) contains this variety of the family. It combines with 3 molecules of  $\text{CaO}$  to a water-insoluble body, while the isomeric cane sugar gives a soluble compound with  $\text{CaO}$ . The so-called *molasses*

or *melasse* in French, is largely composed of this sugar.

*Milk-sugar, lactose*,  $C^{12}H^{22}O^{11} + H^2O$ . Obtained in hard, colorless, tetragonal crystals, by evaporating the liquid obtained by straining curdled milk. Is only soluble in  $6\frac{1}{2}$  parts of cold water. Agreeable sweet taste, but not as strong as the isomeric cane sugar. Does not melt, but loses the water of crystallization at  $130^{\circ}C$ .

*Gum arabic, arabin*,  $C^{12}H^{20}O^{10} + H^2O$ . Contained in the sap of many plants. It flows from the bark and is found adhering in gum drops to the bark of the stem or the twigs. Specially plentiful in the sap of the acacia trees. Forms with little water a thickish, sticky solution. Not soluble in alcohol. Does not reduce  $CuO$  to  $Cu^2O$  as does dextrin. Furnishes about 3 per cent. of ashes.

## CHAPTER XIX.


### ALCOHOL, SPIRITS OF WINE, ETHYLHYDROXYD, $C^2H^5(HO)$ .

THIS important substance is a derivative of the sugars, and especially of glucose—grape-sugar. The grape-juice is essentially a dilute solution of glucose, has a pleasantly sweet taste. When this solution remains standing, uncovered, in a warm place ( $20^{\circ}$  C. to  $30^{\circ}$  C.), it soon becomes turbid, and gas bubbles arise from the liquid more and more abundantly until the liquid begins to froth. As gradually as this action has increased it decreases, leaving a clear liquid, upon which rests a reddish or brownish scum, while a similar material has gone to the bottom. The liquid has now assumed a sour taste, but very agreeable, and when taken in large quantity produces stupefaction (drunkenness); the liquid is then known by the name of *wine* (Latin = *vinum*).

Arab experimenters were first in trying to get at the understanding of this remarkable action. They succeeded, by distillation, in separating the active principle of wine in form of a colorless liquid, very mobile, and of agreeable odor but burning taste. They found that this liquid will burn with a bright and very hot flame. The name *al-Kohol*, the *breath spirit* (from *spirare*, to breath) was given to this re-

markable liquid. The Latin translator of the Arabic treatise translated al-Kohol into *spiritus vini*, French, *esprit de vin*; German, *weingeist* (the spirit or soul of the wine). However, in recent years the chemists of all nations use the word *alcohol* exclusively.

The process described above, by which a solution of sugar converts itself into a solution of alcohol, is now everywhere known as *fermentation*, the Germans alone use an indigenous word *gaehrung*. But similar processes are known to yield other products than alcohol; hence, to be clearly understood, we say *alcoholic fermentation*. The chemical process of this fermentation is of the simplest.



The gas bubbles forming in the process are  $\text{CO}^2$ . The *ferment* or *yeast* is found in the matter, which causes the turbidity of the fermenting liquid, and which either rises to form the scum or sinks to form the sediment. This matter is biological; it is alive. Under sufficient magnifying power the scum differentiates itself into *cells*, single or in clusters. The cells have an envelope of cellulose, and are filled with a clear liquid. The liquid contains *protoplasm*, which can be separated as white flocculæ. The white substance dries into a horny body, the analysis of which gives  $\text{C} = 55$ ;  $\text{H} = 7.5$ ;  $\text{N} = 14$ ;  $\text{O} + \text{S} = 23.4$ . In this body rests the life-potency of the cell. The cells multiply by "budding," as shown in Fig.

100. A yeast cake is merely a multitude of such cells with some potato starch. Botanists classify these yeast cells as the lowest form of the order *fungi*. The specific name is *saccharomycetes cerevisiæ*. Because such cells are in the air with other minute

FIG. 100.



dust, the fruit juices begin to ferment, apparently by themselves, but if filter paper be tied over the vessel, no fermentation takes place. The addition of yeast merely accelerates the process.

*The manufacture of ethyl-alcohol.* It sometimes pays to distill wine (in France), but only in exceptional cases. The bulk of the alcohol is made from grain or potatoes. The process is essentially identical for all raw materials. It comprises the following operations: (1) *Grinding* of the grain without sifting off the bran. (2) *Mashing* of the grain with *malt*. Under the word *malt* is understood sprouted barley dried after sprouting, and ground. The sprouting of the barley grain develops another mysterious substance—*diastase*—which possesses the power to change starch into glucose more rapidly than sulfuric acid. Its composition is not known, *not* supposed to be a living thing as yeast. 10 parts



of grain, 1 part of malt, 80 parts of water heated by steam in a tub to  $75^{\circ}$  C. until the conversion of the starch is completed; that is mashing.

(3) To cool down the mash to  $18^{\circ}$  C. and transfer it to fermenting vats through a strainer. (4) Addition of sound yeast and fermentation of the *wort* (technical name of the strained mash) in a separate room in which the temperature is kept evenly at  $20^{\circ}$  to  $25^{\circ}$  C. The end of the fermentation is indicated by the collapse of the froth over the liquid.


(5) Distillation of the alcoholic liquor in copper stills. Alcohol boils at  $78^{\circ}$  C., water at  $100^{\circ}$  C. From the boiling mixture more alcohol passes into vapor than water. If the liquor contains 5 per cent. alcohol, then this entire quantity will have gone over when 28 per cent. of the liquid has distilled, leaving 72 per cent. of liquid in the still. (This residue is not thrown away, but used in mixing the feed for fattening cattle.) The distillate contains now 20 per cent. alcohol; is milky from minute drops of amyl-alcohol (fusel-oil), and not saleable. It must be redistilled, yielding a 40 per cent. brandy, and this again distilled to get a 60 per cent. spirit, and this again and again until a *95 per cent. proof spirit* results. Beyond this the water cannot be eliminated by distillation alone.

By mixing with fused  $\text{CaCl}_2$ , or calcined copper vitriol,  $\text{CuSO}_4$ , the water is taken up by these bodies, and another distillation yields *absolute alcohol*,  $\text{C}^2\text{H}^5(\text{HO})$ . The latter is only needed for special chemical purposes. The process of concentrating

the alcohol by distillation is called *rectification*. To save the largest part of the fuel required, many improved processes have been invented, which at last resulted in an apparatus yielding proof spirit at the first operation. This is done by carefully cooling the vapors in an apparatus of large cooling surface, the *dephlegmator*, known also as a *column apparatus*, which stands directly above the still. Fusel oil and other complex side-products are removed from the alcohol in the dephlegmator.

*Properties of alcohol.* A colorless, mobile liquid; remains liquid even at  $-90^{\circ}$  C.; boils at  $78.4^{\circ}$  C. Specific gravity at  $+4^{\circ}$  C. = 0.8095, at  $+15^{\circ}$  C. = 0.795. 100 vols. at  $+4^{\circ}$  C. give 109 vols. at  $78^{\circ}$ . Expands therefore thrice as much as water. Taste, burning. Pure alcohol destroys the organisms, is poisonous; dilute alcohol causes intoxication. It is a solvent for many substances which are not soluble in water—its chief use in the laboratory.

*Chemical constitution.* Analysis gives simply the atomic ratio  $C^2H^6O$ . But we write instead  $(C^2H^5)$  (HO) because alcohol acts towards acids like the hydroxyds of metals. In this last form alcohol is meant to appear as the hydroxyd of the monovalent radical  $(C^2H^5)$ . But the latter may be looked upon as a complex radical made up of the simpler hydrocarbons  $CH^3.CH^2$ , or we may say that ethyl alcohol is equal to two groups of methyl in which one H is represented by the monovalent (HO)



One hydroxyd can take up one hydrogen of an acid, hence when we act upon alcohol with conc.  $\text{H}^2\text{SO}^4$ , we obtain  $2(\text{CH}^3.\text{CH}^2.\text{HO}) + \text{H}^2\text{SO}^4 = (\text{CH}^3.\text{CH}^2)^2.\text{SO}^4 + 2\text{H}^2\text{O} = \text{ethyl sulfate}$  or  $\text{CH}^3.\text{CH}^2.\text{HO} + \text{HCl} = \text{CH}^3.\text{CH}^2.\text{Cl} + \text{H}^2\text{O} = \text{ethyl chlorid}$ . Such bodies were formerly named *ethers*, now they are named *esters*, *i. e.*, salts in which the metal is replaced by a compound carbon radical. When alcohol is heated in air or oxygen, its elements become oxydized, it *burns*, producing much heat.




The absolute heat effect,  $A = 2 \times 12 \times 8240 + 6 \times 34000 = 401760$  heat-units, or *one gram* of alcohol, by burning, will raise the temperature of 97.99 grams of water from  $0^\circ \text{C.}$  to  $100^\circ \text{C.}$ , if no heat be lost by radiation and conduction.

## CHAPTER XX.

### SOME IMPORTANT DERIVATIVES OF ALCOHOL.

*Ether, sulfuric ether, ethyl oxyd, C<sup>4</sup>H<sup>10</sup>.O.* This very important substance originates thus :




Bring into a boiling flask 100 grams concentrated  $\text{H}^2\text{SO}^4$ , 20 grams of water and 50 grams of absolute alcohol. This will give ethyl sulfate. Now close the neck with a three-hole stopper. Pass through one hole a long-stemmed funnel, through the second hole a thermometer down into the liquid ; through the third hole a knee-shaped gas evolution tube, and connect the latter by means of a proper reducer with a condenser. Heat until the thermometer shows  $140^{\circ}\text{C}.$  and maintain this temperature by letting in absolute alcohol. Ether + water will go over steadily and collect in receiver. Shake the distillate with  $\text{Ca}(\text{HO})^2$  + water, *milk of lime*, to neutralize acid particles. Separate the upper stratum of the two liquids with syphon. It contains some water. By redistillation in presence of  $\text{CaO}$  (burnt lime) the ether is obtained pure,  $\text{CaO}$  uniting with the water and binding it.

Ether is a thin, very mobile, colorless liquid, of

strong, penetrating, but agreeable odor. Sp. G. = 0.736. Boils at 35° C., hence very volatile. Mixes with alcohol in all proportions, but does not mix with water. 10 parts of water dissolve 1 part of ether. It dissolves many bodies which are neither soluble in alcohol nor in water, notably fats. It ignites easily and burns with luminous sooty flame; be careful in avoiding open flames in the neighborhood of evaporating ether.



It is evident that ether stands to alcohol in the same relation as potassium or sodium oxyd to potassium or sodium hydroxyd,



After once being separated it does not show any tendency to reconvert into alcohol.

Ether produces unconsciousness, stupefaction, when the vapors are taken into the lungs; it is an anaesthetic.

*Chloroform*,  $\text{CHCl}^3$ , can be considered as marsh gas in which 3H have been replaced by 3Cl. It results when 4 parts alcohol, 3 parts water and 1 part bleaching lime are heated in a flask or retort to the boiling point. With the water is found in the receiver a heavy, oil-like liquid—chloroform. The reaction may go thus:



Colorless, thick liquid. Odor pleasant, taste sweet. Insoluble in water. Specific gravity = 1.48. Boils at  $61^{\circ}$  C. Its vapors are more poisonous than ether; it was formerly much used as an anaesthetic.

*Iodoform*,  $\text{CHI}^3$ . A yellow solid in scaly crystals. Insoluble in water, in acids, in alkalies. Soluble in alcohol and ether. Used much in medicine as an antiseptic for burns and wounds. It originates similarly to chloroform. From a mixture of alcohol, KOH or NaOH and iodine. First forms  $\text{K}(\text{IO}) + \text{KI}$ . Then KIO acts on the alcohol just as  $\text{Ca}(\text{ClO})^2$  does. Work out equation.


*Aldehyde*,  $\text{C}^2\text{H}^3\text{HO}$ . A thin, mobile, colorless liquid. Boils at  $21^{\circ}$  C. and has a suffocating odor. Specific gravity = 0.801. Origin:  $\text{C}^2\text{H}^5\text{HO} + \text{MnO}^2 + \text{H}^2\text{SO}^4 + \text{Aq} = \text{C}^2\text{H}^3\text{HO} + \text{MnSO}^4 + 2\text{H}^2\text{O} + \text{Aq}$ . In presence of air it changes into acetic acid,  $\text{C}^2\text{H}^3\text{HO} + \text{O} = \text{C}^2\text{H}^3\text{O}^2.\text{H}$ . It is therefore a strongly *deoxydizing* body. A glass plate can be made into a silvered mirror by pouring upon the well-cleansed surface a liquid composed of  $\text{AgNO}^3 + \text{NH}^4\text{HO} + \text{C}^2\text{H}^3\text{HO} + \text{water}$ . In this liquid we have  $\text{Ag}^2\text{O}.2\text{NH}^4\text{HO} + \text{NH}^4.\text{NO}^3 + \text{C}^2\text{H}^3\text{HO} + \text{water}$ . Now  $\text{Ag}^2\text{O}.2\text{NH}^4\text{HO} + \text{C}^2\text{H}^3\text{HO} = \text{Ag}^2 + \text{NH}^4.\text{C}^2\text{H}^3\text{O}^2 + \text{NH}^4\text{HO} + \text{H}^2\text{O}$ .

*Chloral*,  $\text{C}^2\text{Cl}^3\text{HO}$ . A colorless, thin liquid. Has a penetrating odor, attacks mucus membrane. Soluble in water. This solution does not give a white precipitate of AgCl when  $\text{AgNO}^3$  solution is added. We explain this by saying that chlorine is *intraradical*, inside of the radical, not in the form of



chlorid. Specific gravity 1.502. Boils at  $94^{\circ}$  C. It is prepared by passing dry chlorine into absolute alcohol until the evolution of HCl stops:  $C^2H^5HO + 8Cl = C^2Cl^3HO + 5HCl$ . When chloral is taken in small doses (dissolved in water), it causes a peculiar intoxication and indifference to pain. For this purpose it is given by physicians, but it can become a dangerous habit for the patient.

*Mercuric fulminate*,  $Hg.C^2N^2O^2$ . Discovered by Howard, A. D. 1800. Liebig and Gay-Lussac found correct composition. *Properties*. Minute, white, needle-shaped crystals. Soft to the touch. Sweetish metallic taste. Little soluble in cold, more in boiling, water. Specific gravity = 4.42. *Very poisonous*. In the dry condition it explodes violently by friction, by a blow, or by concentrated  $H^2SO^4$ .



By experiment 1 gram furnished 78 c.c. of gas with a heat generation of 403.5 cal. or enough to heat the products of combustion of explosion to a temperature of  $4200^{\circ}$  C. Although gun-cotton gives more gas per unit, yet the explosive effect of the fulminate is much greater. This effect is often called by the French expression *brisanee*. We say: Mercuric fulminate is the most *brisant* explosive. Why? Probably because of its instantaneous break-up into gas.

*Preparation*. We dissolve 5 grams of mercury in 60 grams of  $HNO^3$  specific gravity 1.34 (45 c.c.) which gives us  $Hg(NO^3) + HNO^3 + N^2O^3 + NO +$  water. When the metal is dissolved we cool the

liquid to  $70^{\circ}\text{C}$ . We bring 50 grams of 90 per cent. alcohol into a strong, well-tempered,  $\frac{1}{2}$ -liter, round flask and pour the first liquid slowly into the alcohol while rotating the flask. A colorless mixture results. Should no reaction develop at once we place the flask upon a water-bath until small bubbles begin to show. Then we set the flask under strongly drawing hood or out of the window. Usually a very violent reaction sets in with large masses of white fumes emerging from the flask. When the reaction is over a white, flour-like precipitate of the fulminate is found on the bottom of the flask. We cool the liquid to ordinary temperature and more fulminate precipitates. Then we pour off the supernatant liquid, wash the crystals with cold water until the acid reaction ceases, collect the powder on a filter and dry it on the water-bath or better yet, in a current of warm air, or still better we keep it wet.

*Manufacture of percussion caps.* To reduce the brisance and to obtain a more penetrating flame jet, the fulminate is mixed with 30 per cent. of niter for mining caps, or 30 per cent. of potassium chlorate for dynamite igniters. 1. The materials are moistened with 30 per cent. water and mixed with wooden rubbers upon a polished slab of marble. 2. The paste is pressed through hair sieves and thus becomes granulated. The granules are very carefully dried, spread upon paper. The dry granules are sifted through hair sieves to remove the dust particles. 3. The granules are filled into the caps by special machines. The sheet copper is 0.26 mm.

thick. The head has a small cavity Fig. A (Fig. 101). Into this drop granules to the extent of 15 milligrams Fig. B; a stamp presses a thin copper foil over the charge Fig. C, and the cap is ready. This latter is the most dangerous of the operations.

FIG. 101.



Since the charges are small, not much damage can ensue. Heavy charges are fired with larger caps containing 300, 500, 750 mgs. of fulminate charge. Torpedo caps contain as much as 1,500 mgs. of fulminate.



*Silver fulminate*,  $\text{Ag}^2.\text{C}^2\text{N}^2\text{O}^2$ , has similar properties.

## CHAPTER XXI.

### SOME ORGANIC ACIDS.

WE find these acid bodies chiefly in the flesh of berries or fruit, in the free state or in the form of salts—esters. There is a multitude of them and only the most important ones can be mentioned here.

*Formic acid, ant acid, H.CHO<sup>2</sup>.* A colorless, mobile liquid of very pungent acid odor. Solid at  $-1^{\circ}$  C. Boils at  $100^{\circ}$  C. Specific gravity = 1.235. Causes a blister on the skin. Found in pyroligneous acid (wood distillation). Causes the stinging sensation produced both by ants and nettles. It forms in many ways by the oxydation of organic bodies, especially the carbohydrates. Most interesting is the synthetic formation by the action of CO gas upon KOH at  $100^{\circ}$  C. Thus:




And also  $\text{CO}^2 + 2\text{K} + \text{H}^2\text{O} = \text{K.CHO}^2 + \text{KOH}$ . Formic acid precipitates many metals from the solution of their salts in metallic form.  $\text{HgO} + \text{H.CHO}^2 + \text{heat} = \text{Hg} + \text{H}^2\text{O} + \text{CO}^2$ .

*Acetic acid, H.C<sup>2</sup>H<sup>3</sup>O<sup>2</sup> or (CH<sup>3</sup>)COOH,* a colorless, mobile liquid of pleasant, pungent odor. Solid


at  $+15^{\circ}\text{C}$ . (glacial acetic acid), boils at  $109^{\circ}\text{C}$ . Sp. gr. at  $18^{\circ}\text{C}$ . = 1.063. On adding water the specific gravity increases until 1.0748 is reached (acid = 78 per cent., water = 22 per cent.), then the gravity decreases with the addition of water until 1.00 is reached. Hence it follows that one specific gravity may mean two very different acids; for instance, 1.069 can mean an acid of 96 per cent. and an acid of 33 per cent. An acid of 54 per cent. has the same specific gravity as acid of 100 per cent. Glacial acetic acid destroys the skin as readily as oil of vitriol. The vapor of the acetic acid burns. Acetic acid dissolves many oils which are insoluble in water. *The acetates of all metals are easily soluble in water*, hence, lead is usually employed in the form of acetate.

Acetic acid finds not only much use as vinegar, but also in the chemical laboratories and in the chemical arts, such as dyeing and printing. It is the oldest acid known to man, as it forms naturally when sugar solutions stand in a warm place in contact with air. This change does not take place unless a peculiar fungus — *mycoderma aceti* — is present.



Pure acetic acid for laboratory use is prepared by distilling two molecules of crystallized sodium acetate with one molecule of concentrated sulfuric acid in glass retorts; in iron retorts for commercial use.

The latter are furnished with an alembic or helmet made of tin.




The sodium acetate is, at this time, obtained exclusively from pyroligneous acid. (See distillation of wood or cellulose.) The pyroligneous acid is first made into  $\text{Ca}(\text{C}^2\text{H}^3\text{O}^2)^2$ , evaporated to dryness, and the mass gently roasted to destroy tar oils. Thereupon distillation with  $\text{HCl}$  yields fairly pure acetic acid, containing water. This acid is neutralized with  $\text{Na}^2\text{CO}^3$  and the  $\text{Na.C}^2\text{H}^3\text{O}^2$  is allowed to crystallize. The crystals are distilled with  $\text{H}^2\text{SO}^4$ .

*Manufacture of vinegar.* Raw materials are all kinds of alcoholic liquids, especially wine, cider, spoilt beer, and fermented worts from the glucose industry. a. *From wine.* In open barrels, into which wine and vinegar are filled. The vinegar brings with it the ferment. The latter grows rapidly into a thick skin over the whole surface. Vinegar is finished in 8–14 days according to temperature of the room, or the outside air. b. *From dilute alcohol, by the rapid process.* The barrel *B*, Fig. 102, is open at the top. At *F*, *F'* are perforated wooden discs, so-called *false bottoms*, the space between the two discs being filled with clean pine shavings. The latter are soaked with good vinegar, and upon the top layer is sown a culture of pure mycoderma cells (all other germs are excluded). The alcoholic liquid is then distributed by the holes in *F'* over



the surface of the shavings, while plenty of warm air enters through the holes 1, 2, 3, 4, etc. The acidified liquid is drawn from the spigot into the pail, and returned to the top of the barrel as many times as may be needed to convert the whole of the alcohol.

FIG. 102.




It is possible to make the resulting vinegar as strong as 30 per cent., in which form it is known as spirit vinegar, which for cooking purposes is diluted with 10 parts of water.



*Oxalic acid*,  $H^2.C^2O^4$ . At ordinary temperature a white, or colorless solid. Forms readily large crystals, monoclinic with 2 molecules of water of crystallization.  $H^2.C^2O^4 + 2H^2O$ . The crystals show a sharp, sour taste, *no odor*. Soluble in 10 parts of cold water, in three parts of boiling water. Soluble in  $2\frac{1}{2}$  parts of cold alcohol, in 1.8 parts of boiling alcohol. The crystals lose their water of crystallization at  $60^\circ C$ . At  $97^\circ C$ . the crystals melt, at higher heat break up into  $CO^2 + CO$  and  $H.CHO^2$  (formic acid). In a current of air the anhydrous

acid can be sublimed at  $150^{\circ}$  to  $155^{\circ}$  C. The vapors attack the mucous membrane and are poisonous.

*Chemical properties.* Oxalic acid is not a good solvent for the metallic oxyds and metals, because the oxalates are either quite *insoluble* or but little soluble in water. The most insoluble oxalate is calcium oxalate  $\text{Ca.C}^2\text{O}^4 + 2\text{H}^2\text{O}$ , (quadratic crystals). This salt is not soluble in acetic acid but soluble in HCl. Oxalic acid is therefore our best agent for separating calcium from a solution. Oxalic acid decomposes quickly in presence of an oxidizing agent.




It precipitates gold and platinum from their chlorids.



(best way to get pure CO for experiments, the  $\text{CO}^2$  being removed by means of NaHO).

*Occurrence of oxalic acid.* It is found in the saps of many plants, as  $\text{KH.C}^2\text{O}^4$ , especially in the leaves of sour clover (oxalis), and in rumex; also in rhubarb (as  $\text{CaC}^2\text{O}^4$ ). It forms the calculi in the bladder and kidneys of man.


*Manufacture.* There are two ways. (1) By action of nitric acid upon the carbohydrates—sugar, starch, wood,



Theoretically 1 part of sugar requires 2.21 parts of  $\text{HNO}^3$  or 5 parts of 50 per cent. acid. Practice shows that 6 parts are needed to effect complete oxydation. Reaction is energetic after once started. When evolution of gas has gone down, put flask on flame and boil until volume of liquid is  $\frac{1}{3}$  of original. On cooling, the acid crystallizes. (2) By action of  $\text{KOH} + \text{NaOH}$  upon sawdust or similar refuse wood. The action is probably as follows :



The temperature should not be above  $240^\circ \text{C}$ . to obtain the best result. At a higher temperature  $\text{Na}^2\text{C}^2\text{O}^4$  changes to  $\text{Na}^2\text{CO}^3 + \text{CO}$ . When all the wood is changed, let cool. Extract with little cold water, which dissolves only  $\text{K}^2\text{CO}^3$ . Then dissolve residue in boiling water and stir with an equivalent of  $\text{Ca}(\text{HO})^2$ .



Filter. Filtrate is evaporated for next batch. Residue is decomposed by dilute  $\text{H}^2\text{SO}^4$  giving  $\text{CaSO}^4 + \text{H}^2\text{C}^2\text{O}^4$ . Separate the liquid (hot) from the solid  $\text{CaSO}^4.2\text{H}^2\text{O}$  by means of a filter press or centrifugal. Evaporate filtrate until crystals appear, then let cool, and the bulk of the oxalic acid will fall out, and after air-drying can be packed and shipped. In spite of the many operations, this second method

gives the acid at considerably less cost than the first method.

*Lactic acid, milk acid*,  $H^2.(C^3H^4O^3)$ . A colorless, thick liquid. Strong sour taste; no odor when pure. Specific gravity = 1.25. Soluble in water, alcohol, ether. When the water-solution is evaporated a part of the acid escapes with the vapor, but most of it remains as a syrup. At  $130^\circ C$ . it loses one  $H^2O$  and becomes so-called anhydrous acid,  $C^6H^{10}O^5$  (isomeric with glucose, starch).

Lactic acid can be separated from other acids by the insolubility of some of its salts, especially  $Sn.C^3H^4O^3$ . The calcium lactate is somewhat soluble in cold and more soluble in hot water. Lactic acid is found in sour milk; in the gastric juice; in certain fermented vegetables, such as cabbage and turnips. It forms easily from glucose or any other sugar in presence of the *lactic ferment*. The latter is obtained from rotten cheese or rotten meat.

*Tartaric acid, wine acid*,  $H^2.C^4H^4O^6$ . As solid, in colorless, monoclinic crystals. Easily soluble in water, much less in alcohol, and not at all in ether. The water-solution rotates the plane of polarization to the right. Strong sour taste. Melts at  $170^\circ C$ ., forming a vitreous body—meta-tartaric acid—without decomposition. If we add tartaric acid to a solution containing a ferric or aluminic salt, and then an excess of  $NH^4HO$ , there will be no precipitate, enabling the chemist to effect certain separations, otherwise difficult.

*Salt of seignette*,  $Na.K.C^4H^4O^6 + 8H^2O$ . Large crystals showing orthorhombic hemihedry.

*Calcium tartrate*,  $\text{Ca.C}^4\text{H}^4\text{O}^6 + 8\text{H}^2\text{O}$  forms by adding lime water to a tartrate solution. Insoluble in water.

*Tartar emetic*,  $\text{K}(\text{SbO})\text{C}^4\text{H}^4\text{O}^6 + \frac{1}{2}\text{H}^2\text{O}$  soluble in 14 parts of cold water, in less boiling water. Obtained by boiling a solution of ordinary tartar ( $\text{K.H.C}^4\text{H}^4\text{O}^6$ ) with  $\text{Sb}^2\text{O}^3$ , the teroxyd of antimony. Used as an emetic; causes death in larger portions.


We obtain the tartaric acid from the tartar or argol which is the name of the hard, stony deposit in the casks in which the grape juice has been fermenting. The Germans call it *weinstein*, winestone. The crude tartar is a mixture of the acid potassium tartrate  $\text{KH.C}^4\text{H}^4\text{O}^6$  with coloring material and yeast cells, and calcium tartrate. The tartrates were originally dissolved in the juice but had to fall out in measure as the percentage of alcohol increased, they being insoluble in the latter.

We boil the fine powder with 15 parts of water for some time, add bone-black or charcoal which absorbs the coloring matter, and filter boiling hot. From the filtrate precipitate small crystals of  $\text{KH.C}^4\text{H}^4\text{O}^6$ . They go under the name *cream of tartar*. We redissolve in boiling water, add one equivalent of  $\text{Ca}(\text{HO})^2$  and keep boiling until we have  $\text{KOH} +$  insoluble  $\text{Ca.C}^4\text{H}^4\text{O}^6$ . We separate by  $\text{H}^2\text{SO}^4$  after filtering, getting  $\text{CaSO}^4.2\text{H}^2\text{O} + \text{H}^2.\text{C}^4\text{H}^4\text{O}^6$ , and proceed the same as for oxalic acid.

The crude tartar is used in assaying as a deoxydizing agent and alkaline flux, for on heating in a



covered crucible to redness a decomposition takes place,



Both C and CO act as deoxydizers upon metallic oxyds; while  $K^2CO^3$  can take up sulfur or silica, or both.

*Citric acid*,  $H^3.C^6H^5O^7$ . A tribasic acid. Large colorless crystals, orthorhombic combinations  $H^3.C^6H^5O^7 + 2H^2O$ . Very sour taste. Easily soluble in water and in alcohol. The acid acts in iron solutions and aluminum solutions the same as tartaric acid. It is used for a number of medical preparations, especially seidlitz powders, magnesium citrate and others. Contained in sour lemons, in gooseberries.

*Preparation.* The juice is pressed from sour lemons, strained and neutralized with  $Ca(HO)^2$ . The calcium citrate is nearly insoluble in water, and is decomposed like the tartrate and oxalate with  $H^2SO^4$ .

*Malic acid*,  $H^2C^4H^4O^5$ . Imperfect crystals, resembling cauliflower. Deliquesces in the air, forming a syrup. Very sour. Soluble in alcohol. Occurs in many fruits, especially in sour apples, plums, gooseberries, etc. Calcium malate is easily soluble, and forms large crystals. This enables the separation of the acid. Insoluble lead malate is, as a rule, prepared and decomposed with  $H^2S$ .

*Tannic acid, tannin*,  $C^{14}H^{20}O^{18}$ . An amorphous powder; color pale-yellow or brownish; somewhat



shining. Taste very astringent. Very soluble in water, barely soluble in ether, very slightly in absolute alcohol. Its water-solution gives with  $\text{FeCl}^3$  or with  $\text{FeSO}^4$  a blue-black color and later a blue-black precipitate, insoluble in water. Its solution gives a flocculent precipitate with a solution of gelatine. If a piece of fresh skin or hide is placed in the tannin solution, the tannin will gradually leave the liquid and precipitate itself on the skin. The tannin is easily oxydizable, and therefore reduces  $\text{CuO}$  to  $\text{Cu}^2\text{O}$  in presence of  $\text{NaHO}$ , or  $\text{KHO}$ , and gives  $\text{Ag}$  with  $\text{AgNO}^3$ , same as glucose. Hence chemists place the tannin among the *glucosids*. Its chief use in medicine is as an *astringent* in affections of the mucous membrane. It is used to make our ordinary *black ink*. We obtain the tannin from the so-called Turkish nut-galls. The latter form on the leaves and twigs of certain plants, especially oak and acacia, after a sting has punctured the leaf. The sting comes from wasps mostly, who deposit their eggs in the outflowing sap. The sap hardens and forms the nut-like excrescence or growth. The nut-galls contain as much as 65 per cent. of tannin.

*Preparation.* We extract the ground galls with a mixture of 30 volumes of crude ether (0.74), 4 volumes of water, 1 volume of alcohol (90 per cent.). The extract separates into a lower layer, a syrup (tannin and water) forming the upper layer is composed of ether-alcohol with a little tannin and all other bodies. The water-syrup is evaporated and leaves shining amorphous tannin.

*Recipe for black ink.* 125 grams of ground nut galls, 24 grams of copperas ( $\text{FeSO}^4 + 7\text{Aq.}$ ) 24 grams gum-arabic, 1 liter of water, 1 gram of creosote. Boil the materials for 1 hour, replacing the evaporating water. Strain through a hair sieve or through muslin. Let settle for a week, then put up into bottles.


*Tanning bodies* similar, yet not quite identical with tannin are contained in the barks of many trees, especially the oak, the hemlock, the willow. In the tanneries the hides are first cleansed from the hair and adhering parts of flesh, together with the uppermost layer of cells—the cuticle—by scraping. The clean skins are spread in water-tight pits so that a layer of ground bark is between two skins, as many as 100 skins in 1 pit. Then the pit is filled with water and allowed to stand quietly for several months, the longer the better. The water draws the tanning bodies from the bark and the skins take it from the water-solution. By cutting off a bit of the skin one recognizes from the cross section, whether the change has taken place to the very core. Then the skin has become *leather*. It is perfectly proof against putrefaction. It has become impenetrable to water and yet it remains pliable when dried. A similar change takes place in the skin in presence of alum, ferric sulfate, and chromium sulfate. So that we have now tan, iron, chrome, alumina leather. The scarcity of bark has brought into use these substitutes.

*Gallic acid*,  $\text{H.C}^7\text{H}^5\text{O}^5 + \text{H}^2\text{O}$  crystallizes in

silky needles of the triclinic system. Colorless or pale yellow. Soluble in 3 parts of boiling water, slightly soluble in ether or alcohol. Action on metallic solutions in presence of NaHO or KHO or  $\text{NH}^4\text{HO}$  like tannin, somewhat more energetic. Forms well crystallized metallic salts. It is found ready in sumac and other plants. Prepared by allowing a solution of tannin to cover itself with fungi (mildew) in presence of yeast.

*Pyrogallie acid, pyrogallol,  $\text{H.C}^6\text{H}^5\text{O}^3$ .* A fluffy mass of thin scales and needles. Soluble in 2.2 parts of water at  $15^\circ \text{C}$ . Solution colorless.  $\text{FeCl}^3$  gives a blue color with this body; free  $\text{N}^2\text{O}^3$  produces a brown color. All oxydizing agents produce a brown color. An alkaline solution turns brown at once in air. It acts more energetically upon metallic solutions than the preceding ones. This gives its high value as a *developer* in photography. Here the AgBr on the plate becomes changed by the action of light and becomes capable of decomposing water in presence of pyrogallate.  $2(\text{AgBr})$  active +  $\text{H}^2\text{O}$  +  $\text{Na.C}^6\text{H}^5\text{O}^3 = \text{Ag}^2 + 2\text{HBr} + \text{Na.C}^6\text{H}^5\text{O}^4$  (the brown body); the ordinary AgBr is not changed.

*Preparation.* By heating gallic acid thus:



hence name *pyrogallie* from *pyro* = fire.

## CHAPTER XXII.

### THE VEGETABLE FATS AND ANIMAL FATS.

MANY plants develop in their seeds an oily substance, in the place of starch, for the nutrition of the embryo plant; the cotton, flax, hemp, rape, sesame, anise, poppy, mustard, and among the trees all the so-called nut-bearing kinds: palm, walnut, beach, hickory, olive, plum, peach, almond and many others, while some grow oil-bearing roots such as the pea-nut. After removing the hard shell from the nuts the soft kernels are crushed between rolls of iron or stone, the pulp is warmed to about  $60^{\circ}$  C, put into strong hair cloth and pressed by mechanical or hydraulic devices. The oil runs off. The remaining cake is very nutritious, containing still a good deal of oil. It is often fed to the cattle, or used as manure after the oil has been fully extracted with carbon disulfid. Immense quantities of these oils are made annually in Europe; in the United States cotton-seed oil and linseed oil (flax) are the only ones. Most of the plant oils remain thick liquids at ordinary temperature, some become pasty (palm-oil) and a few are solids.

All fats and oils repel water, and are practically insoluble in water. They dissolve in alcohol, in ether and in carbon disulfid. Their specific gravity

is mostly smaller than that of water. Between the fingers they produce a slippery sensation, hence they lubricate, *i. e.*, reduce friction. They taste pleasantly or unpleasantly, and each oil has a different odor or flavor. However, taste and flavor are produced mostly by small quantities of special oils,—flavoring oils—*essential oils*, while the mass of the fat varies but slightly in composition and quality, yet often considerably in the relative quantities of its constituents. Hence each oil has a specific boiling-point, and also a different fluidity—*viscosity*—which is measured by the quantity of the oil which will run through a nozzle of standard diameter in one minute at standard temperature.


*The animal organism* develops fat upon and between the muscles, in the hollow bones, partly as lubricant, partly as a non-conductor of heat, partly as a food store (sick persons lose their fat first, because the body lives upon it when the digestive organs are out of gear). The nerve substance, brain substance is fat, and the yoke of the egg is a yellow liquid fat upon which the embryo of the oviparous animals feeds. Viviparous animals have no eggs, because the embryo cell receives its nourishment from the mother's blood through the navel chord.

The animal fats are solid at ordinary temperature or at any rate pasty. The fat lies in form of globules within a loose tissue of large cells. In heating the fat the cells break, the globules unite into one mass and the cell substance (fibrin, chondrin), coagulates and comes to the surface as brown scum.



Otherwise the animal fats act like the vegetable fats or oils.

*Chemical action of fats.* The fats prove themselves to be *esters* or salts, in which fat acids of the marsh gas series,  $C^nH^{2n}O^2$ , with high value of  $n$ , are united with *glycerin*, which has the properties of a triatomic alcohol.



when  $G$  stands for glycerine and  $\bar{A}$  for a monatomic fat acid radical. This view is deduced from the action of metallic hydroxyds upon the fats. (1)  $NaOH + Aq + \text{fat} + \text{boiling heat}$  yields after sufficient boiling a thick paste, *soap*, which is *completely soluble* in much water, and quite soluble in a few volumes of alcohol. Upon adding  $HCl$  or  $H^2SO^4$  (dilute) to the solution in water, a white flocculent body separates. If the liquid be heated to near boiling point, the white substance melts, rises to the surface and forms there a layer of oil. On cooling the oil becomes solid or semi-solid, and represents the fatty acids. The liquid contains  $NaCl + HCl + \text{water} + \text{glycerin}$ . The *glycerin* was discovered by the Swedish chemist Scheele about 130 years ago. Being an apothecary he had to make adhesive plasters by boiling olive oil or other fats with lead oxyd. In this process the fat acid combines with  $PbO$  into a pasty, sticky body, which is quite insoluble in water. Spreading the paste upon linen, muslin or silk, makes the plaster.



Now Scheele noticed one day that the water, with which he had washed the plaster, possessed a sweetish taste, and this led him to work until he had separated this sweet body in the form of a thick liquid, by evaporation; and because of the sweet taste he named it glycerin from *glucos* = sweet. He knew it was not sugar because it would not give alcohol with yeast. The composition and chemical nature was established long afterwards by J. Liebig and others. Accordingly, we know at present that glycerin has a composition represented by the symbol  $C^3H^8O^3$ ; that towards acids it acts like a base—a hydroxyd, with 3 replaceable hydrogen atoms, and hence we write:  $C^3H^5(HO)^3$  with the probable constitution  $CH^2(HO).CH(HO).CH^2(HO)$ .

*Properties.* At ordinary temperature a syrup. At a very low temperature an orthorhombic solid, which only melts at  $17^\circ$ – $20^\circ$  C. Boils at  $290^\circ$  C., and distills without decomposition; more readily in a current of steam. At  $150^\circ$  C. it ignites on approaching a flame and burns with a pale blue, non-luminous flame. If salts are present the glycerin breaks up during the distillation, thus  $C^3H^8O^3 + \text{heat} = C^3H^4O + 2H^2O$ . The product is *acrolein*, a very disagreeably smelling substance (the odor arising from a glowing wick). Glycerine mixes with water in all proportions, is soluble in alcohol. Not soluble in ether nor in chloroform. Pure glycerine is hygroscopic, attracts water from the air, hence it causes a burning sensation on the mucous membrane. Diluted with 1 volume of water it is pleasant on a chapped

skin because it does not become dry, keeps the new skin from chafing. Chemical reaction: A borax bead moistened with glycerin colors flame green; without glycerin colors flame orange-yellow.

*Preparation.* Boil 90 grams of olive oil with 50 grams of lead oxyd and 50 grams of water until the oil has disappeared, the plaster formed. Then add more water, boil a few minutes, let cool and drain the water through a filter. Pass  $H^2S$  through the liquid to remove any dissolved lead, filter, evaporate below the boiling-point, and at last heat the syrup at  $100^{\circ} C.$  to constant weight.

*Manufacture of glycerin* is a part of the manufacture of stearic, oleic, and palmitic acids. These reactions constitute the base of the process:

1. Saponification as above with hydroxyds.
2. Stearate +  $H^2SO^4$  + water + heat = stearic acid + glycerin + sulfate + water.
3. Stearin + water + heat + pressure = stearic acid + glycerin + water.


The last reaction gives the best results. It is carried on in 2 cylindrical steel-plate boilers. As much as 1000 lbs. of fat can be taken at one operation. The boilers stand one above the other vertically, and communicate by means of 2 tubes so that the materials circulate and the temperature equalizes. The temperature is  $200^{\circ} C.$  and the pressure accordingly 15.3 atmospheres or about 230 lbs. per sq. inch. Time of action 10 hours. The molten fatty acids collect above the watery glycerin, the latter is drawn off at the bottom, evaporated in a vacuum,

and furnishes at once a commercial glycerin. It has a yellow color and contains quite a number of other organic bodies in small quantities. Best purification by means of crystallization at  $+2^{\circ}\text{C}$ . and separation of crystals from mother-liquor by means of centrifugal machine.

*Tri-nitro-glycerin*,  $\text{C}^3\text{H}^5(\text{NO}^3)^3$ , an oily liquid at ordinary temperature, but capable of solidification below the freezing-point. Insoluble in water. Easily soluble in ether. Very explosive by friction or blow. Begins to vaporize at  $75^{\circ}\text{C}$ . The vapors when taken into the lungs cause headache, sick stomach. Tri-nitro-glycerin influences the action of the heart, and is used by physicians in cases of collapse. It has become the most important explosive for blasting since Nobel conceived the idea of letting the oily nitro-glycerin be soaked up by *infusorial earth* (4 parts of the oil, 1 part of the earth). In this form it is known as *dynamite*. The white blasting powder now in use in many mines contains: Nitro-glycerin 45–55 per cent.; soda niter 25–30 per cent.; wood pulp 15–20 per cent.; magnesia 2–3 per cent. Wood pulp takes the place of infusorial earth, and the niter burns up the wood in the explosion. Such a composition is not as effective as dynamite, but on the other hand it can be handled with less danger, and it does not shatter the rock as much as dynamite. Nitro-glycerin dissolves gun-cotton, yielding a jelly-like material known as *blasting gelatine*. The latter can be changed into a hard, horny material, granulated,

and then forms, with the addition of ammonium picrate, the so-called *smokeless powder*.

*Manufacture of nitro-glycerine.* Reaction :



temperature  $10^{\circ}$  C. to  $20^{\circ}$  C. Operation : Prepare a mixture of 30 parts of 95 per cent.  $H^2SO^4$  with 28 parts of fuming  $HNO^3$  (Sp. Gr. 1.48). Cool this mixture to  $10^{\circ}$  C., weigh out 10 parts of glycerin, stir together with 0.3 parts of concentrated  $H^2SO^4$ . Pour the glycerin sulfate in a thin stream, or drop by drop, into the acid mixture while stirring the latter with a thermometer. The temperature rises—should it rise above  $20^{\circ}$  C., stop pouring and cool the liquid, then finish pouring. The result is an emulsion or milky liquid which soon begins to separate into two distinct layers. The upper layer is tri-nitro-glycerin ; the lower layer is sulfuric acid, which retains some nitric acid. After some hours' standing the acid part is discharged from the mixing vessel, and the nitro-glycerin run into cold water, washed several times, and finally with a 5 per cent. solution  $Na^2CO^3$  to remove last traces of acid. There must be no brown fumes during the mixing. Their appearance means danger. The sulfuric acid serves to take up the  $3H^2O$  of the process, so that the  $HNO^3$  retains its concentration. It can be used over and over if the water is boiled out of it.


*Tallow* (from beef or mutton) contains three different fats : 1. *Stearin*,  $C^3H^5.(C^{18}H^{35}O^2)^3$ , *glycerin tri-stearate*. A hard, white crystalline solid at ordinary temperatures ; melts at  $55^{\circ}$  C. then returns

to solid condition, but at  $71.6^{\circ}$  C. is permanently liquid.

2. *Palmitin*,  $C^3H^5.(C^{16}H^{31}O^2)^3$ , *glycerin palmitate*. Scaly crystals; white, luster of mother of pearl. Melts at  $46^{\circ}$  C., then solidifies, and becomes permanently liquid at  $62.8^{\circ}$  C.

3. *Olein* =  $C^3H^5.3(C^{18}H^{33}O^2)$ , *glycerin tri-oleate*. Pure olein, is a colorless oil without either taste or smell. In contact with air it absorbs oxygen and turns "rancid" by separation of "free" oleic acid. Oleic acid belongs to the series  $C^nH^{2n-2}O^2$ .

*Soap, saponification.* Soap results from the change of the glycerin esters into sodium or potassium esters, by the action of NaOH or KOH:



890 parts of stearin require 120 NaOH; 100 parts of stearin require 13.4 parts NaHO and furnish 10.3 parts of glycerin; while the quantity of soap depends upon how far the water is removed by the process of boiling and salting, for in a solution of NaCl soap is more and more insoluble as the percentage of NaCl rises in the liquid. This distinguishes soft from hard soap. The process of changing fat into soap is named *saponification*.

*Drying oils.* The oils from flax seed, poppy seed and some other plants, differ from other oils in so much as they become dry in a day or two when spread over wood or any other solid material. The surface has become covered with an elastic film. A second



and third coating increases the thickness of the film. The latter shows luster and is known as *varnish*. Other oils, such as olive oil, nut oil, etc., do not dry, they turn rancid and continue to be greasy for a long time. Hence linseed oil or poppy seed oil is used as the liquid medium in painting. The oil is mixed intimately with the colored solids oxyds, salts, earths after the latter have been ground to exceedingly fine grain. The chemical reason for this drying lies in the fact that the chief part of drying oils is made up of *linolein* instead of olein. *Linolein* is  $C^3H^5 \cdot (C^{16}H^{27}O^2)^3$ . The linoleic acid is  $H(C^{16}H^{27}O^2)$  and is known as an unsaturated acid belonging to the series  $C^nH^{2n-4}O^2$ . The acid absorbs oxygen rapidly and becomes *oxylinoleic acid*; the latter being the varnish.

*Turpentine, spirits of turpentine, resin, balsam.* All coniferous plants exude from their bark an oily liquid, which soon becomes sticky and ultimately a transparent solid. The pitch-pine of the Southern States and the Canadian balsam fir produce this material more abundantly. By means of cuts in the trees the liquid is collected in the woods (same as maple sap). It is usually a thick liquid of pale yellow color and strong pleasant odor. This turpentine or balsam contains an oil named spirits of turpentine and a solid known as *resin* or *rosin*. The oil is obtained by distillation, usually in a current of steam. The residue contains several similar bodies insoluble in water, but soluble in part in absolute alcohol, and partly insoluble. The Canada



balsam contains oil = 24 per cent.; resin soluble in absolute alcohol = 60 per cent.; insoluble resin = 16 per cent. Spirits of turpentine, a clear, colorless, mobile liquid, sharp taste, peculiar pleasant odor. High refraction. Insoluble in water, little soluble in alcohol, easily soluble in ether, chloroform, carbon disulfid. Boiling-point varies between  $160^{\circ}$  to  $180^{\circ}$  C. Specific gravity = 0.85 to 0.89. Rotates plane of polarization. The substance may be represented as a hydrocarbon,  $C^{10}H^{16}$ , but is evidently a mixture of several hydrocarbons, which have not been separated from each other. Turpentine evaporates very rapidly in air at ordinary temperature and leaves a *film*. Hence it is known by painters as a *quick dryer* when used for diluting paint. Its chief use is in the manufacture of different varnishes, because it dissolves the resins.

*Rosin, pine rosin, colophony.* Obtained from the heating or distilling of the turpentine. A hard, brittle substance, of yellow to brown color, sticks to the mortar and pestle when grinding. Specific gravity 1.01 to 1.08. Insoluble in water. Softens when heated, then melts. Soluble in spirits of turpentine, in absolute alcohol, in ether, etc. Chemically this substance must be considered as an acid—alcoholic solution reddens litmus. This body has been named *abietinic acid*,  $C^{44}H^{62}O^4 + H^2O$ . It combines with the metallic hydroxyds to form soluble and insoluble salts, esters. With KOH or NaOH results soluble (water) potassium or sodium abietinate; with  $Ca(OH)^2$ ,  $Pb(OH)^2$ , etc.,

result insoluble bodies. The acid is, therefore, analogous with the fatty acids. Rosin is frequently added in soap-making.

*Pitch* is the name given to the somewhat elastic body obtained by the destructive distillation of rosin, or resinous wood.

There are a great many other rosin-like substances from other plants, each with peculiar properties. They are all used in making varnishes.

*Rubber, caoutchouc, gutta-percha.* Obtained by evaporation of the milky sap of certain tropical trees, and of immense importance to modern civilization. The gutta-percha tree is from 40–70 feet high. The leaves are inverted, oval and leathery. By means of incisions at different heights the sap is made to flow. It sets or coagulates shortly after leaving the tree and then dries out into a tough, elastic, light-colored, leathery substance which is impermeable to water, therefore even the natives knew how to make flasks and bottles of it. Rubbing causes negative electrification. Very poor conductor. In contact with air it absorbs oxygen, increases in weight, becomes brittle. Under gentle heat the gutta-percha becomes soft and plastic, whence its great serviceability. Melts at  $120^{\circ}\text{C}$ . into a thin liquid; at higher heat forms vapors of a disagreeable odor. The analysis of pure gutta gave C, 86.36; H, 12.15; O, 1.49. The great mass of the gutta is probably a hydrocarbon, the oxygen belonging to a secondary body.  $\text{C}^6\text{H}^{10}$  is probably the nearest approach to the formula. The gutta is somewhat

soluble in absolute alcohol. Easily soluble in chloroform and carbon disulfid.

Under the name *caoutchouc* (Indian) goes a variety of rubber which comes from the milk-sap of a number of trees belonging to different families and quite unlike the gutta plant. Some of the trees are of enormous size, 135 feet high, 25 feet diameter. One tree can produce as much as 60 lbs. of rubber per year without suffering. The fresh sap has an acid reaction. Contains in 100 parts caoutchouc = 31.5 ; albumen = 2.0 ; a bitter substance = 7.0 ; insoluble in water and alcohol = 3.0 ; water = 56.5. Handling of sap : The sap is mostly spread upon boards and dried upon a slow fire ; the resulting film is drawn off, and many films are kneaded into one mass with water. By another method the sap is allowed to stand 24 hours. The rubber rises to the top and looks like cream, (a double volume of water having been first added). The water is let out at the bottom and fresh water is added until it runs off clear. A solution of alum is then added to the cream, and the the caoutchouc separates. It is kneaded to remove the liquids and then dried, but often retains much water when it reaches the factory.

Caoutchouc contains the hydrocarbon  $C^6H^{10}$  in varying proportion depending on the species of plant, upon the local climate, the soil, and especially the treatment of the sap. Physical properties : High elasticity, transparency in thin sheets. Indifference towards acids and alkalies except the concentrated  $H^2SO^4$ ,  $HNO^3$  at elevated temperature. Easy adherence of two fresh surfaces (making of

tubes). Impermeable to water, but permeable to gases. Pure rubber changes on exposure to light and air into a more or less brittle body. Can be restored somewhat by exposure to the vapors of alcohol.

*Vulcanizing of rubber.* The fact that rubber becomes soft and smeary at a relatively low heat reduces its usefulness. This defect can be remedied somewhat by impregnating it with sulfur. In molten sulfur the rubber takes up as much as 15 per cent. S. But it also absorbs sulfur if the latter in the form of fine powder is kneaded together with the fine-cut rubber and then rolled out into sheets or blocks. Thus prepared the material is less permeable to gases and does not get smeary when warmed. Ink eraser is made by adding fine quartz sand to the sulfur. Zinc oxyd, or prepared chalk are kneaded into the rubber to make strong tubes and steam-packing washers. A solution of sulfur in  $\text{CS}_2$  or  $\text{SCl}_2$  in contact with fine-cut rubber gives sulfur to the latter. The actual vulcanization consists in *heating* (baking) the impregnated rubber to a temperature of from  $146^\circ$  to  $170^\circ$  C.

*Hard rubber, ebonite.* A horn-like, usually black substance in which one does not recognize any of the usual properties of rubber. Replaces wood, glass, leather, horn, ivory for many articles, notably combs. Can be pressed to assume almost any shape. Excellent non-conductor for electricity. 6 parts of rubber are kneaded with 3 parts of sulfur, 1 part of zinc sulfid or other material, then heated for several several hours to  $140^\circ$  C., and pressed in iron moulds.

## CHAPTER XXIII.

### ORGANIC ALKALOIDS.

THESE bodies are of special interest as strong stimulants or direct and fatal poisons. Combine with acids to form easily crystallizing salts nearly all soluble in water. Some turn red litmus to blue. They are the direct products of certain plants, in which they are found as salts of organic acids.

*Theobromin*,  $C^7H^8N^4O^2$ . The stimulating substance in cocoa and hence in chocolate. Is very slightly soluble in water, its reaction is neutral.

*Caffein*,  $C^8H^{10}N^4O^2$ . The stimulant in coffee and in tea. The tea-leaves contain as much as 4 per cent.; the coffee bean only one per cent. It is not a strong base; the salts are decomposed by water.

*Urea*,  $CH^4N^2O$ . Colorless, orthorhombic prisms. Specific gravity 1.45. Taste cooling like niter. Easily soluble in water. Reaction neutral. Combines with acids to salts, which are mostly little soluble in water, especially the precipitates produced by  $AgNO^3$  and  $Hg(NO^3)^2$ . Urea is the principal substance contained in urine.

*Morphin*,  $C^{17}H^{19}NO^3$ . White or colorless; little soluble in water, but easily soluble in acids. Is a narcotic or sleep producer; overdose is fatal. Contained in opium, and opium itself is the result from



drying the sap which runs from unripe seed capsules of the poppy.

*Chinin or Quinin*,  $C^{20}H^{24}N^2O^2$ . White solid, slightly soluble in water with blueing of litmus; strong base. Easily soluble in acids—chinin sulfate forms white, fluffy needles, and is usually taken as a remedy for malaria or other fevers. Chinin is extracted from the bark of a shrub growing in Peru.

*Strychnin*,  $C^{27}H^{22}N^2O^2$ . Colorless prisms, nearly insoluble in water and only slightly in alcohol or ether. Soluble in acids. The sulfate is used in medicine in minute doses. 50 milligrams of the sulfate constitute a *fatal dose*, inasmuch as the strychnin causes terrible convulsions and lock jaw. It is developed in the fruit and seeds of the strychnin plant.

*Brucin*,  $C^{23}H^{26}N^2O^4 + 8H^2O$ . Colorless tetragonal prisms. Easily soluble in alcohol and ether, slightly soluble in water. Reaction: Brucin +  $HNO^3$  gives *red body*, which turns into a purple precipitate on adding  $SnCl^2$ . Made use of to detect nitrates. Brucin is found with strychnin in the beans of the *Strychnos Ignatii*. Brucin is poisonous but not so violently as strychnin.

*Atropin*,  $C^{17}H^{23}NO^3$ . Needle-shaped prisms, silky luster. Tastes very sharp and bitter. Salts do not crystallize readily. Is very poisonous. When applied in dilute solution to the cornea of the eye the pupil becomes enlarged, so that the iris seems to disappear. It is extracted from the cherry-like fruit of *atropa belladonna*, (belladonna signifies handsome



lady—because the ladies of fashion have made use of the extract to make their eyes more fascinating).

*Cocain*,  $C^{17}H^{31}NO^4$ . Colorless prisms soluble in alcohol. Is extracted from the coca leaves. Produces local stupefaction of the nerves and is therefore used to relieve pain.

*Nicotin*,  $C^{10}H^{14}N^2$ . A colorless oily liquid, usually yellow because not quite pure. Odor penetrating, disagreeable; soluble in water, in alcohol. Changes red litmus to blue. A strong base. Boils at  $250^{\circ}C$ . Very poisonous. A small dose acts as a narcotic. Extracted from the leaves of tobacco. Fine Havana tobacco contains only about 2 per cent. Coarse common tobacco as much as 7 per cent. A poultice of tobacco leaves on any part of the body can cause convulsions and even death. The smoking tobacco has been fermented, most of the nicotin is therefore destroyed and  $NH^3$  is formed.




## CHAPTER XXIV.

### ALBUMEN AND ALBUMENOIDS.

The animal body is composed, from the chemical standpoint, of fat, muscles, bone and derivatives of the muscles. The muscles or flesh, as well as the skin, hair, tendons, are composed of albumenoids; the bone is calcium phosphate.

*Albumen*,  $C^{72}H^{112}N^{18}S^2O^{22}$ . Is obtained by evaporating the white of eggs at a temperature not exceeding  $50^{\circ}$  C., and thus,



obtained represents a yellowish transparent mass with a specific gravity of 1.314. With water it swells and then dissolves. It has an alkaline reaction (due to the presence of KOH and K salts). If the water solution be heated it begins to become turbid at  $60^{\circ}$  C. At  $75^{\circ}$  C. large flakes separate; the albumen is *coagulated*, and a faint odor of  $H^2S$  noticed. The addition of  $HCl$ ,  $H^2SO^4$ ,  $HNO^3$  causes the coagulation without heating; the addition of KOH or NaOH prevents coagulation, even at boiling heat.

Basic lead acetate,  $(HO)Pb.\bar{A}^2$ , forms a precipitate with albumen. The percentage composition of dry albumen is: C = 53.4, H = 7.0, N = 15.6, O = 22.4, S = 1.6. The blood of all animals contains a

colorless substance which acts like the white of eggs, and a similar body is contained in the sap of all plants: they are known as *blood albumen* and *vegetable albumen*. What biologists call protoplasm is chemically not distinguished from albumen, but protoplasm has life and albumen has not. To find the conditions under which ordinary albumen will change into protoplasm will be equivalent to finding the source of life itself. A very grand problem, and one which I think possible of solution, though not probable. \*

All derivatives and homologues of albumen we call *albumenoids*.

*Haemoglobin*,  $C = 53.8$ ,  $H = 7.3$ ,  $N = 16.1$ ,  $S = 0.5$ ,  $Fe = 0.4$ ,  $O = 21.9$ . This body gives the red color to blood. Under the microscope a drop of blood reveals a colorless fluid in which spheroid red bodies float freely, together with pale bodies of the same shape. The first are known as the red blood corpuscles, the second as the white corpuscles or *leucophytes*. Diameter about 0.013 mm. The bulk of the red corpuscles is water and haemoglobin. Note the iron in the composition. The red color is due to this iron. Haemoglobin absorbs oxygen, nitric oxyd, carbon monoxyd, hydrogen cyanid. The reason for the very fatal action of CO and HCN upon man is probably due to this property of the haemoglobin.

*Haematin*,  $C^{48}H^{51}Fe^3N^6O^9$ . This is a genuine chemical unit. Blue-black of color, insoluble in water, alcohol, ether. Soluble in solutions of KOH

NaOH,  $\text{NH}^4\text{HO}$ . Also in dil.  $\text{H}^2\text{SO}^4$ . The alkaline solution is olive green in thin column and red in thick column. Obtained from haemoglobin.

*Fibrin*, C = 52.6, H = 7.0, N = 17.4, O = 21.8, S = 1.2. Fibrin is held in solution in the fresh blood. When blood is in contact with air for a short time it *clots*, becomes thick. Beating with a paddle separates from the blood a red, stringy, fine fibrous mass, which is fibrin mixed with haemoglobin. The clear liquid is named the *serum* of the blood, and contains the blood albumen and many other bodies besides the mineral salts. The muscles are chiefly modified fibrin, so called *hyssin*, *creatinin*, *sarkin*, *xanthine*, all albumenoids.

*Casein* (*cheese stuff*), C = 53.6, H = 7.1, N = 15.7, O = 22.6, S = 1.0. This body is found in the skimmed milk together with milk-sugar and salts. Casein is the chief nutriment in milk. It is well known that milk *curdles* and then has a sour reaction. The curds are composed of casein and water; the water can be removed by pressing and drying. Casein turns into cheese by the action of ferments.

*Horn*, *hair*, *skin* are all closely allied to albumen, although their chemical action is quite different, for these bodies are quite insoluble in water and in all agents except concentrated alkalies, which dissolve them and evolve  $\text{NH}^3$  as other albuminoids.

	<i>hair</i>	<i>horn</i>	<i>finger-nail</i>	<i>wool</i>
C	= 49.8	50.7	50.2	49.8
H	= 6.4	6.7	6.8	7.0
N	= 17.1	17.3	16.9	17.7
O + S	= 26.7	25.3	26.1	25.5

Boiling water changes the skin into *gelatine* or *glue*. *Silk* is partly soluble in boiling water, which extracts *sericin*,  $C^{15}H^{25}N^5O^8$ , and leaves *fibroin*,  $C^{15}H^{23}N^5O^6$ .

*Distillation of albumenoid substances. Animal charcoal.* When albumenoid bodies, more particularly the hoofs and horns, and leather scraps of cattle are subjected to dry distillation, the succession of phenomena and the products are in general similar to those which we observed in the distillation of wood and coal, especially the latter. Yet the quality and quantity of the products are different.

The distillation furnishes (1) a black residue—animal charcoal; (2) a watery ammoniacal liquid, and oils (not tar); (3) gases of exceedingly strong and disagreeable odor. Animal charcoal contains more or less of the original nitrogen according to the degree of the temperature. Much of the nitrogen goes over as  $NH^3.CO^2$  (carbamid) and  $(NH^4)^2CO^3$ . (This used to be the chief source of ammonium salts, hence the expression salts of and spirits of hartshorn).

All charcoal has the faculty of attracting coloring matter which may be either actually dissolved or only suspended in liquids; but animal charcoal, and

especially bone charcoal (bone-black), are more efficient than wood charcoal. The oil resulting from this work is known as the *oil of Dippel*, or also *neat's foot oil* much in use formerly as a lubricant for clocks and watches and other delicate mechanism. It does not turn rancid, nor change into resinous sticky substances as other oils do.




## CHAPTER XXV.

### ORIGIN OF CYANIDS.

WHEN potash (pearlash) is liquefied in an iron crucible or pot at a bright red heat and animal charcoal (except bone-black) be added, the charcoal disappears amid turbulent evolution of gas. (In practice the proportions of potash and charcoal vary, perhaps 1 : 1 is a good average.) If then the fused mass be digested with water, or boiled with the latter in the kettle or iron pot, and then allowed to stand quietly, there will be found, after several days, a crop of yellow quadratic crystals and a mother liquor containing  $K^2CO^3$ ,  $K^2S$ , and some other bodies. This liquor, after evaporating to dryness, can be used in the next operation with fresh potash. The crystals are of exceeding interest. In the drug trade they are known as *yellow prussiate* of potash. The German chemist Diesbach, of Berlin, was the first who discovered that this salt gives with certain salts of iron a beautiful blue precipitate—known as *prussian blue*—but kept his knowledge as a secret; the Englishman Woodward, in 1724, published the first account of its preparation. In 1752, the Frenchman Nacquer obtained the pure salt, the *phlogisticated alkali*, but only through the genius and labors of Scheele (1732), of Gay-Lussac and Porret (1814),


of Gmelin (1822), and lastly of Liebig, came to light the true nature of this remarkable substance, which chemists now call unanimously potassium *ferrocyanate* and give to it the symbol  $K^4(Fe(CN)^6)$ . The symbol assumes the existence of a radical  $(Fe(CN)^6)$  of the valence four. Within this larger radical there are six units of the smaller radical  $(CN) = \text{cyanogen}$ . The larger radical is ferro-cyanogen, which some chemists write  $FeCy$  and therefore the yellow crystals are  $K^4FeCy + 3H^2O$ . One sees here the very positive iron become a part of a very negative complex radical, in such a way that from a solution of this salt the alkaline hydroxyds do *not* precipitate any iron hydroxyd, nor does  $H^2S$  precipitate any  $FeS$  when passed into the alkaline solution. But for an understanding of this strange complex it will be necessary to break down the complex form in order to get at the fundamental unit cyanogen.

*Properties of potassium ferrocyanate.* Commonly the crystals show a light lemon-yellow color; color is sometimes orange. The crystals are very cleavable parallel to the basal plane and the cleavage plates are pliable, elastic. The salt dissolves in 4 parts of water at ordinary temperature, in 2 parts at boiling temperature. It loses its water of crystallization between  $100^\circ$  and  $110^\circ$  C. and becomes an opaque, chalk-like mass. At red heat it melts and decomposes with evolution of nitrogen gas, thus:




The fused mass is grey. Water extracts a colorless solution of  $K(CN)$ , *potassium cyanid*, leaving behind a black mixture of iron and carbon and iron carbid.

*Potassium cyanid*,  $K(CN)$ . C = 18.4, N = 21.5, K = 60.1. It is a white crystalline body. Easily soluble in water. Insoluble in 95 per cent. alcohol; but in warm 50 per cent. alcohol it is fairly soluble. On cooling the solution will deposit isometric cubes or cubo-octohedrons similar to KCl or NaCl. The salt itself as well as the solutions emit the strong and peculiar odor of bitter almonds, because the  $CO^2$  of air decomposes the solution (and the solid salt in moist air also) thus:



The odor is due to  $H(CN)$  = hydrogen cyanid. The solution of  $K(CN)$  in water does not bear heating, and therefore cannot be concentrated by boiling as other salt solutions, because the salt breaks up thus:



That is, potassium cyanid breaks up into potassium formate and ammonia. This behavior should be well remembered.


*Preparation.* Pulverize the yellow ferrocyanate coarsely. Heat in a flat, iron dish or pan to expel water of crystallization. The salt turns white. Fill it into an iron crucible, which stands in a furnace, after admixing dry pearlash in the ratio of 3 parts of pearlash to 8 parts of the dehydrated ferrocyanate. Cover with an iron lid and heat grad-

ually to yellow heat. The  $K^2CO^3$  is added in order to save all the cyanid, thus :



$K(CNO)$  is potassium *cyanate*. The product is thus a mixture of 5 molecules potassium cyanid with 1 molecule potassium cyanate. The cyanate is not harmful for any of the ordinary uses of the cyanid. In order to obtain pure  $KCN$ , the pearlash is left out. When the mass in the crucible fuses quietly, introduce a warm glass rod, and this rod, upon cooling, must be covered with a pure white film of the salts. Tap the crucible several times on the floor to cause a perfect settling of the iron particles and pour out into ingot moulds.


*Chemically pure KCN* can only be obtained by passing  $H(CN)$  into a solution of  $KOH$  in alcohol, the liquid being kept cool. Then  $K(CN)$  separates in small crystals. Collecting these upon a filter they can be melted into solid cakes of desired shape and size.  $K(CN)$  is *very poisonous*. It is used to extract gold from the ores, because  $Au$  dissolves, forming a double cyanid, but only in presence of air thus :



A very dilute solution such as 0.2 per cent.  $KCN$  is just as efficient as a strong solution. Silver dissolves similarly and all silver salts except  $Ag^2S$  and even the latter slowly in presence of air. Hence the


use of KCN in cleaning silverware, in fixing photographic negatives.  $K(CN)$  is used as a solvent in electroplating with silver, with gold, and other metals. It is used for separating nickel from cobalt.

*Hydrogen cyanid, prussic acid,  $H(CN)$* , is at ordinary temperature a colorless liquid. Strong odor of bitter almonds or peach kernels. Boiling-point at  $26.5^{\circ} C.$ , hence exceedingly volatile. One drop upon a glass slide solidifies from the sinking of the temperature due to the rapid evaporation. Point of solidification lies at  $-15^{\circ} C.$  Specific gravity = 0.696.  $H(CN)$  barely reddens blue litmus. This substance is so very dangerous that it should only be handled by very expert persons. Mixes with water and with alcohol in all proportions. The water solution exposed to light decomposes into a brown solid and  $NH^3$ , hence such solution should not be kept, but should be prepared whenever needed. The vapor of  $H(CN)$  burns with a *purple* flame.  $H(CN)$  is changed by concentrated  $HCl$  or  $H^2SO^4$  into formic acid  $H(CHO^2)$  and ammonium salt thus :




(Some water must be present).

That  $(CN)$  is a monad and combines with *one*  $H$  can be shown by decomposing one volume of the vapor or gas over mercury by  $K$ , when  $\frac{1}{2}$  vol. of hydrogen results. The water solution of  $H(CN)$  dissolves oxyds and hydroxids, forming metallic cyanids :



Mercuric cyanid which is soluble in water.


*Preparation of H(CN).* The reaction  $\text{HCl} + \text{K}(\text{CN})$  is not available, because the solution of  $\text{K}(\text{CN})$  does not stand heating. (See above.) But the yellow prussiate is available for this purpose, with the following scheme :



*Recipe.* Bring into a short-neck flask 30 grams pulverized yellow prussiate and a cooled mixture of 21 grams conc.  $\text{H}^2\text{SO}^4$  with 42 c.c. of water. Connect with Liebig's condenser and let the end of the condenser tube dip just under the distilled water (25 c.c.) in the receiver. Then distill.

*Cyanogen,  $\text{C}^2\text{N}^2$ .* (*Kuaneos = Cy.*). This radical is at ordinary temperature a colorless gas of peculiar odor and very poisonous. The gas shows a sp. gr. of 1.806. It is inflammable and burns with *purple* flame. Under a pressure of 3 atmospheres it becomes a mobile liquid which turns into a mass of crystals at  $-35^\circ \text{C}$ . It is soluble in water and in alcohol.

*Preparation.* We heat in a small retort, or in a glass tube, mercuric cyanid (see above), the latter breaking up :



*Composition.* We collect the gas over mercury ; add somewhat more than 2 volumes of O and  $\frac{1}{2}$  volume ( $\text{H}^2 + \text{O}$ ) fulminating gas, and explode. We



introduce KOH to absorb  $\text{CO}^2$ . Next we introduce hydrogen, 2 volumes, and explode after reading the volume. From the contraction we find the remnant of O after first explosion ( $\frac{1}{3}$  contraction and  $\frac{2}{3}$  contraction for H consumed), and subtracting this from the remnant of N + H we get at the vol. of N, and in the absorbed vol. of  $\text{CO}^2$  there is  $\frac{1}{2}$  vol. of C. And then we find that 1 vol. cyanogen gives 1 vol. C + 1 vol. N, hence cyanogen in the free state is  $(\text{C}^2\text{N}^2)$  not (CN). It follows also that here is an exception to the law which we found previously. Here one volume of an element uniting with one volume of another element yields *two* volumes of the combination. In cyanogen the *two* volumes give only *one volume* of the product. This is verified by the volume weight of the gas, for.

1 volume of C weighs	1.072 grs.
1 volume of N weighs	1.250 grs.
<hr/>	
1 volume of $\text{C}^2\text{N}^2$ weighs	2.322 grs.,

while the experiment gives 2.326 for one volume of cyanogen. Cyanogen combines with chlorine to form  $(\text{CN})\text{Cl}$ , a terribly poisonous gas. It also combines with bromine.

*Sulfocyanogen*,  $(\text{CNS})^2$ , forms colorless crystals.

*Hydrogen sulfocyanate*,  $\text{H}(\text{CNS})$ , a colorless liquid of pungent odor like HCl and strong acid reaction.


*Potassium sulfocyanate*,  $\text{K}(\text{CNS})$ . Colorless crystals; easily soluble in water and in alcohol. The solution of this salt gives with ferric salts a *blood-*

*red solution* in presence of free acid. Very delicate reaction.

*Preparation.* Fuse together KCN + S in a crucible, dissolve in water and crystallize.


*Potassium ferricyanate, red prussiate of potash*,  $K^6(Fe^2(CN)^{12})$ . A beautiful salt. The crystals are of deep garnet-red color and dissolve very easily, with an intense yellow color, in water. The crystals contain no water of crystallization.

It is prepared by acting upon the solution of potassium ferrocyanate with chlorine; thus:




We say that 2 molecules of tetravalent ferrocyanogen have become polymerized—have coalesced—in one hexavalent radical, *ferricyanogen*,  $(Fe^2(CN)^{12})$ .

In qualitative analysis potass. ferricyanate serves to reveal the presence of a *ferrous* salt inasmuch as they combine to an intensely blue compound, thus:




$Fe^3(Fe^2(CN)^{12})$  is known as a blue coloring material under the name of *Turnbull's blue*.

Potassium ferrocyanate gives with *ferric* salts a similar combination of equal intense blue color which is named *prussian blue*. The following equation represents this relation:





Of prussian blue as well as of Turnbull's blue there is a water-soluble variety, which results whenever an excess of the potassium ferro- or potassium ferricyanate is added to a ferric or ferrous salt. This soluble form is known as *wash blue* in the laundry business.


*Potassium cyanate*,  $K(CNO)$ . This salt results when  $K(CN)$  is kept liquid in presence of air.



This strong affinity for oxygen is the reason why  $K(CN)$  is a most excellent agent for deoxydation at high heat. It is much used in assaying and in blowpipe work thus:



The aqueous solution of potassium cyanate breaks up even at ordinary temperature; when heated to boiling  $NH^3 + CO^2$  escape and  $K^2CO^3$  remains in the liquid.



*Hydrogen cyanate* can be made, but is exceedingly unstable. It has an odor somewhat like a mixture of  $SO^2 + HC^2H^3O^2$ . It is very curious that if a solution of  $(NH^4)(CNO)$ , ammonium cyanate is evaporated, only water escapes, but the residue has all the properties of *urea*,  $(CH^4N^2O)$ , the principle secretion in the animal and human urine.

*Cyanuric acid*,  $H^3(C^3N^3O^3)$  can be made in large colorless crystals. One recognizes that this is the formula of cyanic acid multiplied by 3. It results

from the heating of dry urea above the boiling-point, when  $\text{NH}^3$  escapes and cyanuric acid is left.

*Fulminic acid* is not known in the free state. But we saw above that the analysis of the mercuric fulminate gives  $\text{HgC}^2\text{N}^2\text{O}^2$ . Now this radical ( $\text{C}^2\text{N}^2\text{O}^2$ ) is just the double of the radical in (CNO) cyanic acid. Hence we have here an intensely interesting polymerism or equal percentage-composition for three totally different substances.

## CHAPTER XXVI.

### BONE-ASH AND PHOSPHORUS.

THE animal skeleton is composed of bones. The bone again can be separated into a mineral part (not combustible), into gelatine (glue) and into fat. The fat may be extracted by any of its solvents (carbon disulfid, ether, chloroform, etc.) The gelatine can be boiled out by water. If the bone after extraction is merely dried and bleached it becomes fit for conversion into useful and ornamental objects (by the lathe and by the carving tool). If it be thrown into a hot furnace it will be converted into *bone-ash*. With the latter and its chemical nature, we shall now occupy ourselves.


Bone-ash is infusible and somewhat luminous at a high temperature. It dissolves in  $\text{HCl}$ , in  $\text{HNO}^3 + \text{aq.}$ , but not in  $\text{H}^2\text{SO}^4$ . There is usually a disengagement of  $\text{CO}^2$  unless the burning of the bones had been done at very high heat. In the latter instance a small quantity of the ash will produce a brown spot (alkaline reaction) when placed upon a strip of yellow turmeric paper. It acts thus like  $\text{CaO}$ . The solution in  $\text{HCl}$  gives a white, gelatinous precipitate when  $\text{NH}^4(\text{HO})$  is added; therefore there must be an *acid* present of as yet unknown properties, because the solution of calcite in  $\text{HCl}$

or of  $\text{CaSO}_4$  in  $\text{HCl}$  does not precipitate with  $(\text{NH}_4)\text{HO}$ . The presence of calcium is, however, made certain by the form of the minute crystals which are separated upon adding conc.  $\text{H}_2\text{SO}_4$  to the  $\text{HCl}$  solution of the bone-ash. These crystals are identical in shape with the calcium sulfate or vitriol.

But since we have learned that oxalic acid forms with calcium an oxalate which is not soluble in water and not soluble in dilute acetic acid, we can apply that knowledge right here to advantage. For we find the precipitate which is thrown down by  $\text{NH}_4\text{HO}$  in an  $\text{HCl}$  solution of the bone-ash to be quite soluble in acetic acid. Adding a solution of ammonium oxalate to the latter a bulky white precipitate falls and thus calcium in large quantities is proved beyond any doubt. Let us designate the unknown acid as  $\text{A}^x$ ; then we can write a preliminary scheme for what we have thus far accomplished.

Bone-ash =  $\text{CaA}^x$ ; solution of bone-ash in  $(m + 2)\text{HCl} = \text{CaCl}^2 + \text{H}^2\text{A}^x + m\text{HCl}$ ; the precipitate by  $\text{NH}_4\text{HO}$  must be  $\text{CaA}^x$ .

The acetic solution will be again




To this solution we add ammonium oxalate  $(\text{NH}_4)^2\text{C}^2\text{O}^4$ , and thus we get a precipitate and a liquid. The precipitate will be  $\text{Ca}(\text{C}^2\text{O}^4)$ , the liquid will contain  $\text{H}^2\text{A}^x + \text{NH}_4\text{Cl} + \text{NH}_4(\text{C}^2\text{H}^3\text{O}^2) + \text{H}.\text{C}^2\text{H}^3\text{O}^2 + (\text{NH}_4)^2\text{C}^2\text{O}^4$  water. We filter and



evaporate to dryness. Then we heat carefully over an open flame, causing the volatilization of  $\text{NH}^4\text{Cl}$  and of  $\text{NH}^4(\text{C}^2\text{H}^3\text{O}^2)$ ; further increase of heat will eliminate the excess of  $(\text{NH}^4)^2\bar{\text{O}}$  and we will have, probably,  $\text{H}^2\text{A}^x$ , the unknown acid, or perhaps  $(\text{NH}^4)^2\text{A}^x$ ; provided that this unknown acid does not easily volatilize. We find that a remnant is left; that this remnant imparts a *green* color to the flame, and that it dissolves in water, giving a colorless solution which shows a strong acid reaction. We also find that the remnant does volatilize at a *red heat*. Thus we have established that this body is probably an acid-forming, non-metallic, oxyd. It is not probable that any of the non-metallic elements of our acquaintance could produce such a residue, since their oxyds are very easily volatilized, or since they form hydrogen compounds which are quite volatile. In order to separate the element a treatment with strong deoxydizing bodies is indicated, to wit, potassium, sodium, hydrogen, carbon, marsh gas. Of these possible agents *carbon* is the only available. (Reasons below.) We make  $\text{H}^2\text{A}^x$  into a thick syrup by using  $\text{H}^2\text{SO}^4$  to precipitate the calcium instead of using  $\text{NH}^4\text{HO}$  and  $(\text{NH}^4)^2\bar{\text{O}}$ , and then evaporating. Mix charcoal powder with it until a sticky mass results. Heat over open flame until quite dry, and then fill the black mass into a tube, F, Fig. 103, of hard infusible glass, which has been closed at one end. By placing the tube upon a brick (1) against a second brick (2) and using a Bunsen blowpipe on the closed end, we can bring

the tube to the required temperature. As the heat rises we begin to observe a peculiar odor at the mouth of the tube, and on approaching a taper a greenish flame develops, a white smoke rising into the air. At the same time wax-like drops condense in the forward part of the tube at (3). At length the flame burns more and more with a pure

FIG. 103.



blue color (CO) and the action is complete. When the tube has become cold we cut off the forward part with a file stroke and a red-hot rod (glass or iron). The new substance is transparent or translucent. Its color varies from pale yellow to bright red. (Probably 2 different substances?) It is soft as wax. It emits a peculiar odor similar to that of *ozone*. A white fume arises from it steadily. It melts at  $45^{\circ}$  C., and begins at once to burn with a bright flame and evolution of white fumes. Under water it can be melted without danger of ignition. It dissolves somewhat in alcohol, more in fat oils. If such solution is rubbed over the hands or the face those parts will shine in a dark room with a pale green light. This property has been the justification for giving to this remarkable elementary body

the name *phosphorus* (*phos* = light ; *phorus* = carrier). It would be more consistent to change the name to *phosgen*, and thus obtain a consonant series : Oxygen, hydrogen, nitrogen, chlorogen, bromogen, iodogen, fluogen, phosgen, and so forth.

(1). *Note.* Brand was the first to obtain phosphorus, in 1674, by distilling, in a clay retort, the residual mass from evaporated urine ; but that bones contain much more phosphorus than does urine, was only discovered 100 years later by Scheele.

(2) *Note.* Experience has shown that a higher percentage of phosphorus can be obtained from bone-ash if only two-thirds of the calcium are removed by means of  $\text{H}^2\text{SO}^4$ , leaving soluble calcium phosphate to be separated by filtration, to be evaporated, mixed with coal and dried before distillation. This plan is followed in the match factories.

There are three *modifications* of phosphorus.

(1). *Common, pale-colored, phosphorus* which crystallizes in octahedrons has essentially the properties already given. Specific gravity = 1.83 at  $10^\circ \text{C}$ . Above  $45^\circ \text{C}$ ., that is, in the liquid state, the specific gravity decreases considerably ; with the temperature at  $100^\circ \text{C}$ . it is 1.695 ; at  $200^\circ \text{C}$ . = 1.603. At the boiling-point = 1.485. Phosphorus boils at  $250^\circ \text{C}$ . and distills over in an atmosphere of hydrogen. Rapid cooling of the vapor throws the phosphorus out in a fluffy, snow-white condition (flowers of phosphorus). But it is quite certain that slight volatilization goes on at ordinary temperature. To this volatilization

is probably due the *phosphorescence*, the power to emit light in a dark room. Surrounded by oxygen alone there is no phosphorescence; but the latter phenomenon appears when the oxygen is diluted with nitrogen. Phosphorus ignites at  $60^{\circ}$  C. in air. Rubbing on a rough surface causes ignition (matches). Mixed with  $\text{KClO}^3$  a very explosive substance results (lucifer matches).

Phosphorus is poison to man and animals when brought into the stomach or esophagus. Every particle of the phosphorus causes an intense local inflammation of the membrane, hence great pain and shock, which result in death. 0.2 gram may be a fatal dose for an adult person. Emptying the stomach by emetics and the pump may save the life in some cases. The workmen in match factories are known to suffer from necrosis of the teeth, the gums and the jaw-bones themselves. Burns made by burning phosphorus on the fingers have even been fatal. Wash out such wounds with utmost haste with a dilute water solution of bleaching lime, or bleaching soda (crude Javelle).

The best solvent for the active form of phosphorus is carbon disulfid.

(2). *Amorphous red phosphorus*. Sunlight, especially the violet portion of it, or heat plus pressure, or the electric current, changes the ordinary phosphorus more or less rapidly into the amorphous modification. In the factories the change is brought about by keeping the yellow phosphorus for 10 days at a steady temperature of  $260^{\circ}$  C.

The amorphous red phosphorus is not poisonous. It does not melt even at red heat but volatilizes. It is *not* soluble in  $\text{CS}^2$  nor in  $\text{KOH}$ . It appears as a scarlet or as a purplish-red, pulverulent mass, sometimes brown-red. In bulk it shows sometimes a weak metallic luster, more often no luster at all. It has neither taste nor odor, does not show phosphorescence. It ignites at  $260^\circ \text{C}$ . It forms with  $\text{KClO}^3$ , with  $\text{PbO}^2$ , and with  $\text{MnO}^2$ , mixtures which ignite by blows or friction. It is the only phosphorus now used in matches, or on the friction surfaces of safety match boxes.

(3). *Black crystallized phosphorus*. By heating red phosphorus in a vacuum to  $447^\circ \text{C}$ . it is obtained as a violet-black mass of conchoidal fracture, or by fusing together in a vacuum phosphorus and metallic lead. After cooling one finds long-stretched rhombohedrons, black in reflected light, red in transmitted light. This phosphorus has a specific gravity of 2.34.

Atomic weight of phosphorus,  $\text{P} = 31$ . The determinations of the vapor density lead to 62. We assume therefore that the element in the free state is  $\text{P}^2$ . According to the combinations into which phosphorus enters with the non-metals, it is trivalent and pentavalent like nitrogen.

The oxyds of phosphorus are  $\text{P}^2\text{O}^5$ ,  $\text{P}^2\text{O}^3$ .

*Phosphorus pentoxyd*,  $\text{P}^2\text{O}^5$ , a colorless, vitreous solid or colorless triclinic crystals. Dissolves in water. Is very hygroscopic (goes slowly into a syrup when standing in moist air), hence it is often



used to dry gases. It has no odor, but a very *sour* taste. Is sometimes called *anhydrous phosphoric acid*. Volatilizes partly at  $250^{\circ}$  C. But when heated quickly it changes its nature by polymerization (aggregation of molecules) and is much less volatile.

*Preparation.* By igniting phosphorus in a flask in a current of perfectly dry air. The product is a mass of minute snowy-white crystals. For larger quantities a tinned sheet-iron cylinder is substituted for the flask.

*Phosphoric acids.* It was Graham who first demonstrated that the pentoxyd can form 3 hydrates and that these hydrates possess very distinct properties. We distinguish these hydrates thus: The trihydrate,  $3\text{H}^2\text{O}.\text{P}^2\text{O}^5$ , the dihydrate,  $2\text{H}^2\text{O}.\text{P}^2\text{O}^5$  and the monohydrate,  $\text{H}^2\text{O}.\text{P}^2\text{O}^5$ . We translate these hydrates into the radical expressions or hydrogen acids thus:

Trihydrate,  $3\text{H}^2\text{O}.\text{P}^2\text{O}^5 = 2\text{H}^3(\text{PO}^4) =$   
orthophosphoric acid.

Dihydrate,  $2\text{H}^2\text{O}.\text{P}^2\text{O}^5 = \text{H}^4(\text{P}^2\text{O}^7) =$   
pyrophosphoric acid.

Monohydrate,  $\text{H}^2\text{O}.\text{P}^2\text{O}^5 = 2\text{H}(\text{PO}^3) =$   
metaphosphoric acid.

*Orthophosphoric acid,  $\text{H}^3(\text{PO}^4)$ .* Orthorhombic, colorless crystals, or a thick syrup of specific gravity = 1.88. Strong acid reaction; easily soluble in water. Gives green coloration to a flame.

*Preparation.* (1). By acting with  $\text{HNO}^3$  (specific



gravity = 1.2) upon ordinary phosphorus in a glass retort, (1 phosphorus, 10 acid), at such a temperature that lively action ensues, but not a violent one, (because explosions may set in). After all the phosphorus has disappeared heat to boiling and distill over about 7 parts of the  $\text{HNO}^3$ . The distillate has a specific gravity of 1.1–1.14 and may be used for another operation by adding enough concentrated acid to bring gravity up to 1.2. Pour the liquid from the retort into an evaporating dish and evaporate carefully to syrup, or until all  $\text{HNO}^3$  has been removed. Sometimes there occurs during this stage another disengagement of NO from the fact that some  $\text{P}^2\text{O}^3$  is still present. The temperature may be brought to  $188^\circ \text{C}$ . but not higher, for pyro-phosphoric acid may form. By adding some alcohol the remainder of  $\text{HNO}^3$  may be removed more easily. The syrup of  $\text{H}^3(\text{PO}^4)$  is called *glacial phosphoric acid*.

(2). By acting upon red amorphous phosphorus with concentrated  $\text{HNO}^3$ . The oxydation is more rapid and can be carried on in a beaker glass.

*Orthophosphates.* The orthophosphoric acid can form 3 series of salts as follows :

Monads,  $\text{Na}^3(\text{PO}^4)$ ,  $\text{HNa}^2(\text{PO}^4)$ ,  $\text{H}^2\text{Na}(\text{PO}^4)$ .

Diads,  $\text{Ca}^3(\text{PO}^4)^2$ ,  $\text{Ca}^2\text{H}^2(\text{PO}^4)^2$ ,  $\text{CaH}^4(\text{PO}^4)^2$ .

Triads,  $\text{Al}^3(\text{PO}^4)^3$ ,  $\text{Al}^2\text{H}^3(\text{PO}^4)^3$ ,  $\text{AlH}^6(\text{PO}^4)^3$ .

The radical  $(\text{PO}^4)$  is trivalent, hence  $\text{Na}^3(\text{PO}^4)$  is a fully saturated combination and so is  $\text{Al}(\text{PO}^4)$ , because aluminum is trivalent. But in order to bring

out the partially saturated series ( $\text{Al}^2\text{H}^3$ ) and ( $\text{AlH}^6$ ), we must treble the saturated molecule into  $\text{Al}^3(\text{PO}^4)^3$ . Any diad-metal will form orthophosphates similar to calcium. For example, copper will make  $\text{Cu}^3(\text{PO}^4)^2$ ,  $\text{Cu}^2\text{H}^2(\text{PO}^4)^2$ ,  $\text{CuH}^4(\text{PO}^4)^2$ . The three series are sometimes called *basic*, *neutral*, *acid*.


$\text{Cu}^3(\text{PO}^4)^2$  is basic copper orthophosphate.

$\text{Cu}^2\text{H}^2(\text{PO}^4)^2$  is neutral copper orthophosphate.

$\text{CuH}^4(\text{PO}^4)^2$  is acid copper orthophosphate.

The orthophosphates of potassium, sodium, ammonium are all soluble in water and all can be crystallized. The most common of them, because most easily obtainable, is  $\text{Na}^2\text{H}(\text{PO}^4) + 12\text{H}^2\text{O}$  in monoclinic crystals, and  $\text{H.NH}^4.\text{Na}(\text{PO}^4) + 4\text{H}^2\text{O}$ , also monoclinic crystals. The latter salt goes under the names: *salt of phosphorus*, and *microcosmic salt*. This salt fuses into a perfect glass and dissolves at red heat most of the metallic oxyds, giving with some of them transparent glasses of constant color. Hence we utilize this salt as a flux in blow-pipe analysis.

*Preparation of microcosmic salt.* Dissolve 353 grams of the crystals of  $\text{Na}^2\text{HPO}^4 + 12\text{H}^2\text{O}$  and 53.5 grams of  $\text{NH}^4\text{Cl}$  (sal ammoniac) in 1700 c.c. of warm water, filter and evaporate until a film of crystals forms at the surface. Let stand for several days in a cool place. A large crop of the microcosmic salt will have been formed. Drain crystals from mother liquor. Dissolve again in water and crystallize, repeating the recrystallization twice. Then you will have crystals sufficiently free from  $\text{NaCl}$  to answer for blow-pipe work. Reaction :



NaCl remaining in the crystals causes the bead, after fusion, to become white and opaque.

*Insoluble orthophosphates.* The neutral solutions of all the metals are precipitated by adding a solution of any alkali phosphate (K, Na,  $\text{NH}^4$ ). These precipitates are soluble in dilute acids—even acetic acid. Two of these precipitates are of special interest because by means of them we distinguish orthophosphoric acid from other acids.

(1). *Silver orthophosphate*,  $\text{Ag}^3(\text{PO}^4)$ , a yellow flocculent precipitate. The solution must be *neutral* before  $\text{AgNO}^3$  is added to the unknown.


(2). *Ammonium-magnesium orthophosphate*,  $\text{NH}^4.\text{Mg}(\text{PO}^4)$ , a colorless, granular or crystalline precipitate which forms when  $\text{MgCl}^2$  or  $\text{MgSO}^4$  solution is added to a slightly ammoniacal solution of an orthophosphate.

The most delicate or sensitive reagent for orthophosphate is the so-called *molybdic solution*. This is a solution of  $(\text{NH}^4)^2\text{MoO}^4$ —ammonium molybdate—in  $\text{HNO}^3$ , specific gravity 1.2; the solution is pale yellow or colorless. Any metallic phosphate is first dissolved in a little  $\text{HNO}^3$ , or any unknown substance is heated with  $\text{HNO}^3$ , water added and after filtering 1 volume of the molybdic solution is added. The temperature of the liquid is brought to about  $50^\circ \text{C}$ ., and the liquid shaken rapidly. A fine granular, citron-yellow precipitate falls if a phosphate be present. The precipitate is  $(\text{NH}^4)^2$ -

$\text{H}(\text{PO}^4).10\text{MoO}^3 + 1\frac{1}{2}\text{H}^2\text{O} = \text{ammonium-hydrogen phosphopolymolybdate}.$

*Pyrophosphoric acid*,  $\text{H}^4(\text{P}^2\text{O}^7)$ , is not known in solid state, only known as an aqueous solution.

*Preparation.* Heat the salt  $\text{Na}^2\text{H}(\text{PO}^4) + 12\text{aq.}$  in a crucible until all the water is driven out and then to redness for a short time. Reaction :



Dissolve in water without heating. To solution add  $\text{PbA}^2$  (lead acetate); a white precipitate  $\text{Pb}^2(\text{P}^2\text{O}^7)$  falls. Filter and wash. Suspend the precipitate in water and pass  $\text{H}^2\text{S}$  into the liquid; then you obtain  $\text{Pb}^2(\text{P}^2\text{O}^7) + 2\text{H}^2\text{S} = 2\text{PbS} + \text{H}^4(\text{P}^4\text{O}^7)$ , which are separated by filtering.

*Properties.* (1). When the acid is neutralized with  $\text{NH}^4\text{OH}$  and  $\text{AgNO}^3$  is added, a *white* flocculent precipitate falls (not yellow as with an orthophosphate). (2).  $\text{MgCl}^2$  does not produce a precipitate.

When the precipitate  $\text{MgNH}^4(\text{PO}^4)$  (see above) is ignited  $\text{Mg}^2(\text{P}^2\text{O}^7)$  magnesium pyrophosphate is left behind. A solution of pyrophosphoric acid reverts into orthophosphoric acid by boiling for several hours.

*Metaphosphoric acid*,  $\text{H}(\text{PO}^3)$ . A colorless glassy substance.

*Preparation.* (1) By dissolving  $\text{P}^2\text{O}^5$  in cold water. (2) By heating the syrup of  $\text{H}^3\text{PO}^4$ , thus  $\text{H}^3\text{PO}^4 + \text{heat} = \text{H}^2\text{O} + \text{H}(\text{PO}^3)$ . (3) By fusing the microcosmic salt at a red heat we get

$\text{Na}(\text{NH}^4)\text{H}(\text{PO}^4) + \text{heat} = \text{Na}(\text{PO}^3) + \text{NH}^3 + \text{H}^2\text{O}$   
 sodium metaphosphate + ammonia + water. The characteristics of metaphosphoric acid are as follows: (a) the acid solution causes coagulation (curdling) in a water solution of albumen. (Neither ortho- nor pyrophosphoric acid coagulates the albumen.) (b) When the solution is neutralized with  $\text{NH}^4\text{HO}$ ,  $\text{AgNO}^3$  solution gives a white gelatinous precipitate. (c) When a solution of  $\text{Na}(\text{PO}^3)$  is added to neutral salts of the metals, a precipitate forms at first, but dissolves on further addition of the metaphosphate. Some of the precipitates separate like tough resin, some separate as oily liquids.

*Phosphorus trioxyd*,  $\text{P}^2\text{O}^3$ . White snowy aggregates of small crystals often in the shape of trees or ferns.

*Preparation.* Heat phosphorus in a tube until it ignites, and allow a very slow current of dry air to pass through the tube. The oxyd sublimes into a receiver. In the dark it remains unchanged. Sunlight brings red or orange colors. In warm oxygen it ignites and changes to  $\text{P}^2\text{O}^5$ .

It smells like phosphorus. Some authors say it is poisonous, others say it is not; I, myself, think it is poisonous. Very slightly soluble in water.

*Phosphorous acid*, *hydrogen phosphite*,  $\text{H}^3(\text{PO}^3)$ . Crystalline-white mass or distinct crystals. Soluble in water. Sour taste. Is a strong deoxydizing agent. In salts of gold, silver, copper, mercury, the acid causes *precipitation of the metal*.

It is a diatomic acid, that is to say, only two of



the hydrogens can be replaced by a metal. There are therefore two series of *phosphites*,  $\text{Na}^2\text{H}(\text{PO}^3)$  and  $\text{NaH}^2.(\text{PO}^3)$ .

*Preparation.* (1). By action of dilute  $\text{HNO}^3$  upon phosphorus. (2). By the action of oxalic acid upon phosphorus trichlorid. Thus :



*Hypophosphorous acid, hydrogen hypophosphite,  $\text{H}^3\text{PO}^2$ .* A colorless substance in large scales or leaves. Its solution in water is even more deoxydizing than the preceding phosphorous acid. It is a monobasic acid. Only one of the three hydrogens can be replaced by a metal. Thus  $\text{NaH}^2(\text{PO}^2)$  or still better  $\text{Na}(\text{HPO}^2\text{H})$  = sodium hypophosphite, or  $\text{Ba}(\text{HPO}^2\text{H})^2$  = barium hypophosphite. The hypophosphites have been recommended as very active stimulants of the nerves and the brain (humbug).

*Preparation.*  $3\text{KHO} + 4\text{P} + 3\text{H}^2\text{O} = 3\text{KH}^2\text{PO}^2 + \text{PH}^3$ . This means that phosphorus in presence of water and KHO will form potassium hypophosphite plus phosphine ( $\text{PH}^3$ ).  $\text{Ca}(\text{HO})^2$  and  $\text{Ba}(\text{HO})^2$  plus P act similarly.

#### COMBINATIONS OF PHOSPHORUS WITH CHLORINE.

*Phosphorus trichlorid,  $\text{PCl}^3$ .* Colorless liquid. Produces white fumes in moist air; refracts the light strongly, smell, penetrating, and the vapor causes tears to flow. Boils at  $76^\circ \text{C}$ .

When  $\text{PCl}^3$  is poured into cold water it sinks to the bottom and collects like a heavy oil; soon a reaction sets up between the water and the chlorid,



$\text{PCl}^3 + 3\text{H}^2\text{O} = \text{H}^3.\text{PO}^3 + 3\text{HCl}$ , the result being a liquid containing phosphorous acid and hydrochloric acid. Such a solution answers as a deoxydizer.

*Preparation.* Place in a retort some pieces of dried stick phosphorus (dry with blotting paper), the retort having been previously filled with  $\text{CO}^2$  gas. In the tubulus fits a cork, and through this passes a glass tube down to the phosphorus. A receiver is tightly connected with the retort, and is well cooled. Fill now the retort with chlorine, and warm the retort until the phosphorus melts, when the action begins and  $\text{PCl}^3$  distills over.

*Phosphorus pentachlorid,  $\text{PCl}^5$ .* A colorless solid. Peculiar odor, fumes at the air. With little water it decomposes into  $\text{HCl}$  and  $\text{POCl}^3$ . It is often used in synthetic laboratory work to put  $\text{Cl}$  into complex molecules. With sodium it gives  $2\text{NaCl} + \text{PCl}^3$ , and the same with other metals. With much water it decomposes into phosphoric acid and  $\text{HCl}$ .


$\text{PCl}^5 + 4\text{H}^2\text{O} = 5\text{HCl} + \text{H}^3(\text{PO}^4)$ , orthophosphoric acid.

*Preparation.* By acting with excess of  $\text{Cl}$  upon  $\text{PCl}^3$ . There are, of less importance,  $\text{PBr}^3$ ,  $\text{PBr}^5$ ,  $\text{PI}^3$ ,  $\text{PI}^5$ ,  $\text{PF}^5$ .


*Phosphine, hydrogen phosphid,  $\text{PH}^3$ .* A gas of disagreeable odor, somewhat resembling that of garlic. In a dark room the gas gives out a pale light like phosphorus itself. It causes a taste on the tongue. Sunlight decomposes the gas into hydrogen and red amorphous phosphorus. The gas

is poisonous because the blood absorbs it like  $\text{H}(\text{CN})$ . Air containing 0.25 per cent.  $\text{PH}^3$  kills animals in 5–10 minutes. Phosphine ignites in air at  $149^\circ \text{C}$ .; the flame is white and yields white smoke. But sometimes the gas is *self-igniting*. This self-ignition is attributed to the admixture of another compound  $\text{PH}^2$ , which latter forms only under specific conditions. When phosphine gas is passed into solutions of Ag, Hg, Cu, Pb, Bi, Au salts, the metals are thrown out; or phosphids of the metals are formed.  $\text{PH}^3$  in a solution of  $\text{AgNO}^3 + \text{water}$  causes first a yellow precipitate of  $\text{Ag}^3\text{P} \cdot 3\text{AgNO}^3$  (? doubtful composition), but black  $\text{Ag}^3\text{P}$  results ultimately.


*Preparation.* (1). Place 5–10 grams of stick phosphorus into a 150 c.c. flask. Fill the flask with KOH solution (1:5) up to the stopper. The latter carries one evolution tube, bent so that it can be made to dip under water in a dish or beaker glass. (The flask is to be filled completely to avoid explosion with air). On heating the flask the gas evolution will set in, and if the water in the dish be warm, each gas bubble will ignite as it breaks over the water, and will form a ring of smoke in the air. (2). Place the phosphorus in the flask as before, but fill the flask with an *alcoholic* solution of KOH. (70 per cent. alcohol.) The gas will *not ignite* by itself. The reason for this is that the self-igniting  $\text{PH}^2$  remains dissolved in the alcohol; does not mix with the gas  $\text{PH}^3$ . In both these actions the  $\text{PH}^3$  is generated by the reaction



(3). Prepare  $\text{Na}^3\text{P}$  by fusing together sodium and phosphorus. Or by fusing together



In either case you get  $\text{Na}^3\text{P}$  and if a drop of water touches  $\text{Na}^3\text{P}$  then phosphine will be disengaged (noticed by strong odor), and  $\text{NaHO}$  will form :



Metallic magnesium can be carried much better than sodium because it does not oxydize so easily at ordinary temperature. Hence the last reaction is the one best adapted to test an unknown mineral for phosphorus, in the field, in the operation of blow-pipe analysis. We call it the *phosphine reaction*. It is all done in a small ignition tube.

*Composition of  $\text{PH}^3$ .* Phosphine breaks up readily at a red heat into  $\text{P} + \text{H}$ . If 20 c.c. of phosphine are collected over mercury in a eudiometer (see ammonia) and the spark is sent through it, complete dissociation results in from 6 to 10 minutes, when the volume has increased to 30 c.c. Phosphorus covers the surface of the tube and the gas consists entirely of hydrogen. Now since the volume of solid phosphorus is so small as to be negligible it follows that 20 c.c. of phosphine contain 30 c.c. of hydrogen and 10 c.c. of phosphorus gas. Hence  $\text{PH}^3$ .

*Phosphonium,  $\text{PH}^4$ ,* corresponds to *ammonium,  $\text{NH}^4$ ,* and is only known hypothetically. Because

$\text{PH}^3$  combines by simple addition with  $\text{HCl}$ , giving  $\text{PH}^4\text{Cl}$  (colorless crystals below  $20^\circ \text{C.}$ ).

*Liquid hydrogen phosphid,  $\text{PH}^2$ .* A colorless liquid, not soluble in water. In contact with air ignites instantaneously. Forms when  $\text{Ca}^3\text{P}^2$  is decomposed with  $\text{H}^2\text{O}$  and the resulting gas is carried through a U-tube standing in the freezing mixture. At the same time with the liquid  $\text{PH}^2$ , condenses a solid substance which has probably the composition  $\text{P}^2\text{H}$ . Phosphorus combines directly with all metals, yielding *metallic phosphids*. Of practical importance are the phosphids of iron (in the metallurgy of iron and steel) and tin phosphid,  $\text{Sn}^4\text{P}$ , beautiful silver-white crystals (in the manufacture of *phosphorbronze*).

## APPENDIX.

THE CHEMICAL ELEMENTS, THEIR SYMBOLS, EQUIVALENTS  
AND SPECIFIC GRAVITIES.

Name.	Symbol.	Mass Unit Weight.	Specific Gravity.
Aluminium . . . . .	Al.	27.5	2.56
Antimony. . . . .	Sb.	122.0	6.70
Arsenic. . . . .	As.	75.0	5.70
Barium. . . . .	Ba.	137.0	4.00
Bismuth . . . . .	Bi.	210.0	9.7
Boron . . . . .	B.	11.0	2.63
Bromine . . . . .	Br.	80.0	5.54
Cadmium . . . . .	Cd.	112.0	8.60
Caesium . . . . .	Cs.	133.0	1.88
Calcium . . . . .	Ca.	40.0	1.58
Carbon . . . . .	C.	12.0	3.50
Cerium . . . . .	Ce.	92.0	6.68
Chlorine . . . . .	Cl.	35.5	2.45
Chromium . . . . .	Cr.	52.5	6.81
Cobalt . . . . .	Co.	58.8	7.7
Columbium . . . . .	Cb.	184.8	6.00
Copper . . . . .	Cu.	63.0	8.96
Didymium . . . . .	Di.	96.0	6.54
Erbium. . . . .	E.	112.6	—
Fluorine . . . . .	F.	19.0	1.32
Gallium . . . . .	Ga.	69.9	5.9
Glucinum. . . . .	Gl.	9.5	2.1
Gold (Aurum). . . . .	Au.	196.0	19.3
Hydrogen . . . . .	H.	1.0	0.069
Indium . . . . .	In.	113.4	7.4
Iodine . . . . .	I.	127.0	4.94
Iridium. . . . .	Ir.	198.0	21.15
Iron (Ferrum). . . . .	Fe.	56.0	7.79
Lanthanum . . . . .	La.	90.2	11.37
Lead (Plumbum) . . . . .	Pb.	207.0	11.44
Lithium . . . . .	Li.	7.0	0.59

Name.	Symbol.	Atomic Weight.	Specific Gravity.
Magnesium . . . . .	Mg.	24.0	1.75
Manganese . . . . .	Mn.	55.0	8.01
Mercury (Hydrargyrum) . .	Hg.	200.0	13.59
Molybdenum . . . . .	Mb.	96.0	8.60
Nickel . . . . .	Ni.	58.8	8.60
Niobium . . . . .	Nb.	94.0	6.27
Nitrogen . . . . .	N.	14.0	0.972
Osmium . . . . .	Os.	199.0	21.40
Oxygen . . . . .	O.	16.0	1.105
Palladium . . . . .	Pd.	106.5	11.60
Phosphorus . . . . .	P.	31.0	1.83
Platinum . . . . .	Pt.	197.4	21.53
Potassium (Kalium) . . . .	K.	39.0	0.865
Rhodium . . . . .	Ro.	104.3	12.1
Rubidium . . . . .	Rb.	85.4	1.52
Ruthenium . . . . .	Ru.	104.4	11.4
Selenium . . . . .	Se.	79.5	4.78
Silicon . . . . .	Si.	28.0	2.49
Silver (Argentum) . . . . .	Ag.	108.0	10.5
Sodium (Natrium) . . . . .	Na.	23.0	0.972
Strontium . . . . .	Sr.	87.6	2.54
Sulphur . . . . .	S.	32.0	2.05
Tantalum . . . . .	Ta.	182.0	10.78
Tellurium . . . . .	Te.	129.0	6.02
Thallium . . . . .	Tl.	204.0	11.91
Thorium . . . . .	Th.	115.7	7.8
Tin (Stannum) . . . . .	Sn.	118.0	7.28
Titanium . . . . .	Ti.	50.0	4.3
Tungsten (Wolfram) . . . .	W.	184.0	17.6
Uranium . . . . .	U.	120.0	18.4
Vanadium . . . . .	V.	51.3	5.50
Yttrium . . . . .	Y.	61.7	—
Zinc . . . . .	Zn.	65.0	7.14
Zirconium . . . . .	Zr.	89.5	4.15



TABLE FOR THE COMPARISON OF THE SCALES OF RÉAUMUR'S,  
CELSIUS'S, AND FAHRENHEIT'S THERMOMETERS.

Réaumur.	Celsius.	Fahrenheit.	Réaumur.	Celsius.	Fahrenheit.
-15	-18.75	—	33	+41.25	+106.25
14	17.50	+0.50	34	42.50	108.50
13	16.25	2.75	35	43.75	110.75
12	15.00	5.00	36	45.00	113.00
11	13.75	7.25	37	46.25	115.25
10	12.50	9.50	38	47.50	117.50
9	11.25	11.75	39	48.75	119.75
8	10.00	14.00	40	50.00	122.00
7	8.75	16.25	41	51.25	124.25
6	7.50	18.50	42	52.50	126.50
5	6.25	20.75	43	53.75	128.75
4	5.00	23.00	44	55.00	131.00
3	3.75	25.25	45	56.25	133.25
2	2.50	27.50	46	57.50	135.50
+1	1.25	29.75	47	58.75	137.75
0	0	32.00	48	60.00	140.00
1	+1.25	34.25	49	61.25	142.25
2	2.50	36.50	50	62.50	144.50
3	3.75	38.75	51	63.75	146.75
4	5.00	41.00	52	66.00	149.00
5	6.25	43.25	53	66.25	151.25
6	7.50	45.50	54	67.50	153.50
7	8.75	47.75	55	68.75	155.75
8	10.00	50.00	56	70.00	158.00
9	11.25	52.22	57	71.25	160.25
10	12.50	54.50	58	72.50	162.50
11	13.75	56.75	59	73.75	164.75
12	15.00	59.00	60	75.00	167.00
13	16.25	61.25	61	76.25	169.25
14	17.50	63.50	62	77.50	171.50
15	18.75	65.75	63	78.75	173.75
16	20.00	68.00	64	80.00	176.00
17	21.25	70.25	65	81.25	178.25
18	22.50	72.50	66	82.50	180.50
19	23.75	74.75	67	83.75	182.75
20	25.00	77.00	68	85.00	185.00
21	26.25	79.25	69	86.25	187.25
22	27.50	81.50	70	87.50	189.50
23	28.75	83.75	71	88.75	191.75
24	30.00	86.00	72	90.00	194.00
25	31.25	88.25	73	91.25	196.25
26	32.50	90.50	74	92.50	198.50
27	33.75	92.75	75	93.75	200.75
28	35.00	95.00	76	95.00	203.00
29	36.25	97.25	77	96.25	205.25
30	37.50	99.50	78	97.50	207.50
31	38.75	101.75	79	98.75	209.75
32	40.00	104.00	80	100.00	212.00

*Rules for the Conversion of the Different Thermometer  
Degrees into each other.*

The thermometers referred to in the table are graduated so that the range of temperature, between the freezing and boiling points of water, is divided by Fahrenheit's scale into 180 (from  $32^{\circ}$  to  $212^{\circ}$ ) by Celsius's into 100 (from 0 to  $100^{\circ}$ ), and by that of Réaumur into 80 (from 0 to  $80^{\circ}$ ) portions or degrees.

The spaces occupied by a degree of each scale are consequently as  $\frac{1}{9}$ ,  $\frac{1}{5}$  and  $\frac{1}{4}$  respectively, or as 1, 1.8 and 2.25; and the number of degrees denoting the same temperature, by the three scales, when reduced to a common point of departure by subtracting 32 from Fahrenheit's, are as 9, 5, and 4. Hence we derive the following equivalents :

A degree of Fahrenheit is equal to 0.5 of Celsius's, or to 0.4 of Réaumur's; a degree of Celsius's is equal to 1.8 of Fahrenheit's, or to 0.8 of Réaumur's; and a degree of Réaumur's is equal to 2.25 of Fahrenheit's, or to 1.25 of Celsius's.

To convert degrees of Fahrenheit into Celsius's or Réaumur's, subtract 32 and multiply the remainder by  $\frac{5}{9}$  for Celsius's, or  $\frac{4}{9}$  for Réaumur's.

To convert degrees of Celsius's or Réaumur's into Fahrenheit's, multiply Celsius's by  $\frac{9}{5}$ , or Réaumur's by  $\frac{9}{4}$ , as the case may be, and add 32 to the product.

TABLE OF THE LITER WEIGHTS OF THE GASES.

*Temp. 0° C. and 760 Mm. Pressure.*

1 liter of air weighs.		1.29300 Gms.
" CO.	"	1.25078 "
" CO <sub>2</sub> .	"	1.96500 "
" O.	"	1.42910 "
" H.	"	0.08988 "
" N.	"	1.25070 "
" NO.	"	1.34260 "
" NO <sub>2</sub> .	"	2.05440 "
" N <sub>2</sub> O.	"	1.96770 "
" NH <sub>3</sub> .	"	0.76170 "
" Cl.	"	3.17240 "
" HCl.	"	1.62850 "
" H <sub>2</sub> S.	"	1.52100 "
" SO <sub>2</sub> .	"	2.86150 "
" CH <sub>4</sub> .	"	0.71570 "
" steam*	"	0.58960 "
" C <sup>2</sup> H <sup>2</sup> .	"	1.16200 "
" Br.	"	7.14259 "
" I.	"	11.27100 "
" S.	"	2.84300 "
" P.	"	5.63180 "
" Hg.	"	9.02100 "
" HI.	"	5.71067 "
" HBr.	"	3.61607 "
" C <sup>2</sup> N <sup>2</sup> .	"	2.32653 "

\* At 100° C. and 760 Mm.

The values in the above table may be calculated from the relation between the molecular weight in grams of the substance and its liter weight. This relation may be expressed as an equation thus :

$$\frac{\text{weight of gram-molecule}}{\text{weight of a liter}} = \text{constant} = 22.38,$$

$$\text{or liter weight} = \frac{\text{weight of gram molecule.}}{22.38}$$



# INDEX.

## ACETATE, 359

Acid, 72, 124, 129  
   abietinic, 379  
   acetic, 278, 358  
   carbolic, 282  
   citric, 366  
   cyanuric, 399  
   formic, 305, 358  
   fulminic, 400  
   gallic, 368  
   gluco-sulfuric, 343  
   hydrobromic, 138  
   hydrochloric, 108  
   hydrofluoric, 148  
   hydroiodic, 142  
   hypophosphorous, 414  
   lactic, 364  
   malic, 366  
   metaphosphoric, 412  
   muriatic, 108  
   nitric, 164  
   orthophosphoric, 408  
   oxalic, 361  
   pan, 227  
   phosphoric, 408  
   phosphorous, 413  
   picric, 282  
   prussic, 395  
   pyrogallic, 369  
   pyroligneous, 276  
   pyrophosphoric, 412  
   sulfuric, 50, 52, 216  
   tannic, 366  
   tartaric, 364  
 Air, weight of, 13  
 Albumen, blood, 386  
   vegetable, 387  
 Albumenoids, 387  
 Alcohol, 346, 348  
   absolute, 349

Aldehyde, 354  
 Alkali, 91  
 Alkaline reaction, 91, 187  
 Alkaloids, 383  
 Aluminum reactions, 246  
 Amalgam, 87  
 Ammonia, 186, 190  
   liquid, 200  
 Ammonium, 194  
 Amylum, 340  
 Anilin, 321  
 Anode, 26  
 Anthracite, 308  
 Aqua fortis, 156  
   regia, 156  
 Arabin, 345  
 Atom, 33  
 Atomic weights, 34, 126  
 Atropin, 384

## BAKING powders, 121

Balsam, 378  
 Base, 72, 129  
 Benzene, 337  
 Bleaching, 104  
   lime, 104  
 Bone-ash, 401  
   black, 390  
 Braise, 85  
 Bromates, 139  
 Bromine, 137  
 Bromo seltzer, 139  
 Brucin, 384  
 Butane, 303

## CAFFEIN, 383

Calcite, 64  
 Calcium reactions, 252  
   sulfid, 241  
   tartrate, 365

- Calcespar, 64  
 Caoutchouc, 380  
 Caps, percussion, 356  
 Carbamid, 389  
 Carbon, 202, 257  
     atomic weight of, 262  
     dioxyd, 259  
     disulfid, 254  
     hydrates, 270  
     monoxyd, 262  
 Carbonates, 90  
 Carboys, 113  
 Casein, 388  
 Cathode, 26  
 Caustic lime, 70  
     soda, 122  
 Cellulose, 264, 270, 305  
 Chamber acid, 227  
 Charcoal, animal, 389  
     making, 274  
 Charmotte, 85  
 Chemical elements, their sym-  
     bols, equivalents, and specific  
     gravities, 419, 420  
 Chinin, 384  
 Chloral, 354  
 Chlorates, 105, 135  
 Chlorid, silver, 111  
 Chlorine, 98, 104  
 Chloroform, 353  
 Clay, fire, 41  
 Coal, bituminous, 308  
     brown, 309  
     cannel, 309, 311  
     distillation of, 316  
     hard, 308  
     origin of, 312  
     -tar, 320  
 Cocaine, 385  
 Coke, 317  
 Collodion, 272  
 Colophony, 379  
 Compounds, 16  
 Copper oxyd, 16, 47  
     reactions, 250  
 Copperas, 35  
     action of, on leather, 36  
 Covellite, 234  
 Creatinin, 388  
 Cyanates, 394, 399  
 Cyanids, 393  
 Cyanogen, 396  
**D**ECREPITATION, 93  
     Deoxydation, 17  
 Dextrin, 341  
 Diamond, 257  
 Dichloroxyd, 134  
 Dichroism, 336  
 Dissociation, 325  
 Dynamite, 375  
**E**BONITE, 382  
     Electrodes, 26  
 Electrolysis, 25  
 Electrolyte, 26  
 Elements, 16  
     their symbols, equivalents,  
     and specific gravities, 419,  
     420  
 Endothermic reaction, 330  
 Etching, 149  
 Ether, 352  
 Ethylene, 324  
 Eudiometer, 31  
 Exothermic reaction, 330  
**F**ATS, 371  
     Fermentation, 199, 347  
 Ferricyanates, 398  
 Ferrocyanates, 392  
 Fibrin, 388  
 Fibroin, 389  
 Fire-clay, 41  
 Flash-point of an oil, 337  
 Fluorescein, 336  
 Fluorids, 148  
 Fluorine, 144, 150  
 Fluorite, 144, 150  
     gas, 146  
 Fluorspar, 144  
 Flux, 145  
 Formate, 358  
 Fuchsin, 322  
 Fuel-gas, 333  
 Fulminate, 355  
 Fulminating gas, 27  
**G**ALENA, 181  
     Galvani, 125  
 Gangue, 144  
 Gas, analysis apparatus for, 285



Gas, fulminating, 27  
 generator, 101, 103  
 natural, 337  
 works, plan of, 326  
 Gases, table of the liter weights  
 of the, 423  
 Gasoline, 337  
 Gay-Lussac tower, 225  
 Gelatine, 389  
 blasting, 375  
 Glass ink, 150  
 Glover tower, 222  
 Glucose, 343  
 Glue, 389  
 Gluten, 209, 339  
 Glycerin, 374  
 nitro-, 375  
 Gold, atomic-weight, 131  
 chlorids, 131  
 reactions, 246  
 Graphite, 257  
 Green vitriol, 35, 36  
 Gum arabic, 345  
 Gun-cotton, 271  
 powder, 166  
 Gutta-percha, 380  
**H**ÆMATIN, 387  
 Haemoglobin, 387  
 Hair, 388  
 Heat, specific, 132  
 Horn, 388  
 Hydrates, of potassium, 79  
 of sulfuric acid, 55  
 Hydrogen, 25, 61  
 chlorid, 107  
 peroxyd, 34  
 sulfate, 216  
 sulfid, 234, 237, 253  
 Hydrometer, 112  
 Hydroxyd, 71  
 Hydroxyl, 129  
 Hypo, 243  
 Hyposulfites, 243  
 Hyssin, 388  
**I**CE, 19  
 Infusorial earth, 375  
 Ink, 36, 368  
 Iodates, 143  
 Iodids, 142

Iodine, 139, 141  
 and starch, 141  
 Iodoform, 354  
 Iodyrite, 144  
 Iron, reactions of, 246  
 Isomerids, 303  
**K**ELP, 122  
 Kerosene, 337  
 Knee tube, 52  
 Koenig generator, 101  
**L**ACTOSE, 345  
 Laming's mass, 328  
 Laughing gas, 182  
 Law of combination by volume,  
 134  
 Dulong and Petit, 133  
 Lead, reactions of, 246  
 Leather, 368  
 Leblanc process, 119  
 Levulose, 344  
 Lignite, 309  
 Lime, burnt, 70  
 caustic, 70  
 gas, 66  
 hydroxyd, 71  
 Limestone, 65  
 Litmus, 15  
 Lunge-Rohrmann plate column,  
 226  
 Lye, 77  
**M**AGNESIUM, 137  
 Manganese chlorid, 100  
 ore, 100  
 Marsh gas, 295, 301  
 series, 302  
 Mass units, 33  
 Mauvanilin, 321  
 Mercury, fulminate, 355  
 reactions of, 249  
 Metals, 17  
 Methane, 295  
 Methylanilin, 321  
 Microcosmic salt, 410  
 Molecular weight, 55, 126, 132  
 Molybdc test for phosphates, 411  
 Morphine, 209, 383  
 Mortar, 75  
 Mother liquor, 137

**N**APHTHALINE, 320  
 Nascent state, 186  
 Natural gas, 337  
 Nicotine, 209, 385  
 Niter, 152, 209, 211  
   plantation, 210  
 Nitrates, 164  
   test for, 178  
 Nitrites, 179  
 Nitro-benzol, 320  
   cellulose, 271  
   glycerin, 375  
 Nitrogen, 154  
   dioxyd, 171  
   monoxyd, 176  
   oxyds, 185  
   weight of, 13, 159  
 Nitrose, 223  
 Nitrous fumes, 170  
 Non-metals, 17

**O**IL, neat's foot, 390  
   of Dippel, 390  
   vitriol, 230

Oils, 370  
   drying, 377  
   essential, 371  
 Olein, 377  
 Orseille, 15  
 Orthophosphates, 409, 411  
 Oxybromids, 138  
   -iodids, 143  
 Oxyd, 16  
   copper, 16, 47  
   lead, 51  
 Oxygen, 15, 31  
   weight of, 13

**P**ALMITIN, 377  
 Papyrine, 270  
 Paraffin, 283, 337  
 Parchment, 270  
 Pearlash, 77  
 Peat, 309  
 Pentane, 304  
 Periodate, 143  
 Peroxyd of hydrogen, 34  
   sulfur, 47, 49  
 Petroleum, 335  
   ether, 337

Phenol, 282  
 Phosphids, 418  
 Phosphine, 415  
   reaction, 417  
 Phosphites, 414  
 Phosphonium, 417  
 Phosphorbronze, 418  
 Phosphorus, black, 407  
   pentachlorid, 415  
   pentoxyd, 407  
   red, 406  
   trichlorid, 414  
   trioxyd, 413  
   yellow, 405  
 Pitch, 380  
 Plants, structure of, 265  
 Platinum stills, 228  
 Polysulfids, 241  
 Potash, 77  
   bulb. Liebig's, 267  
   caustic, 79  
 Potassium, 84  
   chlorate, 105, 106  
   hydroxyd, 79  
   prussiate, red, 398  
   yellow, 391  
   reactions, 249  
 Propane, 303  
 Protoplasm, 209, 258  
 Prussian blue, 391  
 Putrefaction, 199  
 Pyrite, 219  
 Pyrogallol, 369  
 Pyrolusite, 100  
 Pyroxyline, 273

**Q**UININ, 384

**R**ADICAL, 124, 195  
 Rectification, 350  
 Resin, 378  
 Rock-oil, 335  
   salt, 92  
 Rosanilin, 322  
 Rosin, 379  
 Rubber, 380  
 Rules for the conversion of the  
   different thermometer degrees  
   into each other, 422

**SAFETY** lamp, Davy's, 300  
 Sal ammoniac, 189, 197, 204  
   soda, 121  
 Salt, 72, 92, 129  
   cake, 113  
   gas, 94, 106  
   of phosphorus, 410  
   seignette, 364  
 Saltpetre, 152  
 Saponification, 377  
 Sarkin, 388  
 Sericin, 389  
 Series, electrochemical, 125,  
   marsh gas, 302  
 Silk, 389  
 Silver chlorid, 111  
   fulminate, 357  
   reactions, 246  
 Skin, 388  
 Soap, 372, 377  
   soft, 76  
 Soda ash, 119, 121  
 Sodium, 116  
   carbonate, 121  
   reactions, 249  
 Solar oil, 281  
 Soldering, 205  
 Solvay process, 206  
 Spar, 65  
 Specific heats, 132  
 Spirits of niter, 155, 160  
   salt, 94, 106  
 Starch, 41, 340  
 Stearin, 376  
 Strychnine, 384  
 Sugar, 342  
   cane, 342  
   fruit, 344  
   grape, 343  
   milk, 345  
 Sulfid, 115  
 Sulfocyanates, 397  
 Sulfocyanogen, 397  
 Sulfur, 13  
   chlorid, 254  
   oxyd, 16, 17, 52, 59  
   peroxyd, 46, 49  
   test for, 40  
 Superoxyd, 117

**TABLE** for the comparison of  
 the scales of Réau-  
 mur's, Celsius's and  
 Fahrenheit's ther-  
 mometers, 421  
   of the liter weights of the  
   gases, 423  
 Tallow, 376  
 Tannin, 36, 366  
 Tar, coal, 320  
 Tartar emetic, 365  
 Theobromin, 383  
 Thermometers, rules for the con-  
   version of the  
   different degrees  
   of, into each  
   other, 422  
   table for the com-  
   parison of the  
   scales of Réau-  
   mur's, Celsius's,  
   and Fahrenheit's,  
   421

Thiosulfates, 243  
 Tin, reactions, 246  
 Toluidin, 323  
 Turpentine, 378

**UREA**, 199, 383

**VALENCE**, 125  
   Varec, 122  
 Vaseline, 337  
 Vermilion, 234  
 Vinegar, 15, 360  
 Vitriol, calcium, 73  
   copper, 46  
   green, 35, 36  
   hydrogen, 129  
   iron, 54, 58  
   lead, 51  
   oil of, 40, 43, 50, 230  
   silver, 49, 54  
   zinc, 54

Volume weights, 126  
 Vulcanizing, 382

**WATER**, 31  
   -gas, 331

Water, hard, 74  
  properties, 19  
Weights, atomic, 34, 126  
  molecular, 55, 126  
  volume, 126  
Wood distillation, 274  
  gases, 285  
  tar, 277, 279

Wool, 389

**X**ANTHINE, 388

**Z**INC, reactions, 246







# CATALOGUE

OF

## Practical and Scientific Books

PUBLISHED BY

# HENRY CAREY BAIRD & Co.

INDUSTRIAL PUBLISHERS, BOOKSELLERS AND IMPORTERS.

810 Walnut Street, Philadelphia.

~~Any~~ Any of the Books comprised in this Catalogue will be sent by mail, free of postage, to any address in the world, at the publication prices.

~~A~~ A Descriptive Catalogue, 90 pages, 8vo., will be sent free and free of postage, to any one in any part of the world, who will furnish his address.

~~Where~~ Where not otherwise stated, all of the Books in this Catalogue are bound in muslin.

### AMATEUR MECHANICS' WORKSHOP:

A treatise containing plain and concise directions for the manipulation of Wood and Metals, including Casting, Forging, Brazing, Soldering and Carpentry. By the author of the "Lathe and Its Uses." Seventh edition. Illustrated. 8vo. . . . \$2.50

### ANDES.—Animal Fats and Oils:

Their Practical Production, Purification and Uses; their Properties, Falsification and Examination. 62 illustrations. 8vo. . . \$4.00

### ANDES.—Vegetable Fats and Oils:

Their Practical Preparation, Purification and Employment; their Properties, Adulteration and Examination. 94 illustrations. 8vo. . . \$4.00

### ARLOT.—A Complete Guide for Coach Painters:

Translated from the French of M. ARLOT, Coach Painter, for eleven years Foreman of Painting to M. Eherler, Coach Maker, Paris. By A. A. FESQUET, Chemist and Engineer. To which is added an Appendix, containing Information respecting the Materials and the Practice of Coach and Car Painting and Varnishing in the United States and Great Britain. 12mo. . . . \$1.25

**ARMENGAUD, AMOROUX, AND JOHNSON.**—The Practical Draughtsman's Book of Industrial Design, and Machinist's and Engineer's Drawing Companion :

Forming a Complete Course of Mechanical Engineering and Architectural Drawing. From the French of M. Armengaud the elder, Prof. of Design in the Conservatoire of Arts and Industry, Paris, and MM. Armengaud the younger, and Amoroux, Civil Engineers. Rewritten and arranged with additional matter and plates, selections from and examples of the most useful and generally employed mechanism of the day. By WILLIAM JOHNSON, Assoc. Inst. C. E. Illustrated by fifty folio steel plates, and fifty wood-cuts. A new edition, 4to., cloth. . . . . \$6.00

**ARMSTRONG.**—The Construction and Management of Steam Boilers :

By R. ARMSTRONG, C. E. With an Appendix by ROBERT MALLET, C. E., F. R. S. Seventh Edition. Illustrated. 1 vol. 12mo. .60

**ARROWSMITH.**—The Paper-Hanger's Companion :

Comprising Tools, Pastes, Preparatory Work ; Selection and Hanging of Wall-Papers ; Distemper Painting and Cornice-Tinting ; Stencil Work ; Replacing Sash-Cord and Broken Window Panes ; and Useful Wrinkles and Receipts, By JAMES ARROWSMITH. A New, Thoroughly Revised, and Much Enlarged Edition. Illustrated by 25 engravings, 162 pages. (1905) . . . . . \$1.00

**ASHTON.**—The Theory and Practice of the Art of Designing Fancy Cotton and Woollen Cloths from Sample :

Giving full instructions for reducing drafts, as well as the methods of spooling and making out harness for cross drafts and finding any required reed ; with calculations and tables of yarn. By FREDERIC T. ASHTON, Designer, West Pittsfield, Mass. With fifty-two illustrations. One vol. folio . . . . . \$5.00

**ASKINSON.**—Perfumes and their Preparation :

A Comprehensive Treatise on Perfumery, containing Complete Directions for Making Handkerchief Perfumes, Smelling-Salts, Sachets, Fumigating Pastils ; Preparations for the Care of the Skin, the Mouth, the Hair ; Cosmetics, Hair Dyes, and other Toilet Articles. By G.W. ASKINSON. Translated from the German by ISIDOR FURST. Revised by CHARLES RICE. 32 Illustrations. 8vo. \$3.00

**BRONGNIART.**—Coloring and Decoration of Ceramic Ware.

8vo. . . . . \$2.50

**BAIRD.**—The American Cotton Spinner, and Manager's and Carder's Guide :

A Practical Treatise on Cotton Spinning ; giving the Dimensions and Speed of Machinery, Draught and Twist Calculations, etc. ; with notices of recent Improvements : together with Rules and Examples for making changes in the sizes and numbers of Roving and Yarn. Compiled from the papers of the late ROBERT H. BAIRD. 12mo. . . . . \$1.50

**BAKER.—Long-Span Railway Bridges :**

Comprising Investigations of the Comparative Theoretical and Practical Advantages of the various Adopted or Proposed Type Systems of Construction ; with numerous Formulæ and Tables. By

B. BAKER. 12mo. . . . . \$1.00

**BRANNT.—A Practical Treatise on Distillation and Rectification of Alcohol :**

Comprising Raw Materials ; Production of Malt, Preparation of Mashies and of Yeast ; Fermentation ; Distillation and Rectification and Purification of Alcohol ; Preparation of Alcoholic Liquors, Liqueurs, Cordials, Bitters, Fruit Essences, Vinegar, etc.; Examination of Materials for the Preparation of Malt as well as of the Malt itself ; Examination of Mashies before and after Fermentation ; Alcoholometry, with Numerous Comprehensive Tables ; and an Appendix on the Manufacture of Compressed Yeast and the Examination of Alcohol and Alcoholic Liquors for Fusel Oil and other Impurities. By WILLIAM T. BRANNT, Editor of "The Techno-Chemical Receipt Book." Second Edition. Entirely Rewritten. Illustrated by 105

engravings. 460 pages, 8vo. (Dec., 1903) . . . . . \$4.00

**BAKRR.—A Practical Treatise on the Combustion of Coal :**

Including descriptions of various mechanical devices for the Economic Generation of Heat by the Combustion of Fuel, whether solid, liquid or gaseous. 8vo. . . . . \$2.50

**BARR.—A Practical Treatise on High Pressure Steam Boilers:**

Including Results of Recent Experimental Tests of Boiler Materials, together with a description of Approved Safety Apparatus, Steam Pumps, Injectors and Economizers in actual use. By WM. M. BARR. 204 Illustrations. 8vo. . . . . \$3.00

**BAUERMAN.—A Treatise on the Metallurgy of Iron :**

Containing Outlines of the History of Iron Manufacture, Methods of Assay, and Analysis of Iron Ores, Processes of Manufacture of Iron and Steel, etc., etc. By H. BAUERMAN, F. G. S., Associate of the Royal School of Mines. Fifth Edition, Revised and Enlarged. Illustrated with numerous Wood Engravings from Drawings by J. B. JORDAN. 12mo, . . . . . \$2.00

**BRANNT.—The Metallic Alloys : A Practical Guide**

For the Manufacture of all kinds of Alloys, Amalgams, and Solders, used by Metal-Workers : together with their Chemical and Physical Properties and their Application in the Arts and the Industries ; with an Appendix on the Coloring of Alloys and the Recovery of Waste Metals. By WILLIAM T. BRANNT. 45 Engravings. Third, Revised, and Enlarged Edition. 570 pages. 8vo. . . . . Net, \$5.00

**BRANNT.—The Soap Maker's Hand-Book of Materials, Processes and Receipts for Every Description of Soap. Illustrated. 8vo. (In preparation.)****BEANS.—A Treatise on Railway Curves and Location of Railroads :**

By E. W. BEANS, C. E. Illustrated. 12mo. Tucks. . . . . \$1.50

**BELL.—Carpentry Made Easy :**

Or, The Science and Art of Framing on a New and Improved System. With Specific Instructions for Building Balloon Frames, Barn Frames, Mill Frames, Warehouses, Church Spires, etc. Comprising also a System of Bridge Building, with Bills, Estimates of Cost, and valuable Tables. Illustrated by forty-four plates, comprising nearly 200 figures. By WILLIAM E. BELL, Architect and Practical Builder. 8vo. \$5.00

**BEMROSE.—Fret-Cutting and Perforated Carving :**

With fifty-three practical illustrations. By W. BEMROSE, JR. 1 vol. quarto \$2.50

**BEMROSE.—Manual of Buhl-work and Marquetry :**

With Practical Instructions for Learners, and ninety colored designs. By W. BEMROSE, JR. 1 vol. quarto \$3.00

**BEMROSE.—Manual of Wood Carving :**

With Practical Illustrations for Learners of the Art, and Original and Selected Designs. By WILLIAM BEMROSE, JR. With an Introduction by LLEWELLYN JEWITT, F. S. A., etc. With 128 illustrations, 4to. \$2.50

**BERSCH.—Cellulose, Cellulose Products, and Rubber Substitutes :**

Comprising the Preparation of Cellulose, Parchment-Cellulose, Methods of Obtaining Sugar, Alcohol and Oxalic Acid from Wood-Cellulose ; Production of Nitro-Cellulose and Cellulose Esters ; Manufacture of Artificial Silk, Viscose, Celluloid, Rubber Substitutes, Oil-Rubber, and Faktis. By DR. JOSEPH BERSCH. Translated by WILLIAM T. BRANNT. 41 illustrations. (1904.) \$3.00

**BILLINGS.—Tobacco :**

Its History, Variety, Culture, Manufacture, Commerce, and Various Modes of Use. By E. R. BILLINGS. Illustrated by nearly 200 engravings. 8vo. \$3.00

**BIRD.—The American Practical Dyers' Companion :**

Comprising a Description of the Principal Dye-Stuffs and Chemicals used in Dyeing, their Natures and Uses ; Mordants and How Made ; with the best American, English, French and German processes for Bleaching and Dyeing Silk, Wool, Cotton, Linen, Flannel, Felt, Dress Goods, Mixed and Hosiery Yarns, Feathers, Grass, Felt, Fur, Wool, and Straw Hats, Jute Yarn, Vegetable Ivory, Mats, Skins, Furs, Leather, etc., etc. By Wood Aniline, and other Processes, together with Remarks on Finishing Agents, and instructions in the Finishing of Fabrics, Substitutes for Indigo, Water-Proofing of Materials, Tests and Purification of Water, Manufacture of Aniline and other New Dye Wares, Harmonizing Colors, etc., etc. ; embracing in all over 800 Receipts for Colors and Shades, accompanied by 170 Dyed Samples of Raw Materials and Fabrics. By F. J. BIRD, Practical Dyer, Author of "The Dyers' Hand-Book." 8vo. \$5.00



**BLINN.—A Practical Workshop Companion for Tin, Sheet-Iron, and Copper-plate Workers:**

Containing Rules for describing various kinds of Patterns used by Tin, Sheet-Iron and Copper-plate Workers; Practical Geometry; Mensuration of Surfaces and Solids; Tables of the Weights of Metals, Lead-pipe, etc.; Tables of Areas and Circumferences of Circles; Japan, Varnishes, Lackers, Cements, Compositions, etc., etc. By LEROY J. BLINN, Master Mechanic. With One Hundred and Seventy Illustrations. 12mo. . . . . \$2.50

**BOOTH.—Marble Worker's Manual:**

Containing Practical Information respecting Marbles in general, their Cutting, Working and Polishing; Veneering of Marble; Mosaics; Composition and Use of Artificial Marble, Stuccos, Cements, Receipts, Secrets, etc., etc. Translated from the French by M. L. BOOTH. With an Appendix concerning American Marbles. 12mo., cloth \$1.50

**BRANNT.—A Practical Treatise on Animal and Vegetable Fats and Oils:**

Comprising both Fixed and Volatile Oils, their Physical and Chemical Properties and Uses, the Manner of Extracting and Refining them, and Practical Rules for Testing them; as well as the Manufacture of Artificial Butter and Lubricants, etc., with lists of American Patents relating to the Extraction, Rendering, Refining, Decomposing, and Bleaching of Fats and Oils. By WILLIAM T. BRANNT, Editor of the "Techno-Chemical Receipt Book." Second Edition, Revised and in a great part Rewritten. Illustrated by 302 Engravings. In Two Volumes. 1304 pp. 8vo. . . . . \$10.00

**BRANNT.—A Practical Treatise on the Manufacture of Soap and Candles:**

Based upon the most Recent Experiences in the Practice and Science; comprising the Chemistry, Raw Materials, Machinery, and Utensils and Various Processes of Manufacture, including a great variety of formulas. Edited chiefly from the German of Dr. C. Deite, A. Engelhardt, Dr. C. Schaedler and others; with additions and lists of American Patents relating to these subjects. By WM. T. BRANNT. Illustrated by 163 engravings. 677 pages. 8vo. . . . \$12.50

**BRANNT.—India Rubber, Gutta-Percha and Balata:**

Occurrence, Geographical Distribution, and Cultivation, Obtaining and Preparing the Raw Materials, Modes of Working and Utilizing them, Including Washing, Maceration, Mixing, Vulcanizing, Rubber and Gutta-Percha Compounds, Utilization of Waste, etc. By WILLIAM T. BRANNT. Illustrated. 12mo. (1900.) . . . \$3.00

**BRANNT—WAHL.—The Techno-Chemical Receipt Book :**

Containing several thousand Receipts covering the latest, most important, and most useful discoveries in Chemical Technology, and their Practical Application in the Arts and the Industries. Edited chiefly from the German of Drs. Winckler, Elsner, Heintze, Mierzinski, Jacobsen, Koller and Heinzerling, with additions by WM. T. BRANNT and WM. H. WAHL, Ph. D. Illustrated by 78 engravings. 12mo. 495 pages. . . . . \$2.00

**BROWN.—Five Hundred and Seven Mechanical Movements :**

Embracing all those which are most important in Dynamics, Hydraulics, Hydrostatics, Pneumatics, Steam Engines, Mill and other Gearing, Presses, Horology, and Miscellaneous Machinery ; and including many movements never before published, and several of which have only recently come into use. By HENRY T. BROWN. 12mo. . . . . \$1.00

**BUCKMASTER.—The Elements of Mechanical Physics :**

By J. C. BUCKMASTER. Illustrated with numerous engravings. 12mo. . . . . \$1.00

**BULLOCK.—The American Cottage Builder :**

A Series of Designs, Plans and Specifications, from \$200 to \$20,000, for Homes for the People ; together with Warming, Ventilation, Drainage, Painting and Landscape Gardening. By JOHN BULLOCK, Architect and Editor of "The Rudiments of Architecture and Building," etc., etc. Illustrated by 75 engravings. 8vo.

**BULLOCK.—The Rudiments of Architecture and Building :**

For the use of Architects, Builders, Draughtsmen, Machinists, Engineers and Mechanics. Edited by JOHN BULLOCK, author of "The American Cottage Builder." Illustrated by 250 Engravings. 8vo. \$2.50

**BURGH.—Practical Rules for the Proportions of Modern Engines and Boilers for Land and Marine Purposes.**

By N. P. BURGH, Engineer. 12mo. . . . . \$1.50

**BYLES.—Sophisms of Free Trade and Popular Political Economy Examined.**

By a BARRISTER (SIR JOHN BARNARD BYLES, Judge of Common Pleas). From the Ninth English Edition, as published by the Manchester Reciprocity Association. 12mo. . . . . \$1.25

**BOWMAN.—The Structure of the Wool Fibre in its Relation to the Use of Wool for Technical Purposes :**

Being the substance, with additions, of Five Lectures, delivered at the request of the Council, to the members of the Bradford Technical College, and the Society of Dyers and Colorists. By F. H. BOWMAN, D. Sc., F. R. S. E., F. L. S. Illustrated by 32 engravings. 8vo. . . . . \$7.50

**BYRNE.—Hand-Book for the Artisan, Mechanic, and Engineer :**

Comprising the Grinding and Sharpening of Cutting Tools, Abrasive Processes, Lapidary Work, Gem and Glass Engraving, Varnishing and Lackering, Apparatus, Materials and Processes for Grinding and



Polishing, etc. By OLIVER BYRNE. Illustrated by 185 wood engravings. 8vo. \$5.00

**BYRNE.—Pocket-Book for Railroad and Civil Engineers:**

Containing New, Exact and Concise Methods for Laying out Railroad Curves, Switches, Frog Angles and Crossings; the Staking out of work; Levelling; the Calculation of Cuttings: Embankments; Earth-work, etc. By OLIVER BYRNE. 18mo., full bound, pocket-book form \$1.50

**BYRNE.—The Practical Metal-Worker's Assistant:**

Comprising Metallurgic Chemistry; the Arts of Working all Metals and Alloys; Forging of Iron and Steel; Hardening and Tempering; Melting and Mixing; Casting and Founding; Works in Sheet Metal; the Processes Dependent on the Ductility of the Metals; Soldering; and the most Improved Processes and Tools employed by Metal-Workers. With the Application of the Art of Electro-Metallurgy to Manufacturing Processes; collected from Original Sources, and from the works of Holtzapffel, Bergeron, Leupold, Plumier, Napier, Scoffern, Clay, Fairbairn and others. By OLIVER BYRNE. A new, revised and improved edition, to which is added an Appendix, containing The Manufacture of Russian Sheet-Iron. By JOHN PERCY, M. D., F. R. S. The Manufacture of Malleable Iron Castings, and Improvements in Bessemer Steel. By A. A. FESQUET, Chemist and Engineer. With over Six Hundred Engravings, Illustrating every Branch of the Subject. 8vo. \$5.00

**BYRNE.—The Practical Model Calculator:**

For the Engineer, Mechanic, Manufacturer of Engine Work, Naval Architect, Miner and Millwright. By OLIVER BYRNE. 8vo., nearly 600 pages (Scarce.)

**CABINET MAKER'S ALBUM OF FURNITURE:**

Comprising a Collection of Designs for various Styles of Furniture. Illustrated by Forty-eight Large and Beautifully Engraved Plates. Oblong, 8vo. \$1.50

**CALLINGHAM.—Sign Writing and Glass Embossing:**

A Complete Practical Illustrated Manual of the Art. By JAMES CALLINGHAM. To which are added Numerous Alphabets and the Art of Letter Painting Made Easy. By JAMES C. BADENOCH. 258 pages. 12mo. \$1.50

**CAMPIN.—A Practical Treatise on Mechanical Engineering:**

Comprising Metallurgy, Moulding, Casting, Forging, Tools, Workshop Machinery, Mechanical Manipulation, Manufacture of Steam-Engines, etc. With an Appendix on the Analysis of Iron and Iron Ores. By FRANCIS CAMPIN, C. E. To which are added, Observations on the Construction of Steam Boilers, and Remarks upon Furnaces used for Smoke Prevention; with a Chapter on Explosions. By R. ARMSTRONG, C. E., and JOHN BOURNE. (Scarce.)

**CAREY.—A Memoir of Henry C. Carey.**

By DR. WM. ELDER. With a portrait. 8vo., cloth . . . 75

**CAREY.—The Works of Henry C. Carey:**

Harmony of Interests: Agricultural, Manufacturing and Commercial. 8vo. . . . \$1.25

Manual of Social Science. Condensed from Carey's "Principles of Social Science." By KATE MCKEAN. 1 vol. 12mo. . . \$2.00

Miscellaneous Works. With a Portrait. 2 vols. 8vo. . . \$10.00

Past, Present and Future. 8vo. . . . \$2.50

Principles of Social Science. 3 volumes, 8vo. . . \$10.00

The Slave-Trade, Domestic and Foreign; Why it Exists, and How it may be Extinguished (1853). 8vo. . . . \$2.00

The Unity of Law: As Exhibited in the Relations of Physical, Social, Mental and Moral Science (1872). 8vo. . . . \$2.50

**CLARK.—Tramways, their Construction and Working:**

Embracing a Comprehensive History of the System. With an exhaustive analysis of the various modes of traction, including horse-power, steam, heated water and compressed air; a description of the varieties of Rolling stock, and ample details of cost and working expenses. By D. KINNEAR CLARK. Illustrated by over 200 wood engravings, and thirteen folding plates. 1 vol. 8vo. . . \$5.00

**COLBURN.—The Locomotive Engine:**

Including a Description of its Structure, Rules for Estimating its Capabilities, and Practical Observations on its Construction and Management. By ZERAH COLBURN. Illustrated. 12mo. . . \$1.00

**COLLENS.—The Eden of Labor; or, the Christian Utopia.**

By T. WHARTON COLLENS, author of "Humanics," "The History of Charity," etc. 12mo. Paper cover, \$1.00; Cloth . . \$1.25

**COOLEY.—A Complete Practical Treatise on Perfumery:**

Being a Hand-book of Perfumes, Cosmetics and other Toilet Articles With a Comprehensive Collection of Formulæ. By ARNOLD COOLEY. 12mo. . . . \$1.50

**COOPER.—A Treatise on the use of Belting for the Transmission of Power.**

With numerous illustrations of approved and actual methods of arranging Main Driving and Quarter Twist Belts, and of Belt Fastenings. Examples and Rules in great number for exhibiting and calculating the size and driving power of Belts. Plain, Particular and Practical Directions for the Treatment, Care and Management of Belts. Descriptions of many varieties of Beltings, together with chapters on the Transmission of Power by Ropes; by Iron and Wood Frictional Gearing; on the Strength of Belting Leather; and on the Experimental Investigations of Morin, Briggs, and others. By JOHN H. COOPER, M. E. 8vo. . . . \$3.50

**CRAIK.—The Practical American Millwright and Miller.**

By DAVID CRAIK, Millwright. Illustrated by numerous wood engravings and two folding plates. 8vo. . . . (Scarce.)

**CROSS.—The Cotton Yarn Spinner :**

Showing how the Preparation should be arranged for Different Counts of Yarns by a System more uniform than has hitherto been practiced; by having a Standard Schedule from which we make all our Changes. By RICHARD CROSS. 122 pp. 12mo. . 75

**CRISTIANI.—A Technical Treatise on Soap and Candles:**

With a Glance at the Industry of Fats and Oils. By R. S. CRISTIANI, Chemist. Author of "Perfumery and Kindred Arts." Illustrated by 176 engravings. 581 pages, 8vo. \$15.00

**COURTNEY.—The Boiler Maker's Assistant in Drawing, Templating, and Calculating Boiler Work and Tank Work, etc.**

Revised by D. K. CLARK. 102 ills. Fifth edition. . . 80

**COURTNEY.—The Boiler Maker's Ready Reckoner:**

With Examples of Practical Geometry and Templating. Revised by D. K. CLARK, C. E. 37 illustrations. Fifth edition. . \$1.60

**DAVIDSON.—A Practical Manual of House Painting, Grain-ing, Marbling, and Sign-Writing:**

Containing full information on the processes of House Painting in Oil and Distemper, the Formation of Letters and Practice of Sign-Writing, the Principles of Decorative Art, a Course of Elementary Drawing for House Painters, Writers, etc., and a Collection of Useful Receipts. With nine colored illustrations of Woods and Marbles, and numerous wood engravings. By ELLIS A. DAVIDSON. 12mo. \$2.00

**DAVIES.—A Treatise on Earthy and Other Minerals and Mining:**

By D. C. DAVIES, F. G. S., Mining Engineer, etc. Illustrated by 76 Engravings. 12mo. . . \$5.00

**DAVIES.—A Treatise on Metalliferous Minerals and Mining:**

By D. C. DAVIES, F. G. S., Mining Engineer, Examiner of Mines, Quarries and Collieries. Illustrated by 148 engravings of Geological Formations, Mining Operations and Machinery, drawn from the practice of all parts of the world. Fifth Edition, thoroughly Revised and much Enlarged by his son, E. Henry Davies. 12mo., 524 pages . . . \$5.00

**DIETERICHS.—A Treatise on Friction, Lubrication, Oils and Fats:**

The Manufacture of Lubricating Oils, Paint Oils, and of Grease, and the Testing of Oils. By E. F. DIETERICHS, Member of the Franklin Institute; Member National Association of Stationary Engineers; Inventor of Dieterichs' Valve-Oleum Lubricating Oils. 12mo. (1906.) *A practical book by a practical man.* . . \$1.25

**DAVIS.—A Practical Treatise on the Manufacture of Brick, Tiles and Terra-Cotta:**

Including Stiff Clay, Dry Clay, Hand Made, Pressed or Front, and Roadway Paving Brick, Enamelled Brick, with Glazes and Colors, Fire Brick and Blocks, Silica Brick, Carbon Brick, Glass Pots, Re-

torts, Architectural Terra-Cotta, Sewer Pipe, Drain Tile, Glazed and Unglazed Roofing Tile, Art Tile, Mosaics, and Imitation of Intarsia or Inlaid Surfaces. Comprising every product of Clay employed in Architecture, Engineering, and the Blast Furnace. With a Detailed Description of the Different Clays employed, the Most Modern Machinery, Tools, and Kilns used, and the Processes for Handling, Disintegrating, Tempering, and Moulding the Clay into Shape, Drying, Setting, and Burning. By CHARLES THOMAS DAVIS. Third Edition. Revised and in great part rewritten. Illustrated by 261 engravings. 662 pages . . . . . \$20.00

**DAVIS.—A Treatise on Steam-Boiler Incrustation and Methods for Preventing Corrosion and the Formation of Scale:**  
By CHARLES T. DAVIS. Illustrated by 65 engravings. 8vo.

**DAVIS.—The Manufacture of Paper:**

Being a Description of the various Processes for the Fabrication, Coloring and Finishing of every kind of Paper, Including the Different Raw Materials and the Methods for Determining their Values, the Tools, Machines and Practical Details connected with an intelligent and a profitable prosecution of the art, with special reference to the best American Practice. To which are added a History of Paper, complete Lists of Paper-Making Materials, List of American Machines, Tools and Processes used in treating the Raw Materials, and in Making, Coloring and Finishing Paper. By CHARLES T. DAVIS. Illustrated by 156 engravings. 608 pages, 8vo. \$6.00

**DAVIS.—The Manufacture of Leather:**

Being a Description of all the Processes for the Tanning and Tawing with Bark, Extracts, Chrome and all Modern Tannages in General Use, and the Currying, Finishing and Dyeing of Every Kind of Leather; Including the Various Raw Materials, the Tools, Machines, and all Details of Importance Connected with an Intelligent and Profitable Prosecution of the Art, with Special Reference to the Best American Practice. To which are added Lists of American Patents (1884-1897) for Materials, Processes, Tools and Machines for Tanning, Currying, etc. By CHARLES THOMAS DAVIS. Second Edition, Revised, and in great part Rewritten. Illustrated by 147 engravings and 14 Samples of Quebracho Tanned and Aniline Dyed Leathers. 8vo, cloth, 712 pages. Price . . . . . \$12.50

**DAWIDOWSKY—BRANNT.—A Practical Treatise on the Raw Materials and Fabrication of Glue, Gelatine, Gelatine Veneers and Foils, Isinglass, Cements, Pastes, Mucilages, etc.:**

Based upon Actual Experience. By F. DAWIDOWSKY, Technical Chemist. Translated from the German, with extensive additions, including a description of the most Recent American Processes, by WILLIAM T. BRANNT. 2d revised edition, 350 pages. (1905.) Price . . . . . \$3.00

**DE GRAFF.—The Geometrical Stair-Builders' Guide:**

Being a Plain Practical System of Hand-Railing, embracing all its necessary Details, and Geometrically Illustrated by twenty-two Steel Engravings; together with the use of the most approved principles of Practical Geometry By SIMON DE GRAFF, Architect (See . . .)



**DE KONINCK—DIETZ.—A Practical Manual of Chemical Analysis and Assaying:**

As applied to the Manufacture of Iron from its Ores, and to Cast Iron, Wrought Iron, and Steel, as found in Commerce. By L. L. DE KONINCK, Dr. Sc., and E. DIETZ, Engineer. Edited with Notes, by ROBERT MALLET, F. R. S., F. S. G., M. I. C. E., etc. American Edition, Edited with Notes and an Appendix on Iron Ores, by A. A. FESQUET, Chemist and Engineer. 12mo. . . . \$1.00

**DUNCAN.—Practical Surveyor's Guide:**

Containing the necessary information to make any person of common capacity, a finished land surveyor without the aid of a teacher. By ANDREW DUNCAN. Revised. 72 engravings, 214 pp. 12mo. \$1.50

**DUPLAIS.—A Treatise on the Manufacture and Distillation of Alcoholic Liquors:**

Comprising Accurate and Complete Details in Regard to Alcohol from Wine, Molasses, Beets, Grain, Rice, Potatoes, Sorghum, Asphodel, Fruits, etc.; with the Distillation and Rectification of Brandy, Whiskey, Rum, Gin, Swiss Absinthe, etc., the Preparation of Aromatic Waters, Volatile Oils or Essences, Sugars, Syrups, Aromatic Tinctures, Liqueurs, Cordial Wines, Effervescing Wines, etc., the Ageing of Brandy and the improvement of Spirits, with Copious Directions and Tables for Testing and Reducing Spirituous Liquors, etc., etc. Translated and Edited from the French of MM. DUPLAIS, By M. MCKENNIE, M. D. Illustrated. 743 pp. 8vo. \$15.00

**DYER AND COLOR-MAKER'S COMPANION:**

Containing upwards of two hundred Receipts for making Colors, on the most approved principles, for all the various styles and fabrics now in existence; with the Scouring Process, and plain Directions for Preparing, Washing-off, and Finishing the Goods. 12mo. \$1.00

**EIDHERR.—The Techno-Chemical Guide to Distillation:**

A Hand-Book for the Manufacture of Alcohol and Alcoholic Liquors, including the Preparation of Malt and Compressed Yeast. Edited from the German of Ed. Eidherr.

**EDWARDS.—A Catechism of the Marine Steam-Engine,**

For the use of Engineers, Firemen, and Mechanics. A Practical Work for Practical Men. By EMORY EDWARDS, Mechanical Engineer. Illustrated by sixty-three Engravings, including examples of the most modern Engines. Third edition, thoroughly revised, with much additional matter. 12mo. 414 pages. \$2.00

**EDWARDS.—Modern American Locomotive Engines,**

Their Design, Construction and Management. By EMORY EDWARDS. Illustrated 12mo. . . . . \$2.00

**EDWARDS.—The American Steam Engineer:**

Theoretical and Practical, with examples of the latest and most approved American practice in the design and construction of Steam Engines and Boilers. For the use of engineers, machinists, boiler-makers, and engineering students. By EMORY EDWARDS. Fully Illustrated, 419 pages. 12mo. . . . . \$2.00

**EDWARDS.—Modern American Marine Engines, Boilers, and Screw Propellers,**

Their Design and Construction. Showing the Present Practice of the most Eminent Engineers and Marine Engine Builders in the United States. Illustrated by 30 large and elaborate plates. 4to. \$3.00

**EDWARDS.—The Practical Steam Engineer's Guide**

In the Design, Construction, and Management of American Stationary, Portable, and Steam Fire-Engines, Steam Pumps, Boilers. Injectors, Governors, Indicators, Pistons and Rings, Safety Valves and Steam Gauges. For the use of Engineers, Firemen, and Steam Users. By EMORY EDWARDS. Illustrated by 119 engravings. 420 pages. 12mo. . . . . \$2.00

**EISSLER.—The Metallurgy of Silver :**

A Practical Treatise on the Amalgamation, Roasting, and Lixiviation of Silver Ores, including the Assaying, Melting, and Refining of Silver Bullion. By M. EISSLER. 124 Illustrations. 336 pp. 12mo. . . . . \$4.25

**ELDER.—Conversations on the Principal Subjects of Political Economy.**

By DR. WILLIAM ELDER. 8vo. . . . . \$2.00

**ELDER.—Questions of the Day,**

Economic and Social. By DR. WILLIAM ELDER. 8vo. . . \$3.00

**ERNI AND BROWN.—Mineralogy Simplified.**

Easy Methods of Identifying Minerals, including Ores, by Means of the Blow-pipe, by Flame Reactions, by Humid Chemical Analysis, and by Physical Tests. By HENRI ERNI, A. M., M. D. Third Edition, revised, re-arranged and with the addition of entirely new matter, including Tables for the Determination of Minerals by Chemical and Pyrognostic Characters, and by Physical Characters. By AMOS P. BROWN, E. M., Ph. D. 350 pp., illustrated by 96 engravings, pocket-book form, full flexible morocco, gilt edges . . . . . \$2.50

**FAIRBAIRN. —The Principles of Mechanism and Machinery of Transmission :**

Comprising the Principles of Mechanism, Wheels, and Pulleys, Strength and Proportion of Shafts, Coupling of Shafts, and Engaging and Disengaging Gear. By SIR WILLIAM FAIRBAIRN, Bart. C. E. Beautifully illustrated by over 150 wood-cuts. In one volume, 12mo. . . . . \$2.00

**FLEMING.—Narrow Gauge Railways in America :**

A Sketch of their Rise, Progress, and Success. Valuable Statistics as to Grades, Curves, Weight of Rail, Locomotives, Cars, etc. By HOWARD FLEMING. Illustrated, 8vo. . . . . \$1.00

**FORSYTH.—Book of Designs for Headstones, Mural, and other Monuments :**

Containing 78 Designs. By JAMES FORSYTH, With an Introduction by CHARLES BOUTELL, M. A. 4to., cloth . . . . . \$3.50

**FRIEDBERG. Utilization of Bones by Chemical Means; especially the Modes of Obtaining Fat, Glue, Manures, Phosphorus and Phosphates.**

Illustrated. 8vo. (In preparation.)



**FRANKEL—HUTTER.—A Practical Treatise on the Manufacture of Starch, Glucose, Starch-Sugar, and Dextrine:**

Based on the German of LADISLAUS VON WAGNER, Professor in the Royal Technical High School, Buda-Pest, Hungary, and other authorities. By JULIUS FRANKEL, Graduate of the Polytechnic School of Hanover. Edited by ROBERT HUTTER, Chemist, Practical Manufacturer of Starch-Sugar. Illustrated by 58 engravings, covering every branch of the subject, including examples of the most Recent and Best American Machinery. 8vo., 344 pp. \$6.00

**GARDNER.—The Painter's Encyclopædia:**

Containing Definitions of all Important Words in the Art of Plain and Artistic Painting, with Details of Practice in Coach, Carriage, Railway Car, House, Sign, and Ornamental Painting, including Graining, Marbling, Staining, Varnishing, Polishing, Lettering, Stenciling, Gilding, Bronzing, etc. By FRANKLIN B. GARDNER. 158 Illustrations. 12mo. 427 pp. \$2.00

**GARDNER.—Everybody's Paint Book:**

A Complete Guide to the Art of Outdoor and Indoor Painting. 38 illustrations. 12mo, 183 pp. \$1.00

**GEE.—The Jeweller's Assistant in the Art of Working in Gold:**

A Practical Treatise for Masters and Workmen. 12mo. \$3.00

**GEE.—The Goldsmith's Handbook:**

Containing full instructions for the Alloying and Working of Gold, including the Art of Alloying, Melting, Reducing, Coloring, Collecting, and Refining; the Processes of Manipulation, Recovery of Waste; Chemical and Physical Properties of Gold; with a New System of Mixing its Alloys; Solders, Enamels, and other Useful Rules and Recipes. By GEORGE E. GEE. 12mo. \$1.25

**GEE.—The Silversmith's Handbook:**

Containing full instructions for the Alloying and Working of Silver, including the different modes of Refining and Melting the Metal; its Solders; the Preparation of Imitation Alloys; Methods of Manipulation; Prevention of Waste; Instructions for Improving and Finishing the Surface of the Work; together with other Useful Information and Memoranda. By GEORGE E. GEE. Illustrated. 12mo. \$1.25

**GOTHIC ALBUM FOR CABINET-MAKERS:**

Designs for Gothic Furniture. Twenty-three plates. Oblong \$1.50

**GRANT.—A Handbook on the Teeth of Gears:**

Their Curves, Properties, and Practical Construction. By GEORGE B. GRANT. Illustrated. Third Edition, enlarged. 8vo. \$1.00

**GREENWOOD.—Steel and Iron:**

Comprising the Practice and Theory of the Several Methods Pursued in their Manufacture, and of their Treatment in the Rolling-Mills, the Forge, and the Foundry. By WILLIAM HENRY GREENWOOD, F. C. S. With 97 Diagrams, 536 pages. 12mo. \$1.75

**GREGORY.—Mathematics for Practical Men:**

Adapted to the Pursuits of Surveyors, Architects, Mechanics, and Civil Engineers. By OLINTHUS GREGORY. 8vo., plates \$3.00

**GRISWOLD.—Railroad Engineer's Pocket Companion for the Field:**

Comprising Rules for Calculating Deflection Distances and Angles, Tangential Distances and Angles, and all Necessary Tables for Engineers; also the Art of Levelling from Preliminary Survey to the Construction of Railroads, intended Expressly for the Young Engineer, together with Numerous Valuable Rules and Examples. By W. GRISWOLD. 12mo., tucks . . . . . \$1.50

**GRUNER.—Studies of Blast Furnace Phenomena:**

By M. L. GRUNER, President of the General Council of Mines of France, and lately Professor of Metallurgy at the Ecole des Mines Translated, with the author's sanction, with an Appendix, by L. D. B. GORDON, F. R. S. E., F. G. S. 8vo. . . . . \$2.50

**Hand-Book of Useful Tables for the Lumberman, Farmer and Mechanic:**

Containing Accurate Tables of Logs Reduced to Inch Board Measure, Plank, Scantling and Timber Measure; Wages and Rent, by Week or Month; Capacity of Granaries, Bins and Cisterns; Land Measure, Interest Tables, with Directions for Finding the Interest on any sum at 4, 5, 6, 7 and 8 per cent., and many other Useful Tables. 32 mo., boards. 186 pages . . . . . .25

**HASERICK.—The Secrets of the Art of Dyeing Wool, Cotton, and Linen,**

Including Bleaching and Coloring Wool and Cotton Hosiery and Random Yarns. A Treatise based on Economy and Practice. By E. C. HASERICK. *Illustrated by 323 Dyed Patterns of the Yarns or Fabrics.* 8vo. . . . . \$5.00

**HATS AND FELTING:**

A Practical Treatise on their Manufacture. By a Practical Hatter. Illustrated by Drawings of Machinery, etc. 8vo. . . . \$1.00

**HERMANN.—Painting on Glass and Porcelain, and Enamel Painting:**

A Complete Introduction to the Preparation of all the Colors and Fluxes Used for Painting on Glass, Porcelain, Enamel, Faience and Stoneware, the Color Pastes and Colored Glasses, together with a Minute Description of the Firing of Colors and Enamels, on the Basis of Personal Practical Experience of the Art up to Date. 18 illustrations. Second edition. . . . . \$4.00

**HAUPT.—Street Railway Motors:**

With Descriptions and Cost of Plants and Operation of the Various Systems now in Use. 12mo. . . . . \$1.75

**HAUPT.—A Manual of Engineering Specifications and Contracts.**

By LEWIS M. HAUPT, C. E. Illustrated with numerous maps.  
328pp. 8vo. . . . . \$3 00

**HAUPT.—The Topographer, His Instruments and Methods.**

By LEWIS M. HAUPT, A. M., C. E. Illustrated with numerous  
plates, maps and engravings. 247 pp. 8vo. . . . . \$3.00

**HUGHES.—American Miller and Millwright's Assistant:**

By WILLIAM CARTER HUGHES. 12mo. . . . . \$1.50

**HULME.—Worked Examination Questions in Plane Geometrical Drawing:**

For the Use of Candidates for the Royal Military Academy, Woolwich; the Royal Military College, Sandhurst; the Indian Civil Engineering College, Cooper's Hill; Indian Public Works and Telegraph Departments; Royal Marine Light Infantry; the Oxford and Cambridge Local Examinations, etc. By F. EDWARD HULME, F. L. S., F. S. A., Art-Master Marlborough College. Illustrated by 300 examples. Small quarto . . . . . \$1.00

**JERVIS.—Railroad Property:**

A Treatise on the Construction and Management of Railways; designed to afford useful knowledge, in the popular style, to the holders of this class of property; as well as Railway Managers, Officers, and Agents. By JOHN B. JERVIS, late Civil Engineer of the Hudson River Railroad, Croton Aqueduct, etc. 12mo., cloth \$1.50

**KEENE.—A Hand-Book of Practical Gauging:**

For the Use of Beginners, to which is added a Chapter on Distillation, describing the process in operation at the Custom-House for ascertaining the Strength of Wines. By JAMES B. KEENE, of H. M. Customs. 8vo. . . . . \$1 00

**KELLEY.—Speeches, Addresses, and Letters on Industrial and Financial Questions:**

By HON. WILLIAM D. KELLEY, M. C. 544 pages, 8vo. . . . \$2.50

**KOENIG.—Chemistry Simplified:**

A Course of Lectures on the Non-Metals Based upon the Natural Evolution of Chemistry. Designed Primarily for Engineers. By GEORGE AUGUSTUS KOENIG, Ph. D., A. M., E. M., Professor of Chemistry, Michigan College of Mines, Houghton. Illustrated by 103 Original Drawings. 449 pp. 12mo., (1906). . . . \$2.25

**KEMLO.—Watch-Repairer's Hand-Book:**

Being a Complete Guide to the Young Beginner, in Taking Apart, Putting Together, and Thoroughly Cleaning the English Lever and other Foreign Watches, and all American Watches. By F. KEMLO, Tactical Watchmaker. With Illustrations. 12mo. . . . \$1.25

**KENTISH.—A Treatise on a Box of Instruments,**

And the Slide Rule; with the Theory of Trigonometry and Logarithms, including Practical Geometry, Surveying, Measuring of Timber, Cask and Malt Gauging, Heights, and Distances. By THOMAS KENTISH. In one volume. 12mo. . . . \$1.00

**KERL.—The Assayer's Manual:**

An Abridged Treatise on the Docimastic Examination of Ores, and Furnace and other Artificial Products. By BRUNO KERL, Professor in the Royal School of Mines. Translated from the German by WILLIAM T. BRANNT. Second American edition, edited with Extensive Additions by F. LYNWOOD GARRISON, Member of the American Institute of Mining Engineers, etc. Illustrated by 87 engravings. 8vo. (Third Edition in preparation.)

**KICK.—Flour Manufacture.**

A Treatise on Milling Science and Practice. By FREDERICK KICK Imperial Regierungsrath, Professor of Mechanical Technology in the Imperial German Polytechnic Institute, Prague. Translated from the second enlarged and revised edition with supplement by H. H. P. POWLES, Assoc. Memb. Institution of Civil Engineers. Illustrated with 28 Plates, and 167 Wood-cuts. 367 pages. 8vo. . \$10.00

**KINGZETT.—The History, Products, and Processes of the Alkali Trade:**

(including the most Recent Improvements. By CHARLES THOMAS KINGZETT, Consulting Chemist. With 23 illustrations. 8vo. \$2.50

**KIRK.—The Cupola Furnace:**

A Practical Treatise on the Construction and Management of Foundry Cupolas. By EDWARD KIRK, Practical Moulder and Melter, Consulting Expert in Melting. Illustrated by 78 engravings. Second Edition, revised and enlarged. 450 pages. 8vo. 1903. \$3.50

**LANDRIN.—A Treatise on Steel:**

Comprising its Theory, Metallurgy, Properties, Practical Working, and Use. By M. H. C. LANDRIN, JR. From the French, by A. A. FESQUET. 12mo. . . . \$2.50

**LANGBEIN.—A Complete Treatise on the Electro-Deposition of Metals:**

Comprising Electro-Plating and Galvanoplastic Operations, the Deposition of Metals by the Contact and Immersion Processes, the Coloring of Metals, the Methods of Grinding and Polishing, as well as Descriptions of the Electric Elements, Dynamo-Electric Machines, Thermo-Piles and of the Materials and Processes used in Every Department of the Art. From the German of DR. GEORGE LANGBEIN, with additions by WM. T. BRANNT. Fifth Edition, thoroughly revised and much enlarged. 170 Engravings. 694 pages 8vo. 1905. \$4.00

**LARDNER.—The Steam-Engine:**

For the Use of Beginners. Illustrated. 12mo. . . . .60

**LEHNER.—The Manufacture of Ink:**

Comprising the Raw Materials, and the Preparation of Writing, Copying and Hektograph Inks, Safety Inks, Ink Extracts and Powders, etc. Translated from the German of SIGMUND LEHNER, with additions by WILLIAM T. BRANNT. Illustrated. 12mo. \$2.00



**LARKIN.—The Practical Brass and Iron Founder's Guide:**

A Concise Treatise on Brass Founding, Moulding, the Metals and their Alloys, etc.; to which are added Recent Improvements in the Manufacture of Iron, Steel by the Bessemer Process, etc., etc. By JAMES LARKIN, late Conductor of the Brass Foundry Department of Reany, Neafe & Co.'s Penn Works, Philadelphia. New edition, revised, with extensive additions. 414 pages. 12mo. . . . \$2.50

**LEROUX.—A Practical Treatise on the Manufacture of Worsted and Carded Yarns:**

Comprising Practical Mechanics, with Rules and Calculations applied to Spinning; Sorting, Cleaning, and Scouring Wools; the English and French Methods of Combing, Drawing, and Spinning Worsted, and Manufacturing Carded Yarns. Translated from the French of CHARLES LEROUX, Mechanical Engineer and Superintendent of a Spinning-Mill, by HORATIO PAINE, M. D., and A. A. FESQUET, Chemist and Engineer. Illustrated by twelve large Plates. To which is added an Appendix, containing Extracts from the Reports of the International Jury, and of the Artisans selected by the Committee appointed by the Council of the Society of Arts, London, on Woollen and Worsted Machinery and Fabrics, as exhibited in the Paris Universal Exposition, 1867. 8vo. . . . . \$4.00

**LEFFEL.—The Construction of Mill-Dams:**

Comprising also the Building of Race and Reservoir Embankments and Head-Gates, the Measurement of Streams, Gauging of Water Supply, etc. By JAMES LEFFEL & Co. Illustrated by 58 engravings. 8vo. . . . . (Scarce.)

**LESLIE.—Complete Cookery:**

Directions for Cookery in its Various Branches. By MISS LESLIE. Sixtieth thousand. Thoroughly revised, with the addition of New Receipts. 12mo. . . . . \$1.50

**LE VAN.—The Steam Engine and the Indicator:**

Their Origin and Progressive Development; including the Most Recent Examples of Steam and Gas Motors, together with the Indicator, its Principles, its Utility, and its Application. By WILLIAM BARNET LE VAN. Illustrated by 205 Engravings, chiefly of Indicator-Cards. 469 pp. 8vo. . . . . \$2.00

**LIEBER.—Assayer's Guide:**

Or, Practical Directions to Assayers, Miners, and Smelters, for the Tests and Assays, by Heat and by Wet Processes, for the Ores of all the principal Metals, of Gold and Silver Coins and Alloys, and of Coal, etc. By OSCAR M. LIEBER. Revised. 283 pp. 12mo. . . . \$1.50

**Lockwood's Dictionary of Terms:**

Used in the Practice of Mechanical Engineering, embracing those Current in the Drawing Office, Pattern Shop, Foundry, Fitting, Turning, Smith's and Boiler Shops, etc., etc., comprising upwards of Six Thousand Definitions. Edited by a Foreman Pattern Maker, author of "Pattern Making." 417 pp. 12mo. . . . . \$3.75

**LUKIN.—The Lathe and Its Uses :**

Or Instruction in the Art of Turning Wood and Metal. Including a Description of the Most Modern Appliances for the Ornamentation of Plane and Curved Surfaces, an Entirely Novel Form of Lathe for Eccentric and Rose-Engine Turning; A Lathe and Planing Machine Combined; and Other Valuable Matter Relating to the Art. Illustrated by 462 engravings. Seventh edition. 315 pages. 8vo. . . . . \$4.25

**MAIN and BROWN.—Questions on Subjects Connected with the Marine Steam-Engine :**

And Examination Papers; with Hints for their Solution. By THOMAS J. MAIN, Professor of Mathematics, Royal Naval College, and THOMAS BROWN, Chief Engineer, R. N. 12mo., cloth . \$1.00

**MAIN and BROWN.—The Indicator and Dynamometer :**

With their Practical Applications to the Steam-Engine. By THOMAS J. MAIN, M. A. F. R., Ass't S. Professor Royal Naval College, Portsmouth, and THOMAS BROWN, Assoc. Inst. C. E., Chief Engineer R. N., attached to the R. N. College. Illustrated. 8vo. .

**MAIN and BROWN.—The Marine Steam-Engine.**

By THOMAS J. MAIN, F. R. Ass't S. Mathematical Professor at the Royal Naval College, Portsmouth, and THOMAS BROWN, Assoc. Inst. C. E., Chief Engineer R. N. Attached to the Royal Naval College. With numerous illustrations. 8vo. . . . .

**MAKINS.—A Manual of Metallurgy :**

By GEORGE HOGARTH MAKINS. 100 engravings. Second edition rewritten and much enlarged. 12mo., 592 pages . . . .

**MARTIN.—Screw-Cutting Tables, for the Use of Mechanical Engineers :**

Showing the Proper Arrangement of Wheels for Cutting the Threads of Screws of any Required Pitch; with a Table for Making the Universal Gas-Pipe Thread and Taps. By W. A. MARTIN, Engineer. 8vo. . . . . .50

**MICHELL.—Mine Drainage :**

Being a Complete and Practical Treatise on Direct-Acting Under ground Steam Pumping Machinery. With a Description of a large number of the best known Engines, their General Utility and the Special Sphere of their Action, the Mode of their Application, and their Merits compared with other Pumping Machinery. By STEPHEN MICHELL. Illustrated by 247 engravings. 8vo., 369 pages. \$12 50

**MOLESWORTH.—Pocket-Book of Useful Formulæ and Memoranda for Civil and Mechanical Engineers.**

By GUILFORD L. MOLESWORTH, Member of the Institution of Civil Engineers, Chief Resident Engineer of the Ceylon Railway. Full-bound in Pocket-book form . . . . . \$1.00



**MOORE.—The Universal Assistant and the Complete Mechanic:**

Containing over one million Industrial Facts, Calculations, Receipts, Processes, Trades Secrets, Rules, Business Forms, Legal Items, Etc., in every occupation, from the Household to the Manufactory. By R. MOORE. Illustrated by 500 Engravings. 12mo. \$2.50

**MORRIS.—Easy Rules for the Measurement of Earthworks:**

By means of the Prismoidal Formula. Illustrated with Numerous Wood-Cuts, Problems, and Examples, and concluded by an Extensive Table for finding the Solidity in cubic yards from Mean Areas. The whole being adapted for convenient use by Engineers, Surveyors, Contractors, and others needing Correct Measurements of Earthwork. By ELWOOD MORRIS, C. E. 8vo. \$1.50

**MAUCHLINE.—The Mine Foreman's Hand-Book**

Of Practical and Theoretical Information on the Opening, Ventilating, and Working of Collieries. Questions and Answers on Practical and Theoretical Coal Mining. Designed to Assist Students and Others in Passing Examinations for Mine Foremanships. By ROBERT MAUCHLINE. 3d Edition. Thoroughly Revised and Enlarged by F. ERNEST BRACKETT. 134 engravings, 8vo. 378 pages. (1905) \$3.75

**NAPIER.—A System of Chemistry Applied to Dyeing.**

By JAMES NAPIER, F. C. S. A New and Thoroughly Revised Edition. Completely brought up to the present state of the Science, including the Chemistry of Coal Tar Colors, by A. A. FESQUET, Chemist and Engineer. With an Appendix on Dyeing and Calico Printing, as shown at the Universal Exposition, Paris, 1867. Illustrated. 8vo. 422 pages \$2.50

**NEVILLE.—Hydraulic Tables, Coefficients, and Formulæ, for finding the Discharge of Water from Orifices, Notches Weirs, Pipes, and Rivers:**

Third Edition, with Additions, consisting of New Formulæ for the Discharge from Tidal and Flood Sluices and Siphons; general information on Rainfall, Catchment-Basins, Drainage, Sewerage, Water Supply for Towns and Mill Power. By JOHN NEVILLE, C. E. M. R. I. A.; Fellow of the Royal Geological Society of Ireland. Thick 12mo. \$5.50

**NEWBERRY.—Gleanings from Ornamental Art of every style:**

Drawn from Examples in the British, South Kensington, Indian, Crystal Palace, and other Museums, the Exhibitions of 1851 and 1862, and the best English and Foreign works. In a series of 100 exquisitely drawn Plates, containing many hundred examples. By ROBERT NEWBERRY. 4to. (Scarce.)

**NICHOLLS.—The Theoretical and Practical Boiler-Maker and Engineer's Reference Book:**

Containing a variety of Useful Information for Employers of Labor, Foremen and Working Boiler-Makers, Iron, Copper, and Tinsmiths

**Draughtsmen, Engineers, the General Steam-using Public, and for the Use of Science Schools and Classes.** By SAMUEL NICHOLLS. Illustrated by sixteen plates, 12mo. \$2.50

**NICHOLSON.—A Manual of the Art of Bookbinding:**

Containing full instructions in the different Branches of Forwarding, Gilding, and Finishing. Also, the Art of Marbling Book-edges and Paper. By JAMES B. NICHOLSON. Illustrated. 12mo., cloth \$2.25

**NICOLLS.—The Railway Builder:**

A Hand-Book for Estimating the Probable Cost of American Railway Construction and Equipment. By WILLIAM J. NICOLLS, Civil Engineer. Illustrated, full bound, pocket-book form \$2.00

**NORMANDY.—The Commercial Handbook of Chemical Analysis:**

Or Practical Instructions for the Determination of the Intrinsic or Commercial Value of Substances used in Manufactures, in Trades, and in the Arts. By A. NORMANDY. New Edition, Enlarged, and to a great extent rewritten. By HENRY M. NOAD, Ph.D., F.R.S., thick 12mo. Scarce

**NORRIS.—A Handbook for Locomotive Engineers and Machinists:**

Comprising the Proportions and Calculations for Constructing Locomotives; Manner of Setting Valves; Tables of Squares, Cubes, Areas, etc., etc. By SEPTIMUS NORRIS, M. E. New edition. Illustrated, 12mo. \$1.50

**NYSTROM.—A New Treatise on Elements of Mechanics:**

Establishing Strict Precision in the Meaning of Dynamical Terms, accompanied with an Appendix on Duodenal Arithmetic and Metrology. By JOHN W. NYSTROM, C. E. Illustrated. 8vo.

**NYSTROM.—On Technological Education and the Construction of Ships and Screw Propellers:**

For Naval and Marine Engineers. By JOHN W. NYSTROM, late Acting Chief Engineer, U. S. N. Second edition, revised, with additional matter. Illustrated by seven engravings. 12mo. \$1.25

**O'NEILL.—A Dictionary of Dyeing and Calico Printing:**

Containing a brief account of all the Substances and Processes, & use in the Art of Dyeing and Printing Textile Fabrics; with Practical Receipts and Scientific Information. By CHARLES O'NEILL, Analytical Chemist. To which is added an Essay on Coal Tar Colors and their application to Dyeing and Calico Printing. By A. A. FESQUET, Chemist and Engineer. With an appendix on Dyeing and Calico Printing, as shown at the Universal Exposition, Paris, 1867. 8vo., 491 pages \$2.50

**ORTON.—Underground Treasures:**

How and Where to Find Them. A Key for the Ready Determination of all the Useful Minerals within the United States. By JAMES ORTON, A.M., Late Professor of Natural History in Vassar College, N. Y.; author of the "Andes and the Amazon," etc. A New Edition, with An Appendix on Ore Deposits and Testing Minerals (1901). Illustrated \$1.50

**OSBORN.—The Prospector's Field Book and Guide.**

In the Search For and the Easy Determination of Ores and Other Useful Minerals. By Prof. H. S. OSBORN, LL. D. Illustrated by 66 Engravings. Seventh Edition. Revised and Enlarged. 379 pages, 12mo. (March, 1907) . . . . . \$1.50

**OSBORN—A Practical Manual of Minerals, Mines and Mining:**

Comprising the Physical Properties, Geologic Positions, Local Occurrence and Associations of the Useful Minerals; their Methods of Chemical Analysis and Assay; together with Various Systems of Excavating and Timbering, Brick and Masonry Work, during Driving, Lining, Bracing and other Operations, etc. By Prof. H. S. OSBORN, LL. D., Author of "The Prospector's Field-Book and Guide." 171 engravings. Second Edition, revised. 8vo. . . . . \$4.50

**OVERMAN.—The Manufacture of Steel:**

Containing the Practice and Principles of Working and Making Steel. A Handbook for Blacksmiths and Workers in Steel and Iron, Wagon Makers, Die Sinkers, Cutlers, and Manufacturers of Files and Hardware, of Steel and Iron, and for Men of Science and Art. By FREDERICK OVERMAN, Mining Engineer, Author of the "Manufacture of Iron," etc. A new, enlarged, and revised Edition. By A. A. FESQUET, Chemist and Engineer. 12mo. . . . . \$1.50

**OVERMAN.—The Moulder's and Founder's Pocket Guide:**

A Treatise on Moulding and Founding in Green-sand, Dry-sand, Loam, and Cement; the Moulding of Machine Frames, Mill-gear, Hollowware, Ornaments, Trinkets, Bells, and Statues; Description of Moulds for Iron, Bronze, Brass, and other Metals; Plaster of Paris, Sulphur, Wax, etc.; the Construction of Melting Furnaces, the Melting and Founding of Metals; the Composition of Alloys and their Nature, etc., etc. By FREDERICK OVERMAN, M. E. A new Edition, to which is added a Supplement on Statuary and Ornamental Moulding, Ordnance, Malleable Iron Castings, etc. By A. A. FESQUET, Chemist and Engineer. Illustrated by 44 engravings. 12mo. . . . . \$2.00

**PAINTER, GILDER, AND VARNISHER'S COMPANION.**

Comprising the Manufacture and Test of Pigments, the Arts of Painting, Graining, Marbling, Staining, Sign-writing, Varnishing, Glass-staining, and Gilding on Glass; together with Coach Painting and Varnishing, and the Principles of the Harmony and Contrast of Colors. Twenty-seventh Edition. Revised, Enlarged, and in great part Rewritten. By WILLIAM T. BRANNT, Editor of "Varnishes, Lacquers, Printing Inks and Sealing Waxes." Illustrated. 395 pp. 12mo. . . . . \$1.50

**PALLETT.—The Miller's, Millwright's, and Engineer's Guide.**

By HENRY PALLETT. Illustrated. 12mo. . . . . \$2.00

**PERCY.—The Manufacture of Russian Sheet-Iron.**

By JOHN PERCY, M. D., F. R. S. Paper. . . . . 25 cts.

**PERKINS.—Gas and Ventilation:**

Practical Treatise on Gas and Ventilation. Illustrated. 12mo. \$1.25

**PERKINS AND STOWE.—A New Guide to the Sheet-iron and Boiler Plate Roller:**

Containing a Series of Tables showing the Weight of Slabs and Piles to Produce Boiler Plates, and of the Weight of Piles and the Sizes of Bars to produce Sheet-iron; the Thickness of the Bar Gauge in decimals; the Weight per foot, and the Thickness on the Bar or Wire Gauge of the fractional parts of an inch; the Weight per sheet, and the Thickness on the Wire Gauge of Sheet-iron of various dimensions to weigh 112 lbs. per bundle; and the conversion of Short Weight into Long Weight, and Long Weight into Short.

\$1.50

**POSSELT.—Recent Improvements in Textile Machinery Relating to Weaving:**

Giving the Most Modern Points on the Construction of all Kinds of Looms, Warp-ers, Beamers, Slas/ers, Winders, Spoolers, Reeds, Temples, Shuttles, Bobbins, Heddles, Heddle Frames, Pickers, Jacquards, Card Stampers, etc., etc. 600 illus. . . . \$3.00

**POSSELT.—Technology of Textile Design:**

The Most Complete Treatise on the Construction and Application of Weaves for all Textile Fabrics and the Analysis of Cloth. By E.

A. Posselt. 1,500 illustrations. 4to. . . . . \$5.00

**POSSELT.—Textile Calculations:**

A Guide to Calculations Relating to the Manufacture of all Kinds of Yarns and Fabrics, the Analysis of Cloth, Speed, Power and Belt Calculations. By E. A. POSSELT. Illustrated. 4to. . . \$2.00

**REGNAULT.—Elements of Chemistry:**

By M. V. REGNAULT. Translated from the French by T. FORREST BETTON, M. D., and edited, with Notes, by JAMES C. BOOTH, Melter and Refiner U. S. Mint, and WILLIAM L. FABER, Metallurgist and Mining Engineer. Illustrated by nearly 700 wood-engravings. Comprising nearly 1,500 pages. In two volumes, 8vo., cloth . . \$6.00

**RICHARDS.—Aluminium:**

Its History, Occurrence, Properties, Metallurgy and Applications, including its Alloys. By JOSEPH W. RICHARDS, A. C., Chemist and Practical Metallurgist, Member of the Deutsche Chemische Gesellschaft. Illust. Third edition, enlarged and revised (1895) . . \$6.00

**RIFFAULT, VERGNAUD, and TOUSSAINT.—A Practical Treatise on the Manufacture of Colors for Painting:**

Comprising the Origin, Definition, and Classification of Colors; the Treatment of the Raw Materials; the best Formulæ and the Newest Processes for the Preparation of every description of Pigment, and the Necessary Apparatus and Directions for its Use; Dryers; the Testing. Application, and Qualities of Paints, etc., etc. By MM. RIFFAULT, VERGNAUD, and TOUSSAINT. Revised and Edited by M.



**F. MALEPEYRE.** Translated from the French, by A. A. **FESQUET**, Chemist and Engineer. Illustrated by Eighty engravings. In one vol., 8vo., 659 pages . . . . . \$5.00

**ROPER.**—Catechism for Steam Engineers and Electricians: Including the Construction and Management of Steam Engines, Steam Boilers and Electric Plants. By **STEPHEN ROPER**. Twenty-first edition, rewritten and greatly enlarged by E. R. **KELLER** and C. W. **PIKE**. 365 pages. Illustrations. 18mo., tucks, gilt. \$2.00

**ROPER.**—Engineer's Handy Book: Containing Facts, Formulæ, Tables and Questions on Power, its Generation, Transmission and Measurement; Heat, Fuel, and Steam; The Steam Boiler and Accessories; Steam Engines and their Parts; Steam Engine Indicator; Gas and Gasoline Engines; Materials; their Properties and Strength; Together with a Discussion of the Fundamental Experiments in Electricity, and an Explanation of Dynamos, Motors, Batteries, etc., and Rules for Calculating Sizes of Wires. By **STEPHEN ROPER**. 15th edition. Revised and enlarged by E. R. **KELLER**, M. E. and C. W. **PIKE**, B. S. (1899), with numerous illustrations. Pocket-book form. Leather. . . . . \$3.50

**ROPER.**—Hand-Book of Land and Marine Engines: Including the Modelling, Construction, Running, and Management of Land and Marine Engines and Boilers. With illustrations. By **STEPHEN ROPER**, Engineer. Sixth edition. 12mo., tucks, gilt edge. \$3.50

**ROPER.**—Hand-Book of the Locomotive: Including the Construction of Engines and Boilers, and the Construction, Management, and Running of Locomotives. By **STEPHEN ROPER**. Eleventh edition. 18mo., tucks, gilt edge . . . \$2.50

**ROPER.**—Hand-Book of Modern Steam Fire-Engines. With illustrations. By **STEPHEN ROPER**, Engineer. Fourth edition, 12mo., tucks, gilt edge . . . . . \$3.50

**ROPER.**—Questions and Answers for Engineers. This little book contains all the Questions that Engineers will be asked when undergoing an Examination for the purpose of procuring Licenses, and they are so plain that any Engineer or Fireman of ordinary intelligence may commit them to memory in a short time. By **STEPHEN ROPER**, Engineer. Third edition . . . . . \$2.00

**ROPER.**—Use and Abuse of the Steam Boiler. By **STEPHEN ROPER**, Engineer. Eighth edition, with illustrations. 18mo., tucks, gilt edge . . . . . \$2.00

**ROSE.**—The Complete Practical Machinist: Embracing Lathe Work, Vise Work, Drills and Drilling, Taps and Dies, Hardening and Tempering, the Making and Use of Tools, Tool Grinding, Marking out Work, Machine Tools, etc. By **JOSHUA ROSE**. 395 Engravings. Nineteenth Edition, greatly Enlarged with New and Valuable Matter. 12mo., 504 pages. . . . . \$2.50

**ROSE.**—Mechanical Drawing Self-Taught: Comprising Instructions in the Selection and Preparation of Drawing Instruments, Elementary Instruction in Practical Mechanical Draw-

ing, together with Examples in Simple Geometry and Elementary Mechanism, including Screw Threads, Gear Wheels, Mechanical Motions, Engines and Boilers. By JOSHUA ROSE, M. E. Illustrated by 330 engravings. 8vo, 313 pages . . . \$4.00

**ROSE.—The Slide-Valve Practically Explained :**

Embracing simple and complete Practical Demonstrations of the operation of each element in a Slide-valve Movement, and illustrating the effects of Variations in their Proportions by examples carefully selected from the most recent and successful practice. By JOSHUA ROSE, M. E. Illustrated by 35 engravings . . . \$1.00

**ROSS.—The Blowpipe in Chemistry, Mineralogy and Geology :**

Containing all Known Methods of Anhydrous Analysis, many Working Examples, and Instructions for Making Apparatus. By LIEUT.-COLONEL W. A. ROSS, R. A., F. G. S. With 120 Illustrations. 12mo. . . . . \$2.00

**SHAW.—Civil Architecture :**

Being a Complete Theoretical and Practical System of Building, containing the Fundamental Principles of the Art. By EDWARD SHAW, Architect. To which is added a Treatise on Gothic Architecture, etc. By THOMAS W. SILLOWAY and GEORGE M. HARDING, Architects. The whole illustrated by 102 quarto plates finely engraved on copper. Eleventh edition. 4to. . . . . \$6.00

**SHUNK.—A Practical Treatise on Railway Curves and Location, for Young Engineers.**

By W. F. SHUNK, C. E. 12mo. Full bound pocket-book form \$2.00

**SLATER.—The Manual of Colors and Dye Wares.**

By J. W. SLATER. 12mo. . . . . \$3.00

**SLOAN.—American Houses :**

A variety of Original Designs for Rural Buildings. Illustrated by 26 colored engravings, with descriptive references. By SAMUEL SLOAN, Architect. 8vo. . . . . .75

**SLOAN.—Homestead Architecture :**

Containing Forty Designs for Villas, Cottages, and Farm-houses, with Essays on Style, Construction, Landscape Gardening, Furniture, etc., etc. Illustrated by upwards of 200 engravings. By SAMUEL SLOAN, Architect. 8vo. . . . . \$2.50

**SLOANE.—Home Experiments in Science.**

By T. O'CONOR SLOANE, E. M., A. M., Ph. D. Illustrated by 91 engravings. 12mo. . . . . \$1.00

**SMEATON.—Builder's Pocket-Companion :**

Containing the Elements of Building, Surveying, and Architecture; with Practical Rules and Instructions connected with the subject. By A. C. SMEATON, Civil Engineer, etc. 12mo.

**SMITH.—A Manual of Political Economy.**

By E. PESHINE SMITH. A New Edition, to which is added a full Index. 12mo. . . . . \$1.25



**SMITH.—Parks and Pleasure-Grounds:**

Or Practical Notes on Country Residences, Villas, Public Parks, and Gardens. By CHARLES H. J. SMITH, Landscape Gardener and Garden Architect, etc., etc. 12mo. . . . . \$2.00

**SMITH.—The Dyer's Instructor:**

Comprising Practical Instructions in the Art of Dyeing Silk, Cotton, Wool, and Worsted, and Woolen Goods; containing nearly 800 Receipts. To which is added a Treatise on the Art of Padding; and the Printing of Silk Warps, Skeins, and Handkerchiefs, and the various Mordants and Colors for the different styles of such work. By DAVID SMITH, Pattern Dyer. 12mo. . . . . \$1.00

**SMYTH.—A Rudimentary Treatise on Coal and Coal-Mining.**

By WARRINGTON W. SMYTH, M. A., F. R. G., President R. G. S. of Cornwall. Fifth edition, revised and corrected. With numerous illustrations. 12mo. . . . . \$1.40

**SNIVELY.—Tables for Systematic Qualitative Chemical Analysis.**

By JOHN H. SNIVELY, Phr. D. 8vo. . . . . \$1.00

**SNIVELY.—The Elements of Systematic Qualitative Chemical Analysis:**

A Hand-book for Beginners. By JOHN H. SNIVELY, Phr. D. 16mo. . . . . \$2.00

**STOKES.—The Cabinet-Maker and Upholsterer's Companion:**

Comprising the Art of Drawing, as applicable to Cabinet Work; Veneering, Inlaying, and Buhl-Work; the Art of Dyeing and Staining Wood, Ivory, Bone, Tortoise-Shell, etc. Directions for Lacking, Japanning, and Varnishing; to make French Polish, Glues, Cements, and Compositions; with numerous Receipts, useful to work men generally. By STOKES. Illustrated. A New Edition, with an Appendix upon French Polishing, Staining, Imitating, Varnishing, etc., etc. 12mo . . . . . \$1.25

**STRENGTH AND OTHER PROPERTIES OF METALS:**

Reports of Experiments on the Strength and other Properties of Metals for Cannon. With a Description of the Machines for Testing Metals, and of the Classification of Cannon in service. By Officers of the Ordnance Department, U. S. Army. By authority of the Secretary of War. Illustrated by 25 large steel plates. Quarto . . . \$5.00

**SULLIVAN.—Protection to Native Industry.**

By Sir EDWARD SULLIVAN, Baronet, author of "Ten Chapters on Social Reforms." 8vo. . . . . \$1.00

**SHERRATT.—The Elements of Hand-Railing:**

Simplified and Explained in Concise Problems that are Easily Understood. The whole illustrated with Thirty-eight Accurate and Original Plates, Founded on Geometrical Principles, and Showing how to Make Rail Without Centre Joints, Making Better Rail of the Same Material, with Half the Labor, and Showing How to Lay Out Stairs of all Kinds. By R. J. SHERRATT. Folio. . . . . \$2.50

**SYME.—Outlines of an Industrial Science.**

By DAVID SYME. 12mo.

\$2.00

**TABLES SHOWING THE WEIGHT OF ROUND, SQUARE, AND FLAT BAR IRON, STEEL, ETC.,**

By Measurement. Cloth

63

**THALLNER.—Tool-Steel :**

A Concise Handbook on Tool-Steel in General. Its Treatment in the Operations of Forging, Annealing, Hardening, Tempering, etc., and the Appliances Therefor. By OTTO THALLNER, Manager in Chief of the Tool-Steel Works, Bismarckhütte, Germany. From the German by WILLIAM T. BRANNT. Illustrated by 69 engravings. 194 pages. 8vo. 1902.

\$2.00

**TEMPLETON.—The Practical Examiner on Steam and the Steam-Engine :**

With Instructive References relative thereto, arranged for the Use of Engineers, Students, and others. By WILLIAM TEMPLETON, Engineer. 12mo.

\$1.00

**THAUSING.—The Theory and Practice of the Preparation of Malt and the Fabrication of Beer :**

With especial reference to the Vienna Process of Brewing. Elaborated from personal experience by JULIUS E. THAUSING, Professor at the School for Brewers, and at the Agricultural Institute, Mödling, near Vienna. Translated from the German by WILLIAM T. BRANNT, Thoroughly and elaborately edited, with much American matter, and according to the latest and most Scientific Practice, by A. SCHWARZ and DR. A. H. BAUER. Illustrated by 140 Engravings. 8vo., 815 pages

\$10.00

**THOMPSON.—Political Economy. With Especial Reference to the Industrial History of Nations :**

By ROBERT E. THOMPSON, M. A., Professor of Social Science in the University of Pennsylvania. 12mo.

\$1.50

**THOMSON.—Freight Charges Calculator :**

By ANDREW THOMSON, Freight Agent. 24mo.

\$1.25

**TURNER'S (THE) COMPANION :**

Containing Instructions in Concentric, Elliptic, and Eccentric Turning; also various Plates of Chucks, Tools, and Instruments; and Directions for using the Eccentric Cutter, Drill, Vertical Cutter, and Circular Rest; with Patterns and Instructions for working them. 12mo.

\$1.00

**TURNING: Specimens of Fancy Turning Executed on the Hand or Foot-Lathe :**

With Geometric, Oval, and Eccentric Chucks, and Elliptical Cutting Frame. By an Amateur. Illustrated by 30 exquisite Photographs. 4to.

(Scarce.)

**VAILE.—Galvanized-Iron Cornice-Worker's Manual:**

Containing Instructions in Laying out the Different Mitres, and Making Patterns for all kinds of Plain and Circular Work. Also, Tables of Weights, Areas and Circumferences of Circles, and other Matter calculated to Benefit the Trade. By CHARLES A. VAILE. Illustrated by twenty-one plates. 4to. . . . . (Scarce.)

**VILLE.—On Artificial Manures:**

Their Chemical Selection and Scientific Application to Agriculture. A series of Lectures given at the Experimental Farm at Vincennes, during 1867 and 1874-75. By M. GEORGES VILLE. Translated and Edited by WILLIAM CROOKES, F. R. S. Illustrated by thirty-one engravings. 8vo., 450 pages . . . . . \$6.00

**VILLE.—The School of Chemical Manures:**

Or, Elementary Principles in the Use of Fertilizing Agents. From the French of M. GEO. VILLE, by A. A. FESQUET, Chemist and Engineer. With Illustrations. 12mo. . . . . \$1.25

**VOGDES.—The Architect's and Builder's Pocket-Companion and Price-Book:**

Consisting of a Short but Comprehensive Epitome of Decimals, Duodecimals, Geometry and Mensuration; with Tables of United States Measures, Sizes; Weights, Strengths, etc., of Iron, Wood, Stone, Brick, Cement and Concretes, Quantities of Materials in given Sizes and Dimensions of Wood, Brick and Stone; and full and complete Bills of Prices for Carpenter's Work and Painting; also, Rules for Computing and Valuing Brick and Brick Work, Stone Work, Painting, Plastering, with a Vocabulary of Technical Terms, etc. By FRANK W. VOGDES, Architect, Indianapolis, Ind. Enlarged, revised, and corrected. In one volume, 368 pages, full-bound, pocket-book form, gilt edges . . . . . \$2.00  
Cloth . . . . . 1.50

**VAN CLEVE.—The English and American Mechanic:**

Comprising a Collection of Over Three Thousand Receipts, Rules, and Tables, designed for the Use of every Mechanic and Manufacturer. By B. FRANK VAN CLEVE. Illustrated. 500 pp. 12mo. \$2.00

**VAN DER BURG.—School of Painting for the Imitation of Woods and Marbles:**

A Complete, Practical Treatise on the Art and Craft of Graining and Marbling with the Tools and Appliances. 36 plates. Folio, 12 x 20 inches. . . . . \$6.00

**WAHNSCHAFTE.—A Guide to the Scientific Examination of Soils:**

Comprising Select Methods of Mechanical and Chemical Analysis and Physical Investigation. Translated from the German of Dr. F. WAHNSCHAFTE. With additions by WILLIAM T. BRANNT. Illustrated by 25 engravings. 12mo. 177 pages . . . . . \$1.50

**WALTON.—Coal-Mining Described and Illustrated:**

By THOMAS H. WALTON, Mining Engineer. Illustrated by 24 large and elaborate Plates, after Actual Workings and Apparatus. \$2.00

**WARE.—The Sugar Beet.**

Including a History of the Beet Sugar Industry in Europe, Varieties of the Sugar Beet, Examination, Soils, Tillage, Seeds and Sowing Yield and Cost of Cultivation, Harvesting, Transportation, Conservation, Feeding Qualities of the Beet and of the Pulp, etc. By LEWIS S. WARE, C. E., M. E. Illustrated by ninety engravings. 8vo.

\$3.50

**WARN.—The Sheet-Metal Worker's Instructor:**

For Zinc, Sheet-Iron, Copper, and Tin-Plate Workers, etc. Containing a selection of Geometrical Problems; also, Practical and Simple Rules for Describing the various Patterns required in the different branches of the above Trades. By REUBEN H. WARN, Practical Tin-Plate Worker. To which is added an Appendix, containing Instructions for Boiler-Making, Mensuration of Surfaces and Solids, Rules for Calculating the Weights of different Figures of Iron and Steel, Tables of the Weights of Iron, Steel, etc. Illustrated by thirty-two Plates and thirty-seven Wood Engravings. 8vo.

\$2.50

**WARNER.—New Theorems, Tables, and Diagrams, for the Computation of Earth-work:**

Designed for the use of Engineers in Preliminary and Final Estimates of Students in Engineering, and of Contractors and other non-professional Computers. In two parts, with an Appendix. Part I. A Practical Treatise; Part II. A Theoretical Treatise, and the Appendix. Containing Notes to the Rules and Examples of Part I.; Explanations of the Construction of Scales, Tables, and Diagrams, and a Treatise upon Equivalent Square Bases and Equivalent Level Heights. By JOHN WARNER, A. M., Mining and Mechanical Engineer. Illustrated by 14 Plates. 8vo.

\$3.00

**WILSON.—Carpentry and Joinery:**

By JOHN WILSON, Lecturer on Building Construction, Carpentry and Joinery, etc., in the Manchester Technical School. Third Edition, with 65 full-page plates, in flexible cover, oblong. (Scarce.)

**WATSON.—A Manual of the Hand-Lathe:**

Comprising Concise Directions for Working Metals of all kinds, Ivory, Bone, and Precious Woods; Dyeing, Coloring, and French Polishing; Inlaying by Veneers, and various methods practised to produce Elaborate work with Dispatch, and at Small Expense. By EGBERT P. WATSON, Author of "The Modern Practice of American Machinists and Engineers." Illustrated by 78 engravings. \$1.50

**WATSON.—The Modern Practice of American Machinists and Engineers:**

Including the Construction, Application, and Use of Drills, Lathe Tools, Cutters for Boring Cylinders, and Hollow-work generally, with the most Economical Speed for the same; the Results verified by Actual Practice at the Lathe, the Vise, and on the floor. Together



with Workshop Management, Economy of Manufacture, the Steam Engine, Boilers, Gears, Belting, etc., etc. By **EGBERT P. WATSON**. Illustrated by eighty-six engravings. 12mo. . . . \$2.50

**WATT.—The Art of Soap Making :**

A Practical Hand-Book of the Manufacture of Hard and Soft Soaps, Toilet Soaps, etc. Fifth Edition, Revised, to which is added an Appendix on Modern Candle Making. By **ALEXANDER WATT**. Ill. 12mo. . . . \$3.00

**WEATHERLY.—Treatise on the Art of Boiling Sugar, Crystallizing, Lozenge-making, Comfits, Gum Goods,**

And other processes for Confectionery, including Methods for Manufacturing every Description of Raw and Refined Sugar Goods. A New and Enlarged Edition, with an Appendix on Cocoa, Chocolate, Chocolate Confections, etc. 196 pages, 12mo. (1903) . \$1.50

**WILL.—Tables of Qualitative Chemical Analysis :**

With an Introductory Chapter on the Course of Analysis. By Professor **HEINRICH WILL**, of Giessen, Germany. Third American, from the eleventh German edition. Edited by **CHARLES F. HIMES**, Ph. D., Professor of Natural Science, Dickinson College, Carlisle, Pa. 8vo. . . . \$1.50

**WILLIAMS.—On Heat and Steam :**

Embracing New Views of Vaporization, Condensation and Explosion. By **CHARLES WYE WILLIAMS**, A. I. C. E. Illustrated. 8vo. \$2.50

**WILSON.—First Principles of Political Economy :**

With Reference to Statesmanship and the Progress of Civilization. By Professor **W. D. WILSON**, of the Cornell University. A new and revised edition. 12mo. . . . \$1.50

**WILSON.—The Practical Tool-Maker and Designer :**

A Treatise upon the Designing of Tools and Fixtures for Machine Tools and Metal Working Machinery, Comprising Modern Examples of Machines with Fundamental Designs for Tools for the Actual Production of the work; Together with Special Reference to a Set of Tools for Machining the Various Parts of a Bicycle. Illustrated by 189 engravings. 1898. . . . \$2.50

**CONTENTS:** Introductory. Chapter I. Modern Tool Room and Equipment. II. Files, Their Use and Abuse. III. Steel and Tempering. IV. Making Jigs. V. Milling Machine Fixtures. VI. Tools and Fixtures for Screw Machines. VII. Broaching. VIII. Punches and Dies for Cutting and Drop Press. IX. Tools for Hollow-Ware. X. Embossing: Metal, Coin, and Stamped Sheet-Metal Ornaments. XI. Drop Forging. XII. Solid Drawn Shells or Ferrules; Cupping or Cutting, and Drawing; Breaking Down Shells. XIII. Annealing, Pickling, and Cleaning. XIV. Tools for Draw Bench. XV. Cutting and Assembling Pieces by Means of Ratchet Dial Plates at One Operation. XVI. The Header. XVII. Tools for Fox Lathe. XVIII. Suggestions for a Set of Tools for Machining the Various Parts of a Bicycle. XIX. The Plater's Dynamo. XX. Conclusion—With a Few Random Ideas. Appendix. Index.

**WOODS —Compound Locomotives :**

By **ARTHUR TANNATT WOODS**. Second edition, revised and enlarged by **DAVID LEONARD BARNES**, A. M., C. E. 8vo. 330 pp. \$3.00

## HENRY CAREY BAIRD & CO.'S CATALOGUE.

### **WOHLER.—A Hand-Book of Mineral Analysis :**

By F. WÖHLER, Professor of Chemistry in the University of Göttingen. Edited by HENRY B. NASON, Professor of Chemistry in the Rensselaer Polytechnic Institute, Troy, New York. Illustrated. 12mo. \$2.50

### **WORSSAM.—On Mechanical Saws :**

From the Transactions of the Society of Engineers, 1869. By S. W. WORSSAM, JR. Illustrated by eighteen large plates. 8vo. \$1.50

---

## RECENT ADDITIONS.

### **BRANNT.—Varnishes, Lacquers, Printing Inks and Sealing-Waxes :**

Their Raw Materials and their Manufacture, to which is added the Art of Varnishing and Lacquering, including the Preparation of Putties and of Stains for Wood, Ivory, Bone, Horn, and Leather. By WILLIAM T. BRANNT. Illustrated by 39 Engravings, 338 pages. 12mo. \$3.00

### **BRANNT.—The Practical Dry Cleaner, Scourer, and Garment Dyer :**

Comprising Dry or Chemical Cleaning; Purification of Benzine; Removing Stains; Wet Cleaning; Finishing Cleaned Fabrics; Cleaning and Dyeing Furs, Skins, Rugs and Mats; Cleaning and Dyeing Feathers; Bleaching and Dyeing Straw Hats; Cleaning and Dyeing Gloves; Garment Dyeing; Stripping, Analysis of Textile Fabrics. Edited by WILLIAM T. BRANNT, Editor of the "Techno-Chemical Receipt Book." 2nd edition, in great part re-written and much enlarged. Illustrated. 293 pages. 12mo. \$2.50

### **BRANNT.—Petroleum.**

its History, Origin, Occurrence, Production, Physical and Chemical Constitution, Technology, Examination and Uses; Together with the Occurrence and Uses of Natural Gas. Edited chiefly from the German of Prof. Hans Hoefer and Dr. Alexander Veith, by WM. T. BRANNT. Illustrated by 3 Plates and 284 Engravings. 743 pp. 8vo. \$8.50

### **BRANNT.—A Practical Treatise on the Manufacture of Vinegar and Acetates, Cider, and Fruit-Wines :**

Preservation of Fruits and Vegetables by Canning and Evaporation; Preparation of Fruit-Butters, Jellies, Marmalades, Catchups, Pickles, Mustards, etc. Edited from various sources. By WILLIAM T. BRANNT. Illustrated by 79 Engravings. 479 pp. 8vo. \$5.00

### **BRANNT.—The Metal Worker's Handy-Book of Receipts and Processes :**

Being a Collection of Chemical Formulas and Practical Manipulations for the working of all Metals; including the Decoration and Beautifying of Articles Manufactured therefrom, as well as their Preservation. Edited from various sources. By WILLIAM T. BRANNT. Illustrated. 12mo. \$2.50



**DEITE.—A Practical Treatise on the Manufacture of Perfumery:**

Comprising directions for making all kinds of Perfumes, Sachet Powders, Fumigating Materials, Dentifrices, Cosmetics, etc., with a full account of the Volatile Oils, Balsams, Resins, and other Natural and Artificial Perfume-substances, including the Manufacture of Fruit Ethers, and tests of their purity. By Dr. C. DEITE, assisted by L. BORCHERT, F. EICHBAUM, E. KUGLER, H. TOEFFNER, and other experts. From the German, by WM. T. BRANNT. 28 Engravings. 358 pages. 8vo. . . . . \$3.00

**EDWARDS.—American Marine Engineer, Theoretical and Practical:**

With Examples of the latest and most approved American Practice. By EMORY EDWARDS. 85 illustrations. 12mo. . . . . \$2.00

**EDWARDS.—900 Examination Questions and Answers:**

For Engineers and Firemen (Land and Marine) who desire to obtain a United States Government or State License. Pocket-book form, gilt edge . . . . . \$1.50

**FLEMMING.—Practical Tanning:**

A Handbook of Modern Processes, Receipts, and Suggestions for the Treatment of Hides, Skins, and Pelts of Every Description. By Lewis A. Flemming. American Tanner. 472 pp. 8vo. (1903) \$4.00.

**POSSELT.—The Jacquard Machine Analysed and Explained:**

With an Appendix on the Preparation of Jacquard Cards, and Practical Hints to Learners of Jacquard Designing. By E. A. POSSELT. With 230 illustrations and numerous diagrams. 127 pp. 4to. . . . . \$3.00

**POSSELT.—Recent Improvements in Textile Machinery, Part III:**

Processes Required for Converting Wool, Cotton, Silk, from Fibre to Finished Fabric, Covering both Woven and Knit Goods; Construction of the most Modern Improvements in Preparatory Machinery, Carding, Combing, Drawing, and Spinning Machinery, Winding, Warping, Slashing Machinery Looms, Machinery for Knit Goods, Dye Stuffs, Chemicals, Soaps, Latest Improved Accessories Relating to Construction and Equipment of Modern Textile Manufacturing Plants. By E. A. POSSELT. Completely Illustrated. 4to.

\$7.50

**RICH.—Artistic Horse-Shoeing:**

A Practical and Scientific Treatise, giving Improved Methods of Shoeing, with Special Directions for Shaping Shoes to Cure Different Diseases of the Foot, and for the Correction of Faulty Action in Trotters. By GEORGE E. RICH. 62 Illustrations. 153 pages. 12mo . . . . . \$2.00



**RICHARDSON.—Practical Blacksmithing:**

A Collection of Articles Contributed at Different Times by Skilled Workmen to the columns of "The Blacksmith and Wheelwright," and Covering nearly the Whole Range of Blacksmithing, from the Simplest Job of Work to some of the Most Complex Forgings. Compiled and Edited by M. T. RICHARDSON.

Vol. I.	210 Illustrations.	224 pages.	12mo.	.	.	\$1.00
Vol. II.	230 Illustrations.	262 pages.	12mo.	.	.	\$1.00
Vol. III.	390 Illustrations.	307 pages.	12mo.	.	.	\$1.00
Vol. IV.	226 Illustrations.	276 pages.	12mo.	.	.	\$1.00

**RICHARDSON.—The Practical Horseshoer:**

Being a Collection of Articles on Horseshoeing in all its Branches which have appeared from time to time in the columns of "The Blacksmith and Wheelwright," etc. Compiled and edited by M. T.

RICHARDSON. 174 illustrations, . . . . \$1.00

**ROPER.—Instructions and Suggestions for Engineers and Firemen:**

By STEPHEN ROPER, Engineer. 18mo. Morocco . \$2.00

**ROPER.—The Steam Boiler: Its Care and Management:**

By STEPHEN ROPER, Engineer. 12mo., tuck, gilt edges. \$2.00

**ROPER.—The Young Engineer's Own Book:**

Containing an Explanation of the Principle and Theories on which the Steam Engine as a Prime Mover is Based. By STEPHEN ROPER, Engineer. 160 illustrations, 363 pages. 18mo., tuck . \$2.50

**ROSE.—Modern Steam-Engines:**

An Elementary Treatise upon the Steam-Engine, written in Plain language; for Use in the Workshop as well as in the Drawing Office. Giving Full Explanations of the Construction of Modern Steam-Engines: Including Diagrams showing their Actual operation. Together with Complete but Simple Explanations of the operations of Various Kinds of Valves, Valve Motions, and Link Motions, etc., thereby Enabling the Ordinary Engineer to clearly Understand the Principles Involved in their Construction and Use, and to Plot out their Movements upon the Drawing Board. By JOSHUA ROSE, M. E. Illustrated by 422 engravings. Revised. 358 pp. . . \$6.00

**ROSE.—Steam Boilers:**

A Practical Treatise on Boiler Construction and Examination, for the Use of Practical Boiler Makers, Boiler Users, and Inspectors; and embracing in plain figures all the calculations necessary in Designing or Classifying Steam Boilers. By JOSHUA ROSE, M. E. Illustrated by 73 engravings. 250 pages. 8vo. . . \$2.50

**SCHRIBER.—The Complete Carriage and Wagon Painter:**

A Concise Compendium of the Art of Painting Carriages, Wagons, and Sleighs, embracing Full Directions in all the Various Branches, including Lettering, Scrolling, Ornamenting, Striping, Varnishing, and Coloring, with numerous Recipes for Mixing Colors. 73 Illustrations. 177 pp. 12mo. . . \$1.00




br













YB 16934

